



## Morpho-toxicology of chlorpyrifos to prolactin cells of a freshwater catfish, *Heteropneustes fossilis*

Ajai Kumar Srivastav\*, Sanjay Kumar Srivastava, Sarojni Tripathi, Diwakar Mishra and Sunil Kumar Srivastav

Department of the Zoology, University Gorakhpur, 273 009, Gorakhpur, India. \*Author for correspondence. E-mail: ajaiksrivastav@hotmail.com

**ABSTRACT.** In the present study, an organophosphorus compound Coroban (active ingredient chlorpyrifos – E.C. 20%) was used. In short-term exposure the fish were subjected to 0.8 of 96h LC<sub>50</sub> value of chlorpyrifos (1.76 mg L<sup>-1</sup>) for 96h. In long-term exposure the experiment was performed for 28 days by using 0.2 of 96h LC<sub>50</sub> value of chlorpyrifos (0.44 mg L<sup>-1</sup>). Fish were killed on each time intervals from control and experimental (chlorpyrifos) groups after 24, 48, 72, and 96h in short-term exposure and after 7, 14, 21, and 28 days in long-term experiment. Blood samples were collected and sera were analyzed for calcium. Pituitary glands were fixed for histological studies and stained with Herlant tetrachrome and Heidenhain's azan techniques. Short-term exposure of chlorpyrifos caused decrease in the serum calcium levels. No change was noticed in the prolactin cells of chlorpyrifos treated fish. Long-term treatment with chlorpyrifos provoked hypocalcemia. The prolactin cells of treated fish exhibited slight degranulation after 21 days whereas the nuclear volume remained unchanged. After 28 days, the prolactin cells exhibited further degranulation and the nuclear volume recorded an increase. Cytolysis and vacuolization were also visible.

**Keywords:** prolactin cells, serum calcium, organophosphate, chlorpyrifos, fish.

## Morfototoxicologia de clorpirifos para as células de prolactina de bagre de água doce *Heteropneustes fossilis*

**RESUMO.** No estudo presente, o composto organofosforo Coroban (ingrediente ativo clorpirifó – E.C. 20%) foi usado. Na exposição a curto prazo os peixes foram submetido a 0,8 de valor LC<sub>50</sub> de 96h de clorpirifó (1,76 mg L<sup>-1</sup>) durante 96h. Na exposição a longo prazo o experimento foi executado durante 28 dias usando 0,2 de valor LC<sub>50</sub> de 96h de clorpirifos (0,44 mg L<sup>-1</sup>). Os peixes foram mortos a cada intervalo dos grupos controle e experimental (clorpirifos) após 24, 48, 72, e 96h em exposição a curto prazo e após 7, 14, 21, e 28 dias no experimento a longo prazo. As amostras de sangue foram colhidas e o soro foi analisado para cálcio. As glândulas pituitárias foram fixadas para estudos histológicos e colorido por tetracromo de Herlant e por técnicas de azan do Heidenhain. A exposição a curto prazo do clorpirifó diminuiu os níveis de cálcio no soro. Nenhuma mudança foi observada nas células de prolactina nos peixes tratados com clorpirifó. O tratamento a longo prazo com clorpirifó causou hipocalcemia. As células de prolactina dos peixes tratados mostraram uma leve degranulação após 21 dias ao passo que o volume nuclear permaneceu inalterado. Depois de 28 dias, as células de prolactina mostraram mais degranulação e o volume nuclear registrou um aumento. Citólise e vacuolização também eram visíveis.

**Palavras-chave:** células de prolactina, cálcio sérico, organofosfato, clorpirifos, peixes.

### Introduction

Pesticides are being used by the human beings for their benefits - control of insect vectors of diseases and increased yield of many crops. The use of pesticides has caused severe environmental and health hazards to organisms including human beings (ABDOLLAHI et al., 2004; TUZMEN et al., 2008). Persistence of organochlorine pesticides in the environment resulted in the search for less persistent pesticide - the organophosphorus (OP) compounds. The first OP compounds developed were tepp and parathion, followed by malathion.

Now there are scores of OP compounds, some liquids and some solids, mainly in the molecular forms known as phosphorothioates, phosphorodithioates and phosphates.

Chlorpyrifos [O, O-diethyl-O- (3, 5, 6-trichloro-2-pyridil) phosphoro-thioate], is a member of organophosphate class of insecticides that displays broad-spectrum insecticidal activity against a number of important arthropod pests (RACKE, 1993). It is recommended as a most effective insecticide for the control of mustard aphid, *Liapaphis erysimi* (SRIVASTAVA; SRIVASTAVA,

1986). The wide-spread use of chlorpyrifos is probably due to its relatively low mammalian toxicity. The use of chlorpyrifos to area of fish breeding causes toxicological hazards especially in view of its high toxicity to fish (HERZBERG, 1987).

Fish are exposed to aquatic toxicants via the extensive and delicate respiratory surface of the gills and, in sea water, also via drinking (WENDELAAR BONGA, 1997). They are also exposed to these toxicants through feeding. The gill is a multifunctional and complex organ with which fish make intimate contact with the surrounding water. The gills comprise over half the body surface area and only a few microns of delicate gill epithelium separate the internal environment from a continually flowing external environment (WOOD; SOIVIO, 1991). Thus branchial function is very sensitive to environmental toxicants. Hence, for the study of toxicological impacts, fish serves as an excellent bioassay animal and have been widely used for this purpose (BLAXTER; HALLERS-TJABBES, 1992; ESPELID et al., 1996; HOLLIS et al., 1999; KUMAR et al., 2007; PRATAP, 1999; RUGGIERI, 1975; SWARUP et al., 1977; WENDELAAR BONGA, 1997).

Prolactin has a physiological role in certain biological features among higher vertebrates such as stimulation of the pegen crop, stimulation of mammary gland and post-ovulatory corpus luteum in some mammals and induction of water-drive and land-water integumentary changes in the urodeles (MEITES; NICOLL, 1966; VELLANO et al., 1967). However, fish prolactin does not elicit these responses. Prolactin has been considered as hypercalcemic factor in teleosts (FARGHER; McKEOWN, 1989; FLIK et al., 1994; PANG, 1981; SRIVASTAV; SWARUP, 1985; SRIVASTAV et al., 1995a, b; WENDELAAR BONGA; FLIK, 1982; WENDELAAR BONGA et al., 1985). Recently, a role of prolactin in embryonic-stage organogenesis in zebrafish (NGUYEN et al., 2008) and in the nuptial colouration in female fish has been reported (SKOLD et al., 2008). Although, several authors have reported the effects of organophosphate pesticides in fishes (AGBONE et al., 2002; JOHAL et al., 2007; ORUC et al., 2006; PATIL; DAVID, 2008; TILAK; SWARNA KUMARI, 2009; ZHANG et al., 2009), but there exist a single report regarding the effect of these organophosphates on the histological structure of prolactin cells of fish (MISHRA et al., 2008). Thus in the foregoing study we have made an effort to explore further the effects of organophosphate namely chlorpyrifos on the prolactin cells of a fish, *Heteropneustes fossilis*.

## Material and methods

Live specimens of adult freshwater catfish, *Heteropneustes fossilis* (both sexes, body weight 42-53 g; body length 16-20 cm) were collected locally (from Ramgarh lake through a commercial supplier) and acclimatized to the laboratory conditions for 15 days in plastic tanks (50 fishes in each; dimensions 48 x 40 x 22 inch; capacity 125 gallon). The physicochemical characteristics of the tap water were used in the experiment are - photoperiod 11.58-12.38h; temperature  $25.80 \pm 1.80^{\circ}\text{C}$ ; pH  $7.21 \pm 0.06$ ; hardness  $167.32 \pm 5.81 \text{ mg L}^{-1}$  as  $\text{CaCO}_3$ ; electrical conductivity  $306.18 \pm 68.52 \mu\text{mhos cm}^{-1}$ ; dissolved oxygen  $7.78 \pm 0.30 \text{ mg L}^{-1}$  and no free chlorine. During acclimatization the fish were fed daily with wheat flour pellets and ground dried shrimps, 2-3 times per day. Water was renewed daily after cleaning the fecal matter and left overfood. The fish were not fed 24h before and during the experimental period (short- and long-term).

In the present study, an organophosphorus compound Coroban (active ingredient chlorpyrifos-E.C. 20% manufactured by Coromandel Indag Products, Madras) was used. In short-term exposure the fish were subjected to 0.8 of 96h  $\text{LC}_{50}$  value of chlorpyrifos ( $1.76 \text{ mg L}^{-1}$ ) for 96h. In long-term exposure the experiment was performed for 28 days by using 0.2 of 96h  $\text{LC}_{50}$  value of chlorpyrifos ( $0.44 \text{ mg L}^{-1}$ ). Simultaneously, a control group was also used for comparison by using the tap water containing acetone (the volume of acetone used was the same as used for preparing the pesticide solution at different concentration). Fish were kept in groups of 10 in 40 L media (tap water containing acetone for control and pesticide's solution prepared in acetone for experimental). Six fish were killed on each time intervals from control and experimental (chlorpyrifos) groups after 24, 48, 72, and 96h in short-term exposure and after 7, 14, 21, and 28 days in long-term experiment.

Blood samples (approx 1 mL) were collected by sectioning the caudal peduncle. The sera were separated by centrifugation (3000 r.p.m.) and analyzed for calcium (TRINDER, 1960). After collection of blood samples, pituitary glands along with the brain were fixed in aqueous Bouin's fluid and Bouin's-Hollande fixatives for histological studies. Tissues were routinely processed in graded series of alcohols, cleared in xylene and embedded in paraffin wax. Serial sections were cut at  $6 \mu\text{m}$ . The pituitaries were stained with Herlant tetrachrome and Heidenhain's azan techniques.

Nuclear indices (maximal length and maximal width) of the prolactin cells (50 nuclei from each fish, thus 300 nuclei from 6 fish) were taken with

the aid of ocular micrometer and the nuclear volume was calculated as:

$$\text{volume} = 4/3 \pi ab^2$$

where:


‘a’ is the major semiaxis; and

‘b’ is the minor semiaxis. Student’s t test was used to test for significant differences between the control and experimental groups with  $p < 0.05$  being accepted as significant.


## Results and discussion

The serum calcium levels of *H. fossilis* exhibit a decline after 24h following exposure to chlorpyrifos. This decrease continues till the end of the experiment (96h; Figure 1). The serum calcium levels of chronically-exposed fish exhibit a decrease on day 7. Thereafter, the levels continue to fall progressively till the end of the experiment (28 days; Figure 2).


In control fish, the prolactin cells are predominant cell type of rostral pars distalis. The cell boundaries of these cells are indistinct (Figure 3). However, the nuclei are oval to round in shape with dense chromatin granules.



**Figure 1.** Serum calcium levels of short-term chlorpyrifos treated *Heteropneustes fossilis*. Values are mean  $\pm$  S.E. of six specimens. Asterisk indicates significant differences ( $p < 0.05$ ) from control.




**Figure 2.** Serum calcium levels of long-term chlorpyrifos treated *Heteropneustes fossilis*. Values are mean  $\pm$  S.E. of six specimens. Asterisk indicates significant differences ( $p < 0.05$ ) from control.




**Figure 3.** Prolactin cells of control *Heteropneustes fossilis*. Mark indistinct cell boundaries. Herlant tetrachrome X 800.


The prolactin cells of *H. fossilis* exposed to chlorpyrifos exhibit no change in the histological structure throughout the short-term experimentation whereas in long-term treated fish, the prolactin cells show no histological change up to 14 days. After 21 days, these cells exhibit slight degranulation (Figure 4). Following 28 days chlorpyrifos exposure, the prolactin cells exhibit further degranulation and at some places, vacuolization and cytolysis are also observed (Figure 5). The nuclear volume records an increase (Figure 6).



**Figure 4.** Prolactin cells of 21 days chlorpyrifos treated fish showing slight degranulation (arrows). Herlant tetrachrome X 800.



**Figure 5.** Prolactin cells of 28 days chlorpyrifos exposed fish showing degeneration (arrows). Note vacuolization and cytolysis (white broken arrows) at some places. Herlant tetrachrome X 800.



**Figure 6.** Nuclear volume of prolactin cells of long-term chlorpyrifos exposed *Heteropneustes fossilis*. Each value represents mean  $\pm$  S.E. of six specimens. Asterisk indicates significant differences ( $p < 0.05$ ) from control.

In the present study chlorpyrifos exposure caused hypocalcemia in fish *H. fossilis*. This is in agreement with reports of earlier investigators who have observed decreased blood/plasma content of calcium of fish treated with toxicants – aldrin (BANO, 1982; SINGH et al., 1996), malachite green (SRIVASTAVA et al., 1995b), cadmium (GILES, 1984; KUROSHIMA, 1987; LARSSON et al., 1981; MURAMOTO, 1981; PRATAP et al., 1989), propoxur (SINGH et al., 1997) and formothion (SINGH et al., 1997). In contrast, other investigators have observed elevation of plasma calcium levels of fish exposed to pesticides (BANSAL et al., 1979; DALELA et al., 1981; SASTRY; SHARMA, 1978; SHARMA et al., 1982). However, no statistically significant effect on blood/plasma calcium concentrations has been observed in DDT treated flounders *Platichthys flesus* (HAUX, 1979), methoxychlor treated Northern puffer *Sphaeroides maculatus* (EISLER, 1967) and bifenthrin treated rainbow trout, *Oncorhynchus mykiss* (VELISEK et al., 2009).

Prolactin cells of *H. fossilis* exhibit degranulation and increased nuclear volume after chlorpyrifos treatment. The activity of the prolactin cells has been judged by the staining response and nuclear volume. In past, the increased nuclear size/area/volume has been considered as indications for the hyperactivity of various endocrine glands by several investigators (OLIVEREAU; OLIVEREAU, 1982; OLIVEREAU et al., 1986; SRIVASTAV; RANI, 1992a and b). To the best of our knowledge the effect of organophosphate on prolactin cell activity has been studied only by Mishra et al. (2008). The present study is in conformity with studies of Mishra et al. (2008) as they have reported hyperactivity of prolactin cells of catfish after metacid-50 treatment (an organophosphate).

The effect of toxicants on the prolactin cell activity has also been reported after cadmium (JAMES; WIGHAM, 1986) and cypermethrin (MISHRA et al., 2011). The result of Mishra et al. (2011) strengthen this study as they noticed an increased activity of prolactin cells in cypermethrin-treated catfish. The present study derives support from the studies of Thangavel et al. (2005) and Meredith et al. (1999) who have reported elevated levels of prolactin in fish treated with dimecron and DDT, respectively. The observed hyperactivity in the prolactin cells of chlorpyrifos exposed *H. fossilis* may be attributed to the hypocalcemia noticed in the present study after pesticide treatment as prolactin provoke hypercalcemia in various species of fishes (FLIK et al., 1986, 1994; PANG et al., 1978; WENDELAAR BONGA; FLIK, 1982; WENDELAAR BONGA; PANG, 1991).

## Conclusion

It can be concluded from the present study that chlorpyrifos provoked hypocalcemia in the freshwater catfish *H. fossilis* which caused the hyperactivity of prolactin cells. In fishes calcium has been reported to play an important role in growth, reproduction, and several other physiological functions such as muscle contraction, nerve conduction, membrane permeability etc. (WENDELAAR BONGA; FLIK, 1993). Also calcium has been considered as having a direct stimulating effect on sperm motility (AMIN, 1998; BAYNES et al., 1981). In this study we have noticed continued hypocalcemia after chlorpyrifos exposure to the fish, hence it can be concluded that exposed fish may have disturbed physiological functions including reproduction and sperm motility.

## Acknowledgements

The authors are thankful to Indian Council of Agricultural Research, New Delhi for providing financial assistance for this study.

## References

- ABDOLLAHI, M.; MOSTAFALOU, S.; POURNOUR MOHAMMADI, S.; SHADNIA, S. Oxidative stress and cholinesterase inhibition in saliva and plasma of rats following subchronic exposure to malathion. **Comparative Biochemistry and Physiology - Part C**, v. 137, n. 1, p. 29-34, 2004.
- AGBONE, A. O.; OFAJEKWU, P. C.; EZENWAKA, I. S.; ALEGBELEYE, W. O. Acute toxicity of diazinon on Rotifers, *Cyclops*, mosquito larvae and fish. **Journal of Applied Sciences and Environmental Management**, v. 6, n. 1, p. 18-21, 2002.

- AMIN, E. M. Observations on reproduction techniques applicable to the European eels (*Anguilla anguilla* L.). **Cahiers Options Mediterraneennes**, v. 34, n. 3, p. 223-234, 1998.
- BANO, Y. Effect of aldrin on serum and liver constituents of freshwater catfish *Clarias batrachus* (L.). **Proceedings of the Indian Academy of Sciences (Animal Science)**, v. 91, n. 4, p. 27-32, 1982.
- BANSAL, S. K.; VERMA, S. R.; GUPTA, A. K.; DALELA, R. C. Physiological dysfunction of the haemopoietic system in a freshwater teleost *Labeo rohita* following chronic chlordane exposure. Part I. Alteration in certain hematological parameters. **Bulletin of Environmental Contamination and Toxicology**, v. 22, n. 1, p. 666-673, 1979.
- BAYNES, S. M.; SCOTT, A. P.; DAWSON, A. P. Rainbow trout *Salmo gairdneri* R., spermatozoa: effects of cations and pH on motility. **Journal of Fish Biology**, v. 19, n. 3, p. 259-267, 1981.
- BLAXTER, J. H. S.; HALLERS-TJABBES, C. C. T. The effects of pollutants on sensory systems and behaviour of aquatic animals. **Netherlands Journal of Aquatic Ecology**, v. 26, n. 1, p. 43-58, 1992.
- DALELA, R. C.; RANI, S.; KUMAR, V.; VERMA, S. R. In vivo hematological alteration in a freshwater teleost *Mystus vittatus* following subacute exposure to pesticides and their combination. **Journal of Environmental Biology**, v. 2, n. 2, p. 79-86, 1981.
- EISLER, R. U. S. Bureau sport fisheries wildlife. In: EDWARDS, R. P. (Ed.). **Pesticides and ecology**. s/l publishing, 1967. p. 213-251. (Paper 17, p. 15).
- ESPELID, S.; LOKKEN, G. B.; STEIRO, K.; BOGWALD, J. Effects of cortisol and stress on the immune system in Atlantic salmon (*Salmo salar* L.). **Fish and Shellfish Immunology**, v. 6, n. 2, p. 95-110, 1996.
- FARGHER, R. C.; McKEOWN, B. A. The effect of prolactin on calcium homeostasis on coho salmon (*Oncorhynchus kisutch*). **General and Comparative Endocrinology**, v. 73, n. 3, p. 398-403, 1989.
- FLIK, G.; RENTIER-DELRUE, F.; WENDELAAR BONGA, S. E. Calcitropic effects of recombinant prolactins in *Oreochromis mossambicus*. **American Journal of Physiology: Renal Physiology**, v. 266, p. 1302-1308, 1994.
- FLIK, G.; FENWICK, J. C.; KOLAR, Z.; MAYER-GOSTAN, N.; WENDELAAR BONGA, S. E. Effects of ovine prolactin on calcium uptake and distribution in the freshwater cichlid teleost fish, *Oreochromis mossambicus*. **American Journal of Physiology**, v. 250, n. 2, p. 161-166, 1986.
- GILES, M. Electrolyte and water balance in plasma and urine of rainbow trout (*Salmo gairdneri*) during chronic exposure to cadmium. **Canadian Journal of Fisheries and Aquatic Sciences**, v. 41, n. 11, p. 1678-1685, 1984.
- HAUX, C. Effects of DDT on blood plasma electrolytes in the flounder *Platichthys flesus* L. in hypotonic brackish water. **Ambio**, v. 8, n. 4, p. 171-173, 1979.
- HERZBERG, A. M. Toxicity of chlorpyrifos (dursban) in *Oreochromis aureus*, *O. niloticus* and data on its residues in *O. aureus*. **Bamidgeh**, v. 39, p. 13-20, 1987.
- HOLLIS, L.; McGEER, J. C.; McDONALD, D. G.; WOOD, C. M. Cadmium accumulation, gill Cd binding, acclimation, and physiological effects during long term sublethal Cadmium exposure in rainbow trout. **Aquatic Toxicology**, v. 46, n. 2, p. 101-119, 1999.
- JAMES, V. A.; WIGHAM, T. The effects of cadmium on prolactin cell activity and plasma cortisol levels in the rainbow trout (*Salmo gairdneri*). **Aquatic Toxicology**, v. 8, n. 4, p. 273-280, 1986.
- JOHAL, M. S.; SHARMA, M. L.; RAVNEET. Impact of low dose of organophosphate, monocrotophos on the epithelial cells gills of *Cyprinus carpio communis* Linn.-SEM study. **Journal of Environmental Biology**, v. 28, n. 3, p. 663-667, 2007.
- KUMAR, A.; SHARMA, B.; PANDEY, R. S. Preliminary evaluation of the acute toxicity of cypermethrin and lambda-cyhalothrin to *Channa punctatus*. **Bulletin of Environmental Contamination and Toxicology**, v. 79, n. 6, p. 613-616, 2007.
- KUROSHIMA, R. Cadmium accumulation and its effect on calcium metabolism in the *Girella Girella punctata* during a long term exposure. **Nippon Suisan Gakkaishi**, v. 53, n. 3, p. 445-450, 1987.
- LARSSON, A.; BENGTSSON, B. E.; HAUX, C. Disturbed ion balance in flounder *Platichthys flesus* L., exposed to sublethal levels of cadmium. **Aquatic Toxicology**, v. 1, n. 1, p. 19-35, 1981.
- MEITES, J.; NICOLL, C. S. Adenohypophysis: prolactin. **Annual Review of Physiology**, v. 28, n. 17, p. 57-88, 1966.
- MEREDITH, H. O.; RICHMAN, N. H.; COLLIER, J. T.; SEALE, A. P.; RILEY, L. G.; BALL, C. H.; SHIMODA, S. K.; STETSON, M. H.; GRAU, E. G. Pesticide effects on prolactin release from the rostral pars distalis and their effects on growth in the tilapia. In: HENSHEL, D. S.; BLACK, M. C.; HARRASS, M. C. (Ed.). **Environmental toxicology and risk assessment: standardization of biomarkers for endocrine disruption and environmental assessment**. West Conshohocken: American Society for Testing and Materials (ASTM), STP 1364, 1999.
- MISHRA, D.; SRIVASTAV, S. K.; SUZUKI, N.; SRIVASTAV, A. K. Prolactin cells of a teleost *Heteropneustes fossilis* intoxicated with Metacid-50. **International Journal of Biological and Chemical Sciences**, v. 2, n. 3, p. 339-345, 2008.
- MISHRA, D.; RAI, R.; SRIVASTAV, S. K.; SRIVASTAV, A. K. Histological alterations in the prolactin cells of a teleost *Heteropneustes fossilis* after exposure to cypermethrin. **Environmental Toxicology**, v. 26, n. 4, p. 359-363, 2011.
- MURAMOTO, S. Vertebral column damage and decrease of calcium concentrations in fish exposed experimentally to cadmium. **Environmental Pollution (Series A)**, v. 24, n. 1, p. 125-133, 1981.
- NGUYEN, N.; STELLWAG, E. J.; ZHU, Y. Prolactin-dependent modulation of organogenesis in the vertebrate: Recent discoveries in zebrafish. **Comparative Biochemistry and Physiology - Part C: Toxicology and Pharmacology**, v. 148, n. 4, p. 370-380, 2008.



- OLIVEREAU, M.; OLIVEREAU, J. Kinetics of the response of prolactin cells to environmental changes in the eel. **Acta Zoologica**, v. 63, n. 4, p. 239-245, 1982.
- OLIVEREAU, M.; CHAMBOLLE, P.; DUBOURG, P.; OLIVEREAU, J. Cytological and ultrastructural changes in the pituitary pars distalis of the goldfish kept in calcium-free environments. **Biology of the Cell**, v. 57, n. 1, p. 77-88, 1986.
- ORUC, E. O.; UNER, N.; SEVGILER, Y.; USTA, D.; DURMAZ, H. Sublethal effects of organophosphate diazinon on the brain of *Cyprinus carpio*. **Drug and Chemical Toxicology**, v. 29, n. 1, p. 57-67, 2006.
- PANG, P. K. T. Hypercalcemic effects of ovine prolactin on intact killifish *Fundulus heteroclitus*, subjected to different environmental calcium challenges. **General and Comparative Endocrinology**, v. 43, n. 2, p. 252-255, 1981.
- PANG, P. K. T.; SCHREIBMAN, M. P.; BALBONTIN, F.; PANG, R. K. Prolactin and pituitary control of calcium regulation in the killifish, *Fundulus heteroclitus*. **General and Comparative Endocrinology**, v. 36, n. 2, p. 306-316, 1978.
- PATIL, V. K.; DAVID, M. Behaviour and respiratory dysfunction as an index of malathion toxicity in the freshwater fish, *Labeo rohita* (Hamilton). **Turkish Journal of Fisheries Aquatic Sciences**, v. 8, n. 2, p. 233-237, 2008.
- PRATAP, H. B. Effects of dietary cadmium on growth and food conversion in the freshwater cichlid *Oreochromis urolepis*. **Journal of Aquaculture in the Tropics**, v. 14, n. 1, p. 85-91, 1999.
- PRATAP, H. B.; FU, H.; LOCK, R. A. C.; WENDELAAR BONGA, S. E. Effect of water borne and dietary cadmium on plasma ions of the teleost *Oreochromis mossambicus* in relation to water calcium level. **Archives of Environmental Contamination and Toxicology**, v. 18, n. 4, p. 568-575, 1989.
- RACKE, K. D. Environmental fate of chlorpyrifos. **Reviews of Environmental Contamination and Toxicology**, v. 131, n. 1, p. 1-158, 1993.
- RUGGIERI, G. D. Aquatic animals in biomedical research. **Annals of the New York Academy of Sciences**, v. 245, p. 39-54, 1975.
- SASTRY, K. V.; SHARMA, S. K. The effect of endrin on the histopathological changes in the liver of *Channa punctatus*. **Bulletin of Environmental Contamination and Toxicology**, v. 20, n. 1, p. 674-677, 1978.
- SHARMA, M. L.; AGARWAL, V. P.; AWASTHI, A. K.; TYAGI, S. K. Hematological and biochemical characteristics of *Heteropneustes fossilis* under the stress of congo red (diphenyl diszabine pthionic acid). **Toxicology Letters**, v. 14, n. 3-4, p. 237-240, 1982.
- SINGH, N. N.; DAS, V. K.; SINGH, S. Effect of aldrin on carbohydrate, protein and ionic metabolism of a freshwater catfish, *Heteropneustes fossilis*. **Bulletin of Environmental Contamination and Toxicology**, v. 57, n. 2, p. 204-210, 1996.
- SINGH, N. N.; DAS, V. K.; SRIVASTAVA, A. K. Formothion and propoxur induced ionic imbalance and skeletal deformity in a catfish, *Heteropneustes fossilis*. **Journal of Environmental Biology**, v. 18, n. 4, p. 357-363, 1997.
- SKOLD, H. N.; AMUNDSEN, T.; SVENSSON, P. A.; MAYER, I.; BJELVENMARK, J.; FORSGREN, E. Hormonal regulation of female nuptial coloration in a fish. **Hormones and Behavior**, v. 54, p. 549-556, 2008.
- SRIVASTAV, A. K.; RANI, L. Effects of calcitonin administration on the ultimobranchial body and parathyroid glands of the freshwater snake, *Natrix piscator*. **European Archives of Biology**, v. 103, p. 181-185, 1992a.
- SRIVASTAV, A. K.; RANI, L. Ultimobranchial body and parathyroid glands of the freshwater snake *Natrix piscator* in response to vitamin D<sub>3</sub> administration. **Journal of Experimental Zoology**, v. 262, n. 3, p. 255-262, 1992b.
- SRIVASTAV, S. P.; SWARUP, K. Structure and behaviour of the corpuscles of Stannius, the ultimobranchial gland and the prolactin cells in response to prolactin induced hypercalcemia in male catfish *Clarias batrachus* (L.). **Monitore Zoologico Italiano**, v. 19, p. 111-120, 1985.
- SRIVASTAV, A. K.; SINGH, S.; SASAYAMA, Y. Vitamin D<sub>3</sub> induced changes in the prolactin cells of the fish, *Heteropneustes fossilis* reared in artificial freshwater, calcium-rich freshwater or calcium-deficient freshwater. **Journal of Reproductive Biology and Comparative Endocrinology**, v. 7, p. 72-82, 1995a.
- SRIVASTAVA, S. J.; SINGH, N. D.; SRIVASTAVA, A. K.; SINHA, R. Acute toxicity of malachite green and its effect on certain blood parameters of a catfish *Heteropneustes fossilis*. **Aquatic Toxicology**, v. 31, n. 3, p. 241-247, 1995b.
- SRIVASTAVA, S. K.; SRIVASTAVA, A. S. Toxicity of field weathered deposits of certain modern insecticides to *Lipaphis erysimis* (Kalt.) with special reference to the seed yield of mustard. **Pesticides**, v. 20, p. 31-33, 1986.
- SWARUP, K.; LAL, S.; SRIVASTAV, A. K. Evaluation of toxicity of effluent of a fertilizer factory in Gorakhpur to *Puntius sophore* by bioassay. **Proceedings of the National Academy of Sciences**, v. 74, n. 1, p. 247-250, 1977.
- TILAK, K. S.; SWARNA KUMARI, R. Acute toxicity of Nuvan, an organophosphate to freshwater fish *Ctenopharyngodon idella* and its effects on oxygen consumption. **Journal of Environmental Biology**, v. 30, n. 6, p. 1031-1033, 2009.
- THANGAVEL, P.; SUMATHIRAL, K.; KARTHIKEYAN, S.; RAMASWAMY, M. Endocrine response of the freshwater teleost, *Sarotherodon mossambicus* (Peters) to dimecron exposure. **Chemosphere**, v. 61, n. 8, p. 1083-1092, 2005.
- TRINDER, P. Colorimetric microdetermination of calcium in serum. **Analyst**, v. 85, p. 889-894, 1960.
- TUZMEN, N.; CANDAN, N.; KAYA, E.; DEMIRYAS, N. Biochemical effects of chlorpyrifos and deltamethrin on altered antioxidative defense mechanisms and lipid peroxidation in rat liver. **Cell Biochemistry and Function**, v. 26, n. 1, p. 119-124, 2008.
- VELISEK, J.; SVOBODOVA, Z.; PIACKOVA, V. Effects of acute exposure to bifenthrin on some haematological, biochemical and histopathological parameters of rainbow trout *Oncorhynchus mykiss*. **Veterinarni Medicina**, v. 54, n. 3, p. 131-137, 2009.

- VELLANO, C.; PEYROT, A.; MAZZI, V. Effect of prolactin on the pituitothyroid axis, integument and behaviour of the adult male crested newt. **Monitore Zoologico Italiano**, v. 1, n. 1, p. 207-227, 1967.
- WENDELAAR BONGA, S. E. The stress response in fish. **Physiological Reviews**, v. 77, n. 3, p. 591-625, 1997.
- WENDELAAR BONGA, S. E.; FLIK, G. Prolactin and calcium metabolism in a teleost fish *Sarotherodon mossambicus*. In: OGURO, C.; PANG, P. K. T. (Ed.). **Comparative endocrinology of calcium regulation**. Tokyo: Science Society Press, 1982. p. 19-26.
- WENDELAAR BONGA, S. E.; FLIK, G. Calcium regulation in fish. In: LAHILOU, B.; VITIELLO, P. (Ed.). **Aquaculture: fundamental and applied research**. Washington, D.C.: American Geophysical Union, 1993. p. 47-59.
- WENDELAAR BONGA, S. E.; FLIK, G.; LOWIK, C. W.; VAN EYS, C. J. Environmental control of prolactin secretion in the teleost fish *Oreochromis* (formerly *Sarotherodon*) *mossambicus*. **General and Comparative Endocrinology**, v. 57, n. 3, p. 352-359, 1985.
- WENDELAAR BONGA, S. E.; PANG, P. K. T. Control of calcium regulating hormones in the vertebrates: Parathyroid hormone, calcitonin, prolactin and stanniocalcin. **International Review of Cytology**, v. 128, p. 139-213, 1991.
- WOOD, C. M.; SOIVIO, A. Environmental effects on gill function: an introduction. **Physiological Zoology**, v. 64, n. 1, p. 1-3, 1991.
- ZHANG, Z. Y.; YU, X. Y.; WANG, D. L.; YAN, H. J.; LIU, X. J. Acute toxicity to zebrafish of two organophosphates and four pyrethroids and their binary mixtures. **Pest Management Science**, v. 66, n. 1, p. 84-89, 2009.

Received on December 30, 2009.

Accepted on September 8, 2010.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.