

# MONTHLY ACTIVITY REPORT February 2018

# Table of Contents

ABSTRACT	1
MAPPING (HOUSEHOLD CONTAINER MONITORING)	
OVITRAP SURVEILLANCE	4
BG SURVEILLANCE	
RESISTANCE TESTING	22
DENGUE ANTIGEN TEST	24
TRAINING ACTIVITIES	26
PRESENTATIONS	27
APPENDICES	28
APPENDIX I Surveillance Map	
APPENDIX II Resistance Datasheet	
APPENDIX III Resistance Protocol	30
APPENDIX IV Dengue Test Graphics	
APPENDIX V Consent Form	32
APPENDIX VI Glossary	34

### **ABSTRACT**

The Belize Vector and Ecology Center has embarked on vector surveillance initiatives that will allow us to determine changes in distribution and density of vectors and obtain information about population changes. In turn this will allows us to monitor and evaluate adequate control methods. In this report, we have outlined the various activities that we have been conducting for the month of February.

The **Household Container Mapping** is being conducted by teams consisting of BVEC employees and Vector Control representatives. So far we have mapped approximately 19.19% of the total structures in Orange Walk Town. Mapping data was updated for the month of February based on the amount of structures we did.

**Ovitrap Surveillance** is being done in which 120 oviposition cups are set out to 60 pre-consented homes in Orange Walk Town. The hatch rate has been determined to be about 40 to 60% and there is interesting variation in the number of eggs collected per zone which may also be related to the position of ovicups and climatic conditions.

The **BG Sentinel Surveillance** is being conducted in order to collect live adult *Aedes* mosquitoes in the field. There are 12 BG Sentinels in the town at 12 pre-consented homes. The traps so far have collected a number of *Aedes* mosquitoes as well as some from the *Culex* species and other types of flies.

The adult *Aedes* mosquitoes collected in the traps are then sorted, identified and stored for testing using the **Dengue Antigen Kit**. Tests conducted so far have turned up negative which is a good sign and will continue testing while working on a pooling strategy depending on the number of adult females collected in the field.

**Larval Resistance Testing** is another key component to the surveillance which will be done in conjunction with the ovitrap surveillance initiative. Currently eggs are being pooled and hatched so that there is enough larvae to meet the standard testing methods.

BVEC also does **Presentations** and **Training** sessions in which we collaborate with schools and other groups to foster community engagement.

### MAPPING (HOUSEHOLD CONTAINER MONITORING)

The MOH OW Vector Control Department has been undergoing an initiative in which we map the structures within Orange Walk Town by acquiring GPS data points and gathering structure information (i.e. level of structure, type, no. of doors, windows etc). This data is being linked to the daily *Aedes aegypti* yard inspection that the MOH is collecting. All the data is being collected with the use of an online form that is displayed on a tablet and utilized by BVEC research assistants. Data collection is done in teams with the presence of at least one vector control representative. Each structure is coded with a unique structure ID that is generated based on the zone, street name and number of the structure. The zonation is obtained from a previous database that the town council used in order to facilitate proper sanitation of the town. Orange Walk town is divided into 6 Zones labelled from A to F. The Ministry of Health uses these same zones so that each Vector Control representative has their designated zones that they are assigned to. New container data as well as the rate of positivity per container type was established. Below are data collected so far that has been analysed up to the month of February 2018.

Aedes Postive Containers	Zone A	ZoneB	Zone C	Zone D	Zone E	Zone F	Total
Vat/tank	3	0	0	0	0	1	4
Drum	11	2	1	0	4	5	23
Vases	2	0	0	0	0	0	2
Bucket	6	2	3	0	5	1	17
Tires	6	3	1	0	15	2	27
Bottle/cans	0	0	0	0	0	0	0
Coconut Shells	0	0	0	0	0	0	0
Treeholes/wells	1	0	0	0	0	0	1
Watering Pans	0	0	0	0	0	0	0
Other	7	6	5	0	11	3	32
Grand Total	36	13	10	0	35	12	106

Table 1. Aedes positive containers per zone

Common Containers Types	Zone A	Zone B	Zone C	Zone D	Zone E	Zone F	Total
Vat/tank	24	21	8	0	16	9	78
Drum	43	8	6	0	15	16	88
Vase	6	0	0	0	0	0	6
Bucket	91	57	25	0	82	47	302
Tires	52	10	4	0	61	13	140
Bottle/cans	180	172	142	0	260	91	845
Coconut Shells	5	0	0	0	0	0	5
Treeholes/wells	11	0	0	0	2	8	21
Watering Pans	39	0	0	0	0	0	39
Other	214	171	70	0	164	188	807
Grand Total	665	439	255	0	600	372	2331

Table 2. Number of different types of containers found per zone

Structures mapped	Total
House	922
House/store	36
Store	13
Apartment/Hotel	10
Lab	0
Church	9
Bar	2
House/bar	2
Restaurant	1
School	4
Cemetery	1
Office	3
Clinic	1
Business	13
Other	5
TOTAL STRUCTURES	5420
TOTAL MAPPED	1040
PERCENTAGE COMPLETED	19.19%

Table 3. Types of structures mapped along with percentage completion

Container Types	Rate of Positivity
Vat/tank	5.1%
Drum	26.1%
Vases	33.3%
Bucket	5.6%
Tires	19.3%
Bottle/cans	0.0%
Coconut Shells	0.0%
Treeholes/wells	4.8%
Watering Pans	0.0%
Other	4.0%

Table 4. Rate of positivity per container based on total positive over total encountered at the yard.

Note: Map displaying GPS points of structures mapped as of February 2018 is attached in the appendices on Page 28

### **OVITRAP SURVEILLANCE**

The Belize Vector and Ecology Center is currently undergoing an ovitrap/ovicup surveillance program for six (6) months in Orange Walk Town to capture the eggs of *Ae. aegypti* and *Ae. albopictus*, key vectors in the transmission of Dengue, Zika, and Chikungunya. Homes were selected via a random selection process generated by QGIS software. A total of sixty (60) homes were selected throughout the six (6) zones in Orange Walk Town. Two (2) ovicups/ovitraps were placed in each home, giving a total of 120 ovicups/ovitraps placed throughout the entire town. Note that informed consent was obtained from all home/property owners to participate in the study under the supervision of a Ministry of Health (MoH) Vector Control Officer.

Ovitraps/ovicups are left in the field for a total of seven (7) days, after which the egg papers inside every ovitrap/ovicup is retrieved and taken to the BVEC insectary. Egg papers are carefully folded and placed in a plastic container, egg papers from each home remain separate from one another. The ovitraps/ovicups are cleaned with a scourging sponge and a new labelled egg paper and fresh water are added. Once in the BVEC insectary, the freshly collected egg papers are hung for a 24 hour period to remove any excess water. Once dry, the eggs are counted and the eggs are stored. Note that egg hatching for each zone is being done by pooling eggs from previous weeks, mainly to ensure that there are enough eggs to yield the number of larvae needed to perform a larvicide resistance test. Below is an actual picture of an Ovicup with egg paper in the field.



Figure 1. Oviposition cup with egg paper inside

Note: A copy of the consent form that was used is attached in the appendices on Page 32

When hatching eggs, egg papers for a particular zone are carefully inspected for eggs, and then segments of the egg paper containing the eggs are cut and placed in a container with water. The container with eggs is then placed in the vacuum hatcher for hatching. Use of the vacuum hatcher ensures a more synchronized hatch and same size larvae. After a few days, third and early fourth stage larvae are sorted for a larvicide resistance test.

*Table 5* below shows the number of eggs collected for each zone for every week. Note that each zone is different in terms of geographic location within the town and whether it is more of a commercial or housing area. The different weather patterns experienced each week are also an important factor influencing oviposition each week. For this reason, the egg density will vary across zones and time.

ZONE	WEEK 17	WEEK 18	WEEK 19	WEEK 20	TOTAL
Α	371	317	72	284	1044
В	221	143	25	248	637
С	267	257	302	150	976
D	156	390	402	340	1288
E	84	776	360	1263	2483
F	131	53	62	758	1004
TOTAL	1230	1936	1223	3043	7432

Table 5. Number of eggs collected in each zone per week of surveillance for a total of four (4) weeks surveillance period. Zone E has the highest number of eggs for the entire surveillance period of four (4) weeks with n=2483. Zone B has the lowest number of eggs for the entire surveillance period with n=637.




Figure 2. Number of eggs collected per week (4 weeks) for each zone.




Figure 3. Number of eggs collected in each zone for the entire surveillance period of four (4) weeks/1 month.




Figure 4. Total cumulative number of eggs collected in each zone for the entire surveillance period (Weeks 1-20).

#### **ZONE A OVITRAPS**




Figure 5. Number of eggs collected in Zone A and temperature data for weeks 17 – 20 of surveillance. Week 17 has the highest number of eggs collected (n=371) and Week 19 has the lowest number of eggs collected (n=72).




Figure 6. Number of eggs collected in Zone A and rainfall data for weeks 17 – 20 of surveillance.




Figure 7. Number of eggs collected per house in Zone A for weeks 17 – 20 of surveillance. House 5 and 6 (OVI-H5 and OVI-H6) have the lowest number of eggs (n=0) and House 9 (OVI-H9) has the highest number of eggs (n=373).

### **ZONE B OVITRAPS**




Figure 8. Number of eggs collected in Zone B and temperature data for weeks 17 – 20 of surveillance. Week 20 has the highest number of eggs collected (n=248) and Week 19 has the lowest number of eggs collected (n=25).




Figure 9. Number of eggs collected in Zone B and rainfall data for weeks 17 – 20 of surveillance.




Figure 10. Number of eggs collected per house in Zone B for weeks 17 – 20 of surveillance. House 12 (OVI-H12) has the lowest number of eggs (n=0) and House 15 (OVI-H15) has the highest number of eggs (n=144).

### **ZONE C - OVITRAPS**




Figure 11. Number of eggs collected in Zone C and temperature data for weeks 17 - 20 of surveillance. Week 19 has the highest number of eggs collected (n=302) and Week 20 has the lowest number of eggs collected (n=150).




Figure 12. Number of eggs collected in Zone C and rainfall data for weeks 17-20 of surveillance.




Figure 13. Number of eggs collected per house in Zone C for weeks 17 – 20 of surveillance. House 27, 29 and 30 (OVI-H27, OVI-H29, OVI-H30) have the lowest number of eggs (n=0) and House 24 (OVI-H24) has the highest number of eggs (n=400).

# **ZONE D - OVITRAPS**




Figure 14. Number of eggs collected in Zone D and temperature data for weeks 17 – 20 of surveillance. Week 19 has the highest number of eggs collected (n=402) and Week 17 has the lowest number of eggs collected (n=156).




Figure 15. Number of eggs collected in Zone D and rainfall data for weeks 17 - 20 of surveillance.




Figure 16. Number of eggs collected per house in Zone D for weeks 17 – 20 of surveillance. House 34 (OVI-H34) has the lowest number of eggs (n=1) and House 40 (OVI-H40) has the highest number of eggs (n=779).

### **ZONE E - OVITRAPS**




Figure 17. Number of eggs collected in Zone E and temperature data for weeks 17 – 20 of surveillance. Week 20 has the highest number of eggs collected (n=1263) and Week 17 has the lowest number of eggs collected (n=84).




Figure 18. Number of eggs collected in Zone E and rainfall data for weeks 17-20 of surveillance.




Figure 19. Number of eggs collected per house in Zone E for weeks 17 – 20 of surveillance. House 44, 45, 46 and 48 (OVI-H44, OVI-H45, OVI-H46 and OVI-H48) has the lowest number of eggs (n=0) and House 49 (OVI-H49) has the highest number of eggs (n=1941).

### **ZONE F - OVITRAPS**




Figure 20. Number of eggs collected in Zone F and temperature data for weeks 17 – 20 of surveillance. Week 20 has the highest number of eggs collected (n=758) and Week 18 has the lowest number of eggs collected (n=53).




Figure 21. Number of eggs collected in Zone F and rainfall data for weeks 17-20 of surveillance.




Figure 22. Number of eggs collected per house in Zone F for weeks 17 – 20 of surveillance. House 55, 58 and 60 (OVI-H55, OVI-H58, and OVI-H60) have the lowest number of eggs (n=0) and House 52 (OVI-H52) has the highest number of eggs (n=711).

## **BG SURVEILLANCE**

For this particular surveillance activity, Biogent (BG) Sentinel traps are used to collect adult *Ae. aegypti* and *Ae. albopictus* mosquitoes in the field. A total of 12 homes (2 homes per zone) were selected via a random selection process generated by QGIS software. Informed consent was obtained from all home/property owners with the supervision of a MOH Vector Control representative.

BGS traps are set out in the field once a week. Note that traps are placed in the mornings and left to operate for 7-8 hours, and then retrieved in the afternoon of the same day. Below is an image of a BG Sentinel taken from one of the houses that is participating in the surveillance initiative.



Figure 23. BG Sentinel trap setup in the field.

The mosquitoes along with other insect specimens collected are taken to the BVEC laboratory and stored for later identification. Identification is done once a week. *Aedes Spp.* and *Culex Spp.* are stored, while any other insect specimens are discarded. Once mosquitoes are identified, they are stored back in the freezer until they can be processed for testing.

	Total Adult Aedes (♂,♀) Collected Per Week						
		Week 17	Week 18	Week 19	Week 20		
ZONE	HOUSE CODE	02-Feb-18	09-Feb-18	16-Feb-18	23-Feb-18	TOTAL	
Α	BG-H1	5	2	1	0	8	
Α	BG-H2	1	0	0	0	1	
Т	OTAL	6	2	1	0	9	
В	BG-H3	0	18	3	9	30	
В	BG-H4	2	5	1	2	10	
Т	OTAL	2	23	4	11	40	
С	BG-H5	3	4	2	2	11	
С	BG-H6	4	0	0	6	10	
Т	OTAL	7	4	2	8	21	
D	BG-H7	4	0	5	2	11	
D	BG-H8	0	0	0	0	0	
Т	OTAL	4	0	5	2	11	
E	BG-H9	0	1	1	5	7	
E	BG-H10	0	1	0	0	1	
Т	OTAL	0	2	1	5	8	
F	BG-H11	0	0	0	0	0	
F	BG-H12	0	0	0	0	0	
Т	OTAL	0	0	0	0	0	
GRAN	ND TOTAL	19	31	13	26	89	

Table 6. Number of adult Aedes mosquitoes, both male and female, captured per house in each zone for each week of surveillance. Zone C has the highest number (n=40) of adult Aedes mosquitoes captured while Zone F has the lowest number of adult Aedes mosquitoes captured (n=0). In terms of weekly captures, week 18 had the highest number (n=31) of mosquitoes captured while week 19 had the lowest number of mosquitoes captured (n=13).




Figure 24. Number of adult Aedes mosquitoes captured per house for four (4) weeks pf surveillance. House 3 (BG-H3) has the highest number of mosquitoes captured (n=30) and Houses 8, 11, and 12 (BG-H8, BG-H11, BG-H12) have the lowest number with no mosquitoes captured during the four (4) weeks of BGS trap surveillance.




Figure 25. Number of adult Aedes mosquitoes captured per house in all zones for each month (4 weeks) of BG Sentinel surveillance thus far.




Figure 26. Number of adult Aedes mosquitoes captured per week of BGS trap surveillance. Includes both male and female Aedes mosquitoes captured.




Figure 17. Number of adult Aedes mosquitoes captured per house in all zones for week 17 of BG Sentinel surveillance. House 1 (BG-H1) has the highest number of adult Aedes mosquitoes captured for that day (n=5).




Figure 28. Number of adult Aedes mosquitoes captured per house in all zones for week 18 of BG Sentinel surveillance. House 3 (BG-H3) has the highest number of adult Aedes mosquitoes captured for that day (n=18).




Figure 29. Number of adult Aedes mosquitoes captured per house in all zones for week 19 of BG Sentinel surveillance. House 7 (BG-H7) has the highest number of adult Aedes mosquitoes captured for that day (n=5).




Figure 30. Number of adult Aedes mosquitoes captured per house in all zones for week 20 of BG Sentinel surveillance. House 3 (BG-H3) has the highest number of adult Aedes mosquitoes captured for that day (n=9).

### **RESISTANCE TESTING**

Insecticide resistance testing is being done currently on mosquito larvae that has been reared up from *Aedes* eggs collected via oviposition cups. The following procedure is being conducted based on the World Health Organization standard protocol for determining the susceptibility of mosquito larvae to insecticides. Third and fourth instar larvae are being tested against liquid Temefos (in a solution of 70% ethanol) at varying concentrations. Temefos is the active ingredient found in the granular Abate being used by the Ministry of Health to treat positive larval habitats.

Larvae are reared up in the BVEC insectary based on the zone that they were collected from. After reaching the appropriate size, the larvae are then prepped for testing by separating them into groups of 20 per container. For each trial, 5 containers are used (2 'controls' and 3 'treatments') per concentration per zone which equals to 100 larvae per trial. Containers being used for the test are small glass bowls. Ideally we want to obtain a total of 5 trials per concentration per zone. Depending on the amount of larvae obtained, that will determined how much trials can be done with varying concentrations at a certain time. The 5 glass bowl replicates, are labelled appropriately and are filled with 225ml distilled water. Using a micropipette, 1ml of 70% ethanol is added to both 'control' containers and 1ml of 'x-mg/l Temefos concentration solution' is are added to the three 'treatment' containers.

The contents of each container are stirred with a glass stirring rod for 30 seconds to ensure proper mixing. The larvae are then introduced into the containers and the investigator records time introduced, room temperature and relative humidity. These procedures are then repeated for every other trial. The investigator will then check for mortality after 24 hours and after 48 hours while recording any temperature and humidity changes. At the end of the experiment, larvae that have been exposed to the Temefos are then discarded in a large liquid waste disposal container while the control larvae reared up to adulthood and identified thereafter to verify species as either *Aedes aegypti* or *Aedes albopictus*.

The baseline data for Zones B and D were established for this month. On the following pages are graphs generated from the QCal software, which we use to calculate the models. Note that specimens from each Zone are tested against a control (ethanol) and 4 different concentrations of Temefos solution (0.005mg/L, 0.025mg/L, 0.125mg/L, and 0.625 mg/L)

# **ZONE B FEB 14, 2018**




Figure 31. Larvae Resistance Curve for Zone B. LD50 concentration is at 0.007 mg/L while LD90 concentration is at 0.0139 mg/L

# **ZONE D FEB 13, 2018**




Figure 32. Larvae Resistance Curve for Zone D. LD50 concentration is at 0.0048 mg/L while LD90 concentration is at 0.0116 mg/L

Note: Sample datasheet used to record resistance data and model used to conduct testing are attached in the appendices on Pages 29-30

### **DENGUE ANTIGEN TEST**

The adult female *Ae. aegypti* and *Ae. albopictus* mosquitoes that are collected from the BG Sentinel traps in the field are used in a Dengue Antigen Test to detect the presence of Dengue (DENV 1-4) in potentially infected Aedes mosquitoes. The VectorTest® Dengue Assay is a rapid test that identifies the presence or absence of antigens to any of the four Dengue serotypes in mosquitoes. The principle behind the VectorTest® Dengue Assay is the dual monoclonal antibody "sandwich" principle, similar to a pregnancy test, where there is a control zone that will always develop and a test zone that will only develop if there is antigen presence in the mosquito solution.

Prior to processing, all identified Aedes mosquitoes are stored at  $-20^{\circ}$ C. At no point in time are mosquito samples taken out of the fridge unless for processing. For samples of 25 mosquitoes or less, conical grinding tubes/Eppendorf tubes are used for grinding of mosquitoes. For samples of 26-50 mosquitoes, plastic culture tubes are used for the grinding of mosquitoes. Once mosquitoes are ground, test strips can be placed in the mosquito suspension and left to react for 20-30 minutes, after which the test results can be read and recorded. All dip sticks with test results are placed in a graphics template and a picture is taken for recording purposes.

	Female Adul	t Aedes for Week	cs 17-20 used in Vecto	orTest®		
		Date of Test: N	/Jarch 9, 2018			
ZONE	Ae. aegypti	Ae. albopictus	Aedes Spp. (un-ID)	Total	# of Samples Run	# of Positive Samples
Α	7	0	0	7	2	0
В	26	0	0	26	2	0
С	8	0	2	10	3	0
D	5	0	1	6	2	0
E	3	0	0	3	2	0
F	0	0	0	0	0	0
TOTAL	49	0	3	52	11	

Table 7. Number of female adult Aedes mosquitoes per zone, sorted by species, used in Dengue Antigen Detection Test.

	Surveillance Period: Weeks 17 - 20								
	DATE OF TEST: March 9, 2018								
ВС	SENTINEL I		Number o	Number of Female Aedes Adult Mosquitoes Tested in VectorTest®					
ZONE	HOUSE CODE	BG SENTINEL NO.	Ae. aegypti	Ae. albopictus	Aedes Spp (un- ID).	Total/House			
Α	BG-H1	1	6	0	0	6			
Α	BG-H2	2	1	0	0	1			
В	BG-H3	3	18	0	0	18			
В	BG-H4	4	8	0	0	8			
С	BG-H5	5	3	0	2	5			
С	BG-H6	6	5	0	0	5			
D	BG-H7	7	5	0	1	6			
D	BG-H8	8	0	0	0	0			
E	BG-H9	9	2	0	0	2			
E	BG-H10	10	1	0	0	1			
F	BG-H11	11	0	0	0	0			
F	BG-H12	12	0	0	0	0			

Table 8. Number of female adult Aedes mosquitoes per house, sorted by species, used in Dengue Antigen Detection Test.

NOTE: Diagram with test results on the dipsticks is attached in Appendices on Page 31

# TRAINING ACTIVITIES

 $Training\ sessions\ are\ conducted\ by\ BVEC\ staff\ and\ International\ partners\ to\ collaborators\ that\ work\ and\ conduct\ research\ in\ Country.\ No\ Training\ activities\ we\ conducted\ in\ the\ month\ of\ February\ 2018$ 

### **PRESENTATIONS**

Presentations are given by BVEC staff and International partners to collaborators as well as local partners to promote community engagement. Below are a few presentations that have been given so far.

# • Belmopan Comprehensive High School Biology Class Presentation


o Date: February 2018

o Venue: Belmopan Comprehensive High School

o Purpose: Education on Common Vectors and Vector Borne Diseases in Belize

o Presenter/s: Donovan Leiva

# **APPENDICES**



Appendix 1. Map of Orange Walk Town divided into zones and displaying points of structures mapped

# APPENDIX II Resistance Datasheet


### Small-scale field testing and evaluation of larvicides against mosquito larvae

Experiment N	To: <u>2</u>	Starting date:	8/29/17	Locati	on: Orange	<u> Walk Town</u>
Investigator:		Assessm	ent date: _		Pre or day	s posttreatment:
-	Types of Habit	at:	Species	Aedes ase	nati	_

Treatment	C1-	Zone: C		Zone: D	Zone: D		Zone:	
(Temephos)	Sample	24 hrs.	48 hrs.	24 hrs.	48 hrs.	24 hrs.	48 hrs.	
Control	1 10	L 10	L 9/P1	L 10	L 10			
	2 10	L 10	L 10	L 10	L 10			
	3 10	L 10	L 10	L 10	L 10			
	4 10	L 10	L 10	L 10	L 10			
	5							
	Total 40	40	19	40	40			
	Mean 10	10	9.5	10	10			
	%red							
T1 0.005mg/l	1 10	L 10	L 8/P1	L 10	L 9			
	2 10	L 10	L9	L 10	L 10			
	3 10	L 10	L9	L 10	L 7			
	4 10	L 10	L 10	L 10	L8			
	5							
	Total 40	40	36	40	34			
	Mean 10	10	9	10	8.5			
	%red							
T2 0.025mg/l	1 10	L 5	L 2	L 6	L 5			
	2 10	L 6	L 6	L 5	L 5			
	3 10	L 3	L 2	L 2	L 8			
	4 10	L 6	L 1	L 7	L 2			
	5							
	Total 40	20	11	20	20			
	Mean 10	5	2.75	5	5			
	%red							


Appendix 2. Sample Datasheet showing results obtained from a small scale resistance test on Zone C and Zone D in Orange Walk Town. Two concentrations of Temephos were used (0.005mg/L and 0.025mg/L)

# APPENDIX III Resistance Protocol



Appendix 3. Model outlining resistance testing protocol

# APPENDIX IV Dengue Test Graphics



Appendix 4. Diagram of Vector Test dipsticks displaying results post testing.

## APPENDIX V Consent Form

# **Consent for Participation in Research**

**Title:** Ovicup/Ovitrap Surveillance for *Ae. aegypti* and *Ae. albopictus* eggs **Principal Investigator(s):** The Belize Vector and Ecology Center (BVEC)

#### Introduction

The purpose of this form is to provide you with information that may affect your decision as to whether or not to participate in this research study. Read the information below and ask any questions that you might have before deciding whether or not to take part. The person(s) performing this study will answer any of your questions. If you decide to be involved in this study, this form will be used to record your consent.

#### **Purpose of the Study**

You have been asked to participate in this study by having ovicups/ovitraps set out in your yard for capturing *Ae. aegypti* and *Ae. albopictus* mosquito eggs. The eggs of *Ae. aegypti* and/or *Ae. albopictus* collected in this study will be collected from your yard and hatched in the BVEC laboratory to evaluate mosquito larvicide resistance that may impair the effectiveness of larvicide applications currently being used by the Belize Ministry of Health. This study will also help us to better understand the density of mosquito vectors in the area as well as the abundance of female *Ae. aegypti* and *Ae. albopictus*.

### What will you be asked to do?

If you agree to participate in this study, you will be asked to provide us with access to your yard so that two ovicups/ovitraps can be placed outside and then retrieved after 7 days of installation. Therefore, ovicups/ovitraps will be out in your yard for an entire week to collect maximum amount of *Ae. aegypti* and/or *Ae. albopictus* eggs. You and individuals at your property will be asked to not move the ovicups/ovitraps from their location and/or disturb them so that the water in the cup or egg paper is not lost. The ovicups/ovitraps will remain in your yard for a total of 6 months. If the study is determined to need to be extended more than 6 months, we will ask if you would like to continue participating.

#### What are the risks involved in this study?

The risks involved with participation in this study are low and may include invasion of privacy. Risk level however, will be reduced by attempting to identify times of the day for ovicup/ovitrap placement and egg paper collection that fit homeowner habits (i.e., time of waking, time of preparing meals, returning from work).

#### How will your privacy and confidentiality be protected if you participate in this research study?

Your privacy and the confidentiality of the data from ovicups/ovitraps placed in your yard will be protected. Household locations will be coded so as not to use family names on data records. The signed consent forms will be stored in a locked, separate location from data records. Your name and any information that would allow you to be identified as an individual will not be used in data presentations, publications or reports resulting from this study.

If it becomes necessary to review study records, information that can be linked to you will be protected to the extent permitted by law. Your research records will not be released without your consent unless required by law or a court order.

#### What are the possible benefits of this study?

You will receive no direct benefit from participating in this study; however, there may be societal benefits as the data collected from this study may be used by the Belize Ministry of Health, Division of Vector Control to guide activities in improving or modifying mosquito larvicide control measures that target *Ae. aegypti* and *Ae. albopictus* mosquitoes that may have the potential to spread diseases such as Zika, dengue and Chikungunya.

### Do you have to participate?

No, your participation is voluntary. You may decide not to participate at all or, if you start the study, you may withdraw at any time. Withdrawal or refusing to participate will not affect your relationship with the Belize Vector Ecology Center (BVEC) in anyway.

If you would like to participate in this study, sign and return the form to the principal investigator(s). You will receive a copy of this form.

### Will there be any compensation?

Signature of Person obtaining consent

You will not receive any type of payment, compensation or reimbursement for participating in this study.

### Whom to contact with questions about the study?

Questions can be answered prior to, during or after your participation by contacting BVEC researchers Marla Magaña, Donovan Leiva or Jonathan Kay at 322 -1149 or send an email to byec.bz@gmail.com.

### Whom to contact with questions concerning your rights as a research participant?

For questions about your rights or any dissatisfaction with any part of this study, you can contact, anonymously if you wish, BVEC researchers Marla Magaña, Donovan Leiva or Jonathan Kay at 322 - 1149 or send an email to <a href="mailto:bvec.bz@gmail.com">bvec.bz@gmail.com</a>.

#### **Participation**

If you agree to participate please sign below and return the form to the primary researcher(s).

# Signature

this form. You have been given the opportunity to ask	questions before you sign, and you have been told that you can ask articipate in this study. By signing this form, you are 18 years or older
Printed Name	
Signature	Date
As a representative of this study, I have explained the study.	purpose, procedures, benefits, and the risks involved in this research
Print Name of Person obtaining consent	

Date

# **APPENDIX VI Glossary**

## **Glossary of Terms**

## Types of Structures for Mapping Surveillance:

- **1. House:** Any household dwelling unit.
- 2. House/Store: Is a household dwelling unit with a small grocery store attached to the structure.
- 3. Store: An independently established grocery store/supermarket/shop.
- **4. Apartment/Hotel:** A structure that has rooms or living quarters for rent.
- **5.** Lab: Any Medical diagnostic center or laboratory.
- **6. Church:** A structure used for public worship.
- 7. Bar: An establishment that is authorized to sell solely alcoholic beverages and light food. May have entertainment such as slot machines and pool tables. Commonly known also as "Cool Spots" or "Lounge" or "Pubs")
- **8. House/Bar:** A structure with the combination of a house and a bar.
- **9. Restaurant:** An establishment that is authorized to sell food and drinks. May also sell alcoholic beverages. This also includes small vendor establishments.
- **10. School:** An institution that is used for learning and higher education. Either Primary, Secondary or Tertiary level.
- 11. Cemetery: Grounds for which they deceased are laid.
- **12. Office:** A government or non-governmental organization office/department. Example: (Immigration, Customs, etc.)
- **13. Clinic:** Regional Hospital or any other privately owned establishment where outpatients are given medical treatment or advice, especially of a specialist nature.
- **14. Business:** Refers to any other small business establishments such as beauty salons, barbershops, garages, car wash etc.
- 15. Other: Any structure that is distinct from the ones previously named

### **Types of Containers:**

- 1. Vat/Tank: A large metallic or plastic container used for storing rain water at homes. Usually collects rain water from gutters that run at the end of roofs.
- **2. Drum:** Refers to a metallic or plastic 50 gallon container used for rain water storage. May be used in place of a vat/tank.
- 3. Bucket: A small 1-5 gallon container used to store water (either rain or tap) on a temporary basis.
- **4. Bottles/Cans:** Refers to any small empty plastic, glass, aluminum bottles or cans (usually soft-drink or beer) that is found at the premises.
- **5. Tires:** Any discarded vehicle tires that do not contain a rim and are able to hold water.
- **6. Tree-holes/wells:** Are holes that are natural or man-made in a tree (tree-holes) or in the ground (uncovered wells) that can collect water.
- 7. Other: Any container that is distinct from the ones previously named