

Original article

Comparison of microRNA expression in high-count monoclonal B-cell lymphocytosis and Binet A chronic lymphocytic leukemia



Felipe Magalhães Furtado^{a,*}, Priscila Santos Scheucher^a, Bárbara Amélia Santana^a, Dalila Lucíola Zanette^b, Rodrigo do Tocantins Calado^a, Eduardo Magalhães Rego^a, Daniel Mazza Matos^c, Roberto Passetto Falcão^a

^a Faculdade de Medicina de Ribeirão Preto (FMRP), Ribeirão Preto, SP, Brazil

^b Fundação Oswaldo Cruz (FIOCRUZ), Salvador, BA, Brazil

^c Hospital Universitário Walter Cantídio (HUWC), Fortaleza, CE, Brazil

ARTICLE INFO

Article history:

Received 30 November 2016

Accepted 24 March 2017

Available online 22 April 2017

Keywords:

Monoclonal B-cell lymphocytosis
Chronic lymphocytic leukemia
microRNA

ABSTRACT

Background: Evidence suggests that monoclonal B-cell lymphocytosis precedes all chronic lymphocytic leukemia cases, although the molecular mechanisms responsible for disease progression are not understood. Aberrant miRNA expression may contribute to the pathogenesis of chronic lymphocytic leukemia. The objective of this study was to compare miRNA expression profiles of patients with Binet A chronic lymphocytic leukemia with those of subjects with high-count monoclonal B-cell lymphocytosis and healthy volunteers (controls). **Methods:** Twenty-one chronic lymphocytic leukemia patients, 12 subjects with monoclonal B-cell lymphocytosis and ten healthy volunteers were enrolled in this study. Flow cytometry CD19⁺CD5⁺-based cell sorting was performed for the chronic lymphocytic leukemia and monoclonal B-cell lymphocytosis groups and CD19⁺ cells were sorted to analyze the control group. The expressions of miRNAs (miR-15a, miR-16-1, miR-29b, miR-34a, miR-181a, miR-181b and miR-155) were determined by quantitative reverse transcriptase polymerase chain reaction (qRT-PCR).

Results: Significant differences between the expressions in the chronic lymphocytic leukemia and monoclonal B-cell lymphocytosis groups were restricted to the expression of miR-155, which was higher in the former group. A comparison between healthy controls and monoclonal B-cell lymphocytosis/chronic lymphocytic leukemia patients revealed higher miR-155 and miR-34a levels and lower miR-15a, miR-16-1, miR-181a and miR-181b in the latter group.

* Corresponding author at: Laboratório de Hematologia – Hospital das Clínicas de Ribeirão Preto, Bandeirantes Avenue, 3900, Monte Alegre, 14051-140 Ribeirão Preto, SP, Brazil.

E-mail address: felipemf@usp.br (F.M. Furtado).

<http://dx.doi.org/10.1016/j.bjhh.2017.03.006>

1516-8484/© 2017 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Conclusions: Our results show a progressive increase of miR-155 expression from controls to monoclonal B-cell lymphocytosis to chronic lymphocytic leukemia. The role of miR-155 in the development of overt chronic lymphocytic leukemia in individuals with monoclonal B-cell lymphocytosis must be further analyzed.

© 2017 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Chronic lymphocytic leukemia (CLL) is a malignant neoplasm characterized by an excess of monoclonal B lymphocytes in peripheral blood, bone marrow, the spleen and lymph nodes.¹ CLL is the most common leukemia in the Western world, with an annual incidence of 5.1 cases/100,000 people.²

Over the last years, advances in multi-parameter flow cytometry allowed the identification of small populations of monoclonal B lymphocytes in the blood of apparently healthy subjects.³ The presence of monoclonal B-cells in the peripheral blood of up to $5 \times 10^9/L$ is classified as monoclonal B-cell lymphocytosis (MBL) in the absence of other lymphomatous features.^{4,5} This condition is found in 0.6–12% of healthy individuals.^{3,6,7}

MBL is classified as high-count or low-count MBL. The first is usually diagnosed in asymptomatic subjects with mild lymphocytosis and the latter occurs in asymptomatic subjects with normal blood counts who have been submitted to flow cytometry screening.^{8–10} The most recent World Health Organization (WHO) classification defines low-count MBL as a peripheral blood CLL count of $<0.5 \times 10^9/L$ and no extramedullary disease.⁴ As in CLL, MBL is more common in men and in relatives of CLL patients^{11–13} with its frequency increasing with age.^{11,12,14–17}

Virtually, all CLL cases are preceded by MBL.¹⁸ Even though the mechanisms underlying CLL pathogenesis and MBL progression to CLL are still not well understood, it is speculated that the initial genetic lesion in MBL may occur in the immature bone marrow B cell compartment and afterwards repetitive antigenic stimulation may induce additional genetic lesions, eventually leading to neoplastic transformation.¹⁹

MicroRNAs (miRNA) are 19–25 nucleotide single strand RNAs responsible for gene expression and cellular metabolism regulation.^{20,21} Changes in miRNA expression have been associated with solid and hematologic tumors.²¹ Some microRNAs have been shown to be abnormally expressed in CLL.²² Studies addressing the molecular and genetic basis of MBL may help to elucidate initial steps of CLL pathogenesis and, therefore, increase knowledge about CLL origin. Moreover, only a few studies have investigated miRNAs in MBL.^{9,23} Here we hypothesized that abnormalities in some miRNA expressions may be present in MBL and may have a role in initial monoclonal B-cell expansion.

Methods

Cell samples from normal controls, individuals with high-count MBL and CLL patients

Samples from 12 individuals with high-count MBL and 21 patients with stage A CLL according to the Binet classification were analyzed. All the patients were being monitored in a university hospital in Ribeirao Preto, Sao Paulo, Brazil. Ten healthy individuals were studied as a control group. The study was approved by the Institution Ethics Research Review Board and written informed consent was obtained from all participants in accordance with the Declaration of Helsinki and its revisions.

Using cell sorting by flow cytometry, CD19⁺CD5⁺ lymphocytes were isolated from the peripheral blood of individuals in the two study groups, and CD19⁺ lymphocytes from the peripheral blood of controls as previously described.²⁴ The mean percentage of the desired cell population after isolation was 89.48% ($\pm 8.48\%$).

The clinical characteristics of the enrolled subjects are summarized in [Table 1](#).

RNA isolation and miRNA expression

RNA was extracted from CD5⁺ B lymphocytes isolated from the study groups and B lymphocytes from the control group, using Trizol (Life Technologies, California, USA). Seven miRNAs known to have abnormal expressions in CLL were selected for analysis (miR-15a, miR-16-1, miR-29b, miR-34a, miR-181a, miR-181b and miR-155). miRNA expression was quantified by TaqMan miRNA quantitative reverse-transcription polymerase chain reaction (qRT-PCR) as previously described.^{25,26} Briefly, 5 ng of total RNA were reverse transcribed using the microRNA reverse-transcription kit (Applied Bio-systems, California, USA) with specific stem-loop primers. qRT-PCR analysis was performed using the miRNA-specific Taqman assay (Applied Bio-systems). Expression levels of the nucleolar RNAs RNU24, RNU44 and RNU48 were similar for all groups and the geometric mean of their expression was used to normalize the expression of the miRNAs, as previously described²⁷ using the $2^{-\Delta\Delta Ct}$ formula. The coefficients of variation (CV) of the reactions were calculated. If the CV of a reaction was greater than 2% in two successive experiments,

Table 1 – Characteristics of the enrolled subjects.

Characteristic	Control	High-count monoclonal B-cell lymphocytosis	Chronic lymphocytic leukemia
Number of subjects	10	12	21
Medium age	38	78	70
Range (years)	25–60	57–97	58–81
Gender (n)			
Male	2	9	11
Female	8	3	10
Medium number of lymphocytes	2.064	3.933	41.714
Range (μ L)	1.360–3.300	2.900–5.900	7.500–147.000
% CD5 ⁺ B-Lymphocytes of total lymphocytes	–	34.9	78.02
Range (%)	–	5.97–61.01	58.63–91.09

its value was considered unavailable. All miRNAs were studied in duplicate.

Statistical analysis

The Kolmogorov–Smirnov normality test was initially performed to compare miRNA expressions. The expressions of the miRNAs, miR-15a, miR-16-1, miR-29b, miR-181b and miR-155, did not deviate from the normal distribution and variance analysis (ANOVA) was used followed by the Bonferoni *post hoc* test. The Kruskal–Wallis non-parametric test (non-parametric ANOVA) was used followed by the Dunn *post hoc* test for the miRNAs miR-34a and miR-181a that had non-normal distributions. All tests with p -values ≤ 0.05 were considered statistically significant.

Results

miRNA expression

Of the seven miRNAs deemed relevant for CLL pathogenesis, miRNA-155 and miR-34a were differentially expressed in the CLL and high-count MBL groups compared to the controls (p -value=0.33 and p -value=0.003 for MBL and controls and p -value <0.001 and p -value=0.02 for CLL and controls, respectively) with the highest values detected in CLL (Figure 1). miRNA-155 expression in high-count MBL had a tendency to be greater than in healthy subjects, but this difference was not statistically significant, maybe due to the small number of samples studied (Figure 1). miR-34a expression was not statistically different between CLL and MBL (Figure 1). miR-15a, miR-16-1, miR-181a and miR-181b were down-regulated in CLL and MBL. Their expression was lower in CLL patients when compared to controls (p -value <0.001 for all miRNAs). High-count MBL subjects also had lower expression than controls for miR-15a, miR-16-1 and miR-181b (p -value=0.003) and for miR-181a (p -value <0.001) (Figure 1).

Expression of miR-29b was similar for all three groups (Figure 1).

Discussion

This study with a Binet A CLL group and a high-count MBL group demonstrated new molecular differences between these two entities.

Statistical similarities have been identified regarding the frequencies of immunoglobulin heavy chain variable (IGHV) genes between high-count MBL and the initial stages of CLL, although findings from low-count MBL were different between these groups.^{28,29} Other authors also found biological similarities between these three entities. Not only high-count MBL but also low-count MBL bear cytogenetic abnormalities common in CLL, including 13q-, 17p- and trisomy 12.^{6,30–32}

Aiming to better understand genetic alterations in CLL, this study compared the expressions of miRNAs previously described as altered in this disease with their expression in MBL. Knowledge of miRNA expression in different monoclonal proliferation stages may improve the comprehension of CLL pathophysiology.

Several studies using different methodologies have identified overexpression of miR-155 in CLL when compared to normal controls.^{23,33–37} Our results confirmed the seminal findings of Ferrajoli et al. who showed that the expression of miR-155 is greater in Binet A CLL than in high-count MBL and that its expression in these individuals is greater than in normal controls.²³ Thus, our findings suggest that miR-155 may be used as a progression marker for MBL individuals, but this must be confirmed by further specific studies.

As is usual for miRNAs, miR-155 has distinct functions in different cell types. In B-lymphocytes, it represses the expression of PU.1 transcription factor and the inositol 5-phosphatase Src homology-2 domain-containing inositol 5-phosphatase 1 (SHIP1).³⁸ As an antagonist of the phosphatidylinositol 3-kinase (PI3K) pathway, SHIP1 is a negative regulator of B cell receptor (BCR) signaling and it was identified as a target of miR-155 in diffuse B-cell lymphoma.³⁹ In CLL, the regulation of miR-155 and its target genes are not well understood. Its expression may be regulated by transcription factors that have important roles in its evolution. Furthermore, miR-155 may directly regulate coding genes that are important for the transition of MBL to CLL.²³




Figure 1 – MicroRNA expression of control, monoclonal B-cell lymphocytosis and chronic lymphocytic leukemia groups: (A) miR-155; (B) miR-34a; (C) miR-15a; (D) miR-16-1; (E) miR-181a; (F) miR-181b; (G) miR-29b.

This study also identified miR-34a to be overexpressed in CLL when compared to healthy controls.^{35,40,41} This miRNA has been associated to the regulation of the tumor protein p53 (TP53) pathway⁴¹⁻⁴⁵ and may be related to fludarabine resistance in CLL.⁴⁴

The study that first demonstrated down-regulation of miR-15a and miR-16-1 in CLL patients was also the first to demonstrate abnormal expressions of miRNAs in cancer.⁴⁶ Subsequent studies using different techniques have confirmed this finding, particularly in patients with del 13q14.⁴⁷⁻⁴⁹ Other authors found similar expressions of these miRNAs in CLL and normal controls using microarrays³⁵ and RT-qPCR.³³ Klein et al. demonstrated that mouse models that have the chromosome area (13q14) responsible for the transcription of these miRNAs deleted developed monoclonal expansion of lymphocytes in blood.⁵⁰

In this study, the miRNAs miR-181a and miR-181b were down-regulated in CLL when compared to normal controls. Other authors found similar results in studies using RT-qPCR,^{36,48} Northern Blot³⁴ and microarrays.³⁵ Visone et al. demonstrated that down-regulation of miR-181b is a better marker for worse prognosis in CLL than the IGHV mutation status and ZAP-70 expression, and suggest that its expression should be monitored in CLL patients.⁵¹

The pattern of miR-29b expression in CLL is not well understood. Here the results for CLL Binet A, high-count MBL and normal controls were similar. Sampath et al. found that 70% of the CLL patients have a similar expression to normal controls.⁴⁹ Fulci et al. also had results similar to this study.³³ However, Santanam et al. found this miRNA to be overexpressed in CLL, and Zhu et al. found it to be down-regulated in CLL.^{48,52} All these studies have heterogeneous groups of CLL patients regarding the stage of the disease. These different results suggest that miR-29b expression may change according to disease stage. However, this hypothesis should be confirmed in future studies.

Apart from the study of Ferrajoli et al. on miR-155, only one other group has evaluated miRNA expression in high-count MBL.²³ Thus, Morabito et al. used microarrays to study individuals previously diagnosed with CLL before the change of diagnosis criteria proposed by the International Workshop on Chronic Lymphocytic Leukemia in 2008.^{9,53} They found miR-130a was the only one with different expressions between CLL and high-count MBL. Using another technique and a more heterogeneous high-count MBL population, this study found the expressions of the miR-34a, miR-15a, miR-16-1, miR-181a, miR-181b and miR-29b to be similar in CLL and high-count MBL.

The results reported here should be confirmed by further studies, since the groups were small and more individuals must be studied to allow robust conclusions. Moreover, although the CLL and MBL groups were composed mainly of older individuals and the MBL group mainly of males, our control group was composed mainly of females and the medium age for this group was lower than for the study groups, although there are no data suggesting that the levels of these miRNAs change with sex and age.

In conclusion, the finding that some miRNAs have abnormal expressions in MBL and that this condition is also present in CLL suggests that these genetic changes may be part of the

initial events responsible for monoclonal CD5⁺ B-cell proliferation. The differential expression of miR-155 in MBL and CLL suggests its role in signaling pathways that are important to the development of the disease.

Author contributions

FMF, DLZ, RTCSR, EMR, DMM and RPF designed the research protocols. FMF, PSS and BS performed the research. FMF, BS, DMM and RPF analyzed the data. FMF, RTCSR, EMR, DMM and RPF wrote the manuscript.

Conflict of interests

The authors declare no conflicts of interest.

Acknowledgements

This research was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPQ) grant 573.754/2008-0.

REFERENCES

1. Müller-Hermelink HK, Montserrat E, Catovsky D, Campo E, Harris NL, Stein H. Chronic lymphocytic leukaemia/small lymphocytic lymphoma. In: Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri SA, Stein H, et al., editors. WHO classification of tumors of haematopoietic and lymphoid tissues. 4th ed Lyon: World Health Organization Classification of Tumors. International Agency for Research on Cancer; 2008. p. 180-2.
2. Dores GM, Anderson WF, Curtis RE, Landgren O, Ostroumova E, Bluhm EC, et al. Chronic lymphocytic leukaemia and small lymphocytic lymphoma: overview of the descriptive epidemiology. *Br J Haematol*. 2007;139(5):809-19.
3. Rawstron AC, Green MJ, Kuzmicki A, Kennedy B, Fenton JA, Evans PA, et al. Monoclonal B lymphocytes with the characteristics of indolent chronic lymphocytic leukemia are present in 3.5% of adults with normal blood counts. *Blood*. 2002;100(2):635-9.
4. Swerdlow SH, Campo E, Pileri SA, Harris NL, Stein H, Siebert R, et al. The 2016 revision of the World Health Organization classification of lymphoid neoplasms. *Blood*. 2016;127(20):2375-90.
5. Marti GE, Rawstron AC, Ghia P, Hillmen P, Houlston RS, Kay N, et al. Diagnostic criteria for monoclonal B-cell lymphocytosis. *Br J Haematol*. 2005;130(3):325-32.
6. Rawstron AC, Bennett FL, O'Connor SJ, Kwok M, Fenton JA, Plummer M, et al. Monoclonal B-cell lymphocytosis and chronic lymphocytic leukemia. *N Engl J Med*. 2008;359(6):575-83.
7. D'Arena G, Musto P. Monoclonal B-cell lymphocytosis. *Transl Med UniSa*. 2014;8:75-9.
8. Ghia P, Caligaris-Cappio F. Monoclonal B-cell lymphocytosis: right track or red herring? *Blood*. 2012;119(19):4358-62.
9. Morabito F, Mosca L, Cutrona G, Agnelli L, Tuana G, Ferracin M, et al. Clinical monoclonal B lymphocytosis versus Rai 0 chronic lymphocytic leukemia: a comparison of cellular, cytogenetic, molecular, and clinical features. *Clin Cancer Res*. 2013;19(21):5890-900.

10. Rawstron AC. Monoclonal B cell lymphocytosis – what does it really mean? *Curr Hematol Malig Rep*. 2013;8(1):52–9.
11. Rawstron AC, Yuille MR, Fuller J, Cullen M, Kennedy B, Richards SJ, et al. Inherited predisposition to CLL is detectable as subclinical monoclonal B-lymphocyte expansion. *Blood*. 2002;100(7):2289–90.
12. Matos DM, Ismael SJ, Scrideli CA, de Oliveira FM, Rego EM, Falcao RP. Monoclonal B-cell lymphocytosis in first-degree relatives of patients with sporadic (non-familial) chronic lymphocytic leukaemia. *Br J Haematol*. 2009;147(3):339–46.
13. Goldin LR, Lanasa MC, Slager SL, Cerhan JR, Vachon CM, Strom SS, et al. Common occurrence of monoclonal B-cell lymphocytosis among members of high-risk CLL families. *Br J Haematol*. 2010;151(2):152–8.
14. Shanafelt TD, Kay NE, Jenkins G, Call TG, Zent CS, Jelinek DF, et al. B-cell count and survival: differentiating chronic lymphocytic leukemia from monoclonal B-cell lymphocytosis based on clinical outcome. *Blood*. 2009;113(18):4188–96.
15. Ghia P, Prato G, Scielzo C, Stella S, Geuna M, Guida G, et al. Monoclonal CD5+ and CD5– B-lymphocyte expansions are frequent in the peripheral blood of the elderly. *Blood*. 2004;103(6):2337–42.
16. Mulligan CS, Thomas ME, Mulligan SP. Monoclonal B-lymphocytosis: demographics, nature and subclassification in 414 community patients. *Leuk Lymphoma*. 2011;52(12):2293–8.
17. Kern W, Bacher U, Haferlach C, Dicker F, Alpermann T, Schnittger S, et al. Monoclonal B-cell lymphocytosis is closely related to chronic lymphocytic leukaemia and may be better classified as early-stage CLL. *Br J Haematol*. 2012;157(1):86–96.
18. Landgren O, Albitar M, Ma W, Abbasi F, Hayes RB, Ghia P, et al. B-cell clones as early markers for chronic lymphocytic leukemia. *N Engl J Med*. 2009;360(7):659–67.
19. Chiorazzi N, Rai KR, Ferrarini M. Chronic lymphocytic leukemia. *N Engl J Med*. 2005;352(8):804–15.
20. Alvarez-Garcia I, Miska EA. MicroRNA functions in animal development and human disease. *Development*. 2005;132(21):4653–62.
21. Chen B, Li H, Zeng X, Yang P, Liu X, Zhao X, et al. Roles of microRNA on cancer cell metabolism. *J Transl Med*. 2012;10:228.
22. Ward BP, Tsongalis GJ, Kaur P. MicroRNAs in chronic lymphocytic leukemia. *Exp Mol Pathol*. 2011;90(2):173–8.
23. Ferrajoli A, Shanafelt TD, Ivan C, Shimizu M, Rabe KG, Nouraei N, et al. Prognostic value of miR-155 in individuals with monoclonal B-cell lymphocytosis and patients with B chronic lymphocytic leukemia. *Blood*. 2013;122(11):1891–9.
24. Basu S, Campbell HM, Dittel BN, Ray A. Purification of specific cell population by fluorescence activated cell sorting (FACS). *J Vis Exp*. 2010;41.
25. Chen C, Ridzon DA, Broomer AJ, Zhou Z, Lee DH, Nguyen JT, et al. Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Res*. 2005;33(20):e179.
26. Schmittgen TD, Lee EJ, Jiang J, Sarkar A, Yang L, Elton TS, et al. Real-time PCR quantification of precursor and mature microRNA. *Methods*. 2008;44(1):31–8.
27. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, et al. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol*. 2002;3(7). Research0034.
28. Dagklis A, Fazi C, Sala C, Cantarelli V, Scielzo C, Massacane R, et al. The immunoglobulin gene repertoire of low-count chronic lymphocytic leukemia (CLL)-like monoclonal B lymphocytosis is different from CLL: diagnostic implications for clinical monitoring. *Blood*. 2009;114(1):26–32.
29. Vardi A, Dagklis A, Scarfo L, Jelinek D, Newton D, Bennett F, et al. Immunogenetics shows that not all MBL are equal: the larger the clone, the more similar to CLL. *Blood*. 2013;121(22):4521–8.
30. Nieto WG, Almeida J, Romero A, Teodosio C, Lopez A, Henriques AF, et al. Increased frequency (12%) of circulating chronic lymphocytic leukemia-like B-cell clones in healthy subjects using a highly sensitive multicolor flow cytometry approach. *Blood*. 2009;114(1):33–7.
31. Rossi D, Sozzi E, Puma A, De Paoli L, Rasi S, Spina V, et al. The prognosis of clinical monoclonal B cell lymphocytosis differs from prognosis of Rai 0 chronic lymphocytic leukaemia and is recapitulated by biological risk factors. *Br J Haematol*. 2009;146(1):64–75.
32. Fazi C, Scarfo L, Pecciarini L, Cottini F, Dagklis A, Janus A, et al. General population low-count CLL-like MBL persists over time without clinical progression, although carrying the same cytogenetic abnormalities of CLL. *Blood*. 2011;118(25):6618–25.
33. Fulci V, Chiaretti S, Goldoni M, Azzalin G, Carucci N, Tavolaro S, et al. Quantitative technologies establish a novel microRNA profile of chronic lymphocytic leukemia. *Blood*. 2007;109(11):4944–51.
34. Marton S, Garcia MR, Robello C, Persson H, Trajtenberg F, Pritsch O, et al. Small RNAs analysis in CLL reveals a deregulation of miRNA expression and novel miRNA candidates of putative relevance in CLL pathogenesis. *Leukemia*. 2008;22(2):330–8.
35. Pallasch CP, Patz M, Park YJ, Hagist S, Eggle D, Claus R, et al. miRNA deregulation by epigenetic silencing disrupts suppression of the oncogene PLAG1 in chronic lymphocytic leukemia. *Blood*. 2009;114(15):3255–64.
36. Li S, Moffett HF, Lu J, Werner L, Zhang H, Ritz J, et al. MicroRNA expression profiling identifies activated B cell status in chronic lymphocytic leukemia cells. *PLoS One*. 2011;6(3).
37. Vargova K, Curik N, Burda P, Basova P, Kulvait V, Pospisil V, et al. MYB transcriptionally regulates the miR-155 host gene in chronic lymphocytic leukemia. *Blood*. 2011;117(14):3816–25.
38. Danger R, Braza F, Giral M, Soullillou JP, Brouard S. MicroRNAs major players in B cells homeostasis and function. *Front Immunol*. 2014;5:98.
39. Schott J, Stoecklin G. Networks controlling mRNA decay in the immune system. *Wiley Interdiscip Rev RNA*. 2010;1(3):432–56.
40. Zanette DL, Rivadavia F, Molfetta GA, Barbuzano FG, Proto-Siqueira R, Silva-Jr WA, et al. miRNA expression profiles in chronic lymphocytic and acute lymphocytic leukemia. *Braz J Med Biol Res*. 2007;40(11):1435–40.
41. Asslaber D, Pinon JD, Seyfried I, Desch P, Stocher M, Tinhofer I, et al. microRNA-34a expression correlates with MDM2 SNP309 polymorphism and treatment-free survival in chronic lymphocytic leukemia. *Blood*. 2010;115(21):4191–7.
42. Bommer GT, Gerin I, Feng Y, Kaczorowski AJ, Kuick R, Love RE, et al. p53-mediated activation of miRNA34 candidate tumor-suppressor genes. *Curr Biol*. 2007;17(15):1298–307.
43. Hermeking H. The miR-34 family in cancer and apoptosis. *Cell Death Differ*. 2010;17(2):193–9.
44. Merkel O, Asslaber D, Pinon JD, Egle A, Greil R. Interdependent regulation of p53 and miR-34a in chronic lymphocytic leukemia. *Cell Cycle*. 2010;9(14):2764–8.
45. Dufour A, Palermo G, Zellmeier E, Mellert G, Duchateau-Nguyen G, Schneider S, et al. Inactivation of TP53 correlates with disease progression and low miR-34a expression in previously treated chronic lymphocytic leukemia patients. *Blood*. 2013;121(18):3650–7.
46. Calin GA, Dumitru CD, Shimizu M, Bichi R, Zupo S, Noch E, et al. Frequent deletions and down-regulation of micro-RNA genes miR15 and miR16 at 13q14 in chronic lymphocytic leukemia. *Proc Natl Acad Sci U S A*. 2002;99(24):15524–9.
47. Calin GA, Liu CG, Sevignani C, Ferracin M, Felli N, Dumitru CD, et al. MicroRNA profiling reveals distinct signatures in B cell

- chronic lymphocytic leukemias. *Proc Natl Acad Sci U S A*. 2004;101(32):11755-60.
48. Zhu DX, Miao KR, Fang C, Fan L, Zhu W, Zhu HY, et al. Aberrant microRNA expression in Chinese patients with chronic lymphocytic leukemia. *Leuk Res*. 2011;35(6):730-4.
49. Sampath D, Liu C, Vasani K, Sulda M, Puduvalli VK, Wierda WG, et al. Histone deacetylases mediate the silencing of miR-15a, miR-16, and miR-29b in chronic lymphocytic leukemia. *Blood*. 2012;119(5):1162-72.
50. Klein U, Lia M, Crespo M, Siegel R, Shen Q, Mo T, et al. The DLEU2/miR-15a/16-1 cluster controls B cell proliferation and its deletion leads to chronic lymphocytic leukemia. *Cancer Cell*. 2010;17(1):28-40.
51. Visone R, Veronese A, Rassenti LZ, Balatti V, Pearl DK, Acunzo M, et al. miR-181b is a biomarker of disease progression in chronic lymphocytic leukemia. *Blood*. 2011;118(11):3072-9.
52. Santanam U, Zanesi N, Efanov A, Costinean S, Palamarchuk A, Hagan JP, et al. Chronic lymphocytic leukemia modeled in mouse by targeted miR-29 expression. *Proc Natl Acad Sci U S A*. 2010;107(27):12210-5.
53. Hallek M, Cheson BD, Catovsky D, Caligaris-Cappio F, Dighiero G, Dohner H, et al. Guidelines for the diagnosis and treatment of chronic lymphocytic leukemia: a report from the International Workshop on Chronic Lymphocytic Leukemia updating the National Cancer Institute-Working Group 1996 guidelines. *Blood*. 2008;111(12):5446-56.