

Knowledge and attitude: profile of diabetics in dialysis

Conhecimento e Atitude: perfil de pessoas com diabetes em diálise

Conocimiento y actitud: perfil de personas con diabetes en diálisis

Claudia Capellari[✉], Ana Elizabeth Prado Lima Figueiredo[✉]

ABSTRACT

Objective: to identify knowledge of, and coping with, diabetes mellitus among diabetics undergoing dialysis. **Method:** in this cross-sectional study of patients with type 2 diabetes, the instruments used were the Diabetes Knowledge Scale Questionnaire (DKN-A) and Diabetes Attitude Questionnaire (ATT-19). **Results:** mean age of the 71 participants was 61.81 ± 14.93 years. The most prevalent complication was diabetic retinopathy (81.69%), and the most prevalent comorbidity was systemic arterial hypertension (83.09%). Median fasting glycemia and glycated hemoglobin were 152 (124-228.5) mg/dl and 7.5 (6.42-8.27) mg/dl, respectively. Average DKN-A score was 7.84 ± 2.55 ; the highest success rate was on how to respond to hypoglycemia; the lowest was about ketones in urine and substitute foods. Mean ATT-19 score was 50.26 ± 11.7 . **Conclusion:** the diabetics in dialysis showed deficient knowledge of diabetes and had negative attitudes to the disease.

Descriptors: Kidney Failure, Chronic; Diabetes Mellitus; Nephrology Nursing; Health Education; Knowledge.

RESUMO

Objetivo: identificar o conhecimento e o enfrentamento do diabetes junto a pessoas com diabetes em diálise. **Método:** estudo transversal, realizado com pacientes com diabetes tipo 2. Os instrumentos utilizados foram o *Diabetes Knowledge Scale Questionnaire* (DKN-A) e o *Diabetes Attitude Questionnaire* (ATT-19). **Resultados:** participaram 71 pacientes, com idade média de $61,81 \pm 14,93$ anos. A retinopatia diabética foi a complicação prevalente (81,69%); hipertensão arterial sistêmica foi a comorbidade (83,09%). A glicemia em jejum apresentou mediana de 152 (124-228,5) mg/dl e a hemoglobina glicada de 7,5 (6,42-8,27) mg/dl. O DKN-A apresentou escore médio de $7,84 \pm 2,55$ pontos; seu item com maior número de acertos foi a conduta em caso de hipoglicemia; enquanto o com menor número de acertos foi em relação à cetonúria e substituições alimentares. O instrumento ATT-19 obteve média de $50,26 \pm 11,7$ pontos. **Conclusão:** pessoas com diabetes, em diálise, apresentam conhecimento deficiente em relação ao diabetes, assim como baixo enfrentamento da doença.

Descritores: Insuficiência Renal Crônica; Diabetes Mellitus; Enfermagem em Nefrologia; Educação em Saúde; Conhecimento.

RESUMEN

Objetivo: identificar el conocimiento y el afrontamiento de la diabetes mellitus entre los diabéticos en diálisis. **Método:** en este estudio transversal de pacientes con diabetes tipo 2, los instrumentos utilizados fueron el *Diabetes Knowledge Scale Questionnaire* (DKN-A) y *Diabetes Attitude Questionnaire* (ATT-19). **Resultados:** la edad media de los 71 participantes fue de $61,81 \pm 14,93$ años. La complicación más prevalente fue la retinopatía diabética (81,69%) y la comorbilidad más prevalente fue la hipertensión arterial sistémica (83,09%). La mediana de la glucemia en ayunas y la hemoglobina glucosilada fueron 152 (124-228,5) mg / dl y 7,5 (6,42-8,27) mg / dl, respectivamente. La puntuación promedio de DKN-A fue de $7,84 \pm 2,55$; la tasa de éxito más alta fue sobre cómo responder a la hipoglucemia; el más bajo fue sobre las cetonas en la orina y los alimentos sustitutos. La puntuación media de ATT-19 fue $50,26 \pm 11,7$. **Conclusión:** los diabéticos en diálisis mostraban un conocimiento deficiente de la diabetes y actitudes negativas hacia la enfermedad.

Descriptores: Fallo Renal Crónico; Diabetes Mellitus; Enfermería en Nefrología; Educación en Salud; Conocimiento.

INTRODUCTION

Diabetes Mellitus (DM) is one of the most common chronic diseases in the world, affecting nearly 425 million people¹. In Brazil, it reaches around 7.6% of the population, representing more than 15 million individuals².

It is well described that adequate glycemic control is fundamental for the reduction of harms and chronic complications. Major studies show that glycemic levels are directly proportional to the complications presented by DM carriers^{3,4}. In this sense, the maintenance of acceptable levels of glucose has revealed benefits related to the complications, in special those associated with microvasculopathy and neuropathy^{3,5,6}.

One of most critical consequences of diabetes is Chronic Kidney Disease (CKD). For people with both these conditions, recommendations are that glycemic control be part of a multifactorial intervention strategy to prevent microvascular lesions and others injuries^{7,8}.

Acknowledgments to the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Brazil.
Corresponding author: Claudia Capellari. E-mail: capellaric@gmail.com.
Editor in charge: Cristiane Helena Gallasch.

For treatment success, it is important that the patient participates in the care process. In this way, it is believed that knowledge of the disease and its control alternatives are essential for minimizing adverse events and for better quality of life. In this context, health education emerges with potential for an improvement in patient empowerment. From the aforementioned, it is inferred that professionals are able to act as motivators, facilitators and promoters of the awareness of individuals, contributing to adherence to treatment, to the development of self-care capacity, and to lifestyle change^{9,10}.

This study aimed to identify the knowledge and attitudes of people with diabetes and chronic kidney disease on dialysis.

THEORETICAL REFERENTIAL

For healthcare professionals, being aware of how well patients understand their pathology and their attitude towards it is valuable. Such data may guide health education strategies and guidelines for adherence to treatment. Although there are few studies regarding the focus population of this study, a case study recommends interventions planned by the nurse¹¹.

In this sense, some instruments may support the decision-making of professionals by providing necessary information on the patients' knowledge about their own disease and the strategies they use to address it. Two of them are the Diabetes Knowledge Questionnaire (DKN-A)^{12,13} and the Diabetes Attitude Questionnaire (ATT-19)^{13,14}, validated for the Brazilian reality.

DKN-A has 15 multiple-answer items and covers aspects related to general knowledge on diabetes mellitus. The questions are about basic physiology, food groups and DM management. The measuring scale used is from 0 to 15. A score of one (1) is attributed to the correct answer and of zero (0) for the incorrect answer. A score higher than eight (8) indicates knowledge about diabetes mellitus¹⁵.

ATT-19 is an instrument that seeks to measure psychological adjustment for diabetes mellitus, developed in response to the needs of evaluation of the psychological and emotional aspects of the disease. It contains nineteen items that include six factors: a) DM-associated stress, b) treatment receptivity, c) trust in the treatment, d) personal efficiency, e) health perception, and f) social acceptance, with the answers measured using a five-point *Likert* scale (completely disagree - score 1; up to completely agree - score 5). The total value of the score can vary from 19 to 95 points. A score higher than 70 indicates a positive attitude towards the disease¹⁵. In this instrument, attitude is related to the decision of the individual to adopt or not the self-care measures for diabetes control¹⁶.

These questionnaires have been referred to in a series of studies that measure the impact of interventions on people with diabetes, especially those involving health education, or which have investigated knowledge, coping and self-care as subsidiary data for health team actions¹⁶⁻²⁵. Thus, assessing knowledge as well as emotional aspects is important to provide care and information consistent with the population served.

METHOD

This is a cross-sectional study, approved by the Ethics and Research Committee of *Pontifícia Universidade Católica do Rio Grande do Sul* (PUCRS) under Certificate of Presentation for Ethical Appreciation (*Certificado de Apresentação para Apreciação Ética*, CAAE) number 51381215.4.0000.5336.

The study was conducted in two dialysis services in the state of Rio Grande do Sul, Brazil. Data was collected between November 2016 and April 2017.

The population consisted of diabetics on dialysis (hemodialysis and peritoneal dialysis), linked to the aforementioned services: a total of 118 individuals. In the sample, the individuals included were those with type 2 diabetes, over 18 years of age, literate, and who agree to participate in the study by signing the free and informed consent form.


The patients excluded were those with a serious deficiency not corrected in hearing, speaking, and with total amaurosis or with degenerative neurological disease, reported in medical records. Also, all those who obtained less than 20 points in the Mini Mental State Examination (MMSE)²⁶.

The selection of the sample followed the diagram shown in Figure 1.

The patients who met the inclusion criteria were invited to participate and those who accepted signed the free and informed consent form and answered the MMSE.

For those with a score of 20 or more on the MMSE, an instrument was applied to collect sociodemographic and clinical variables, which included age, skin color, gender, marital status, schooling, income, distance from home in relation to the dialysis service, time of diabetes diagnosis, medications, chronic complications and comorbidities, laboratory values

of fasting blood glucose, glycated hemoglobin, cholesterol and triglycerides. Sociodemographic data were collected at the bedside, during a dialysis session, by the researcher. Clinical data were obtained by consulting the medical record.



*MMSE: Mini Mental State Examination

FIGURE 1: Sample selection diagram.

On the same occasion of the collection of sociodemographic data, the participants answered the DKN-A and ATT-19 questionnaires¹⁵. For that, the Survey Monkey® application was used, via a tablet with Internet access. The patients themselves, with the equipment offered by the researcher, answered the questionnaires, after being explained how to fill them out.

The data were extracted directly from the Survey Monkey software, to an Excel file. After organizing the spreadsheets, statistical analysis was performed using the Statistical Package for the Social Sciences (SPSS) software, version 20.0. The results were presented using descriptive statistics - absolute and relative distribution (n - %), measures of central tendency and variability (mean \pm standard deviation, median (1st - 3rd quartile). The confidence interval was 95% (p<0.05).

RESULTS

A total of 71 patients participated in the study, with a mean age of 61.81 \pm 14.93 years old, 62 (87.32%) were white-skinned, 37 (52.11%) were male and 37 (52.11%) were married or in a stable relationship. Among the participants, the median schooling time was 5 (3.5-8.5) years; and 6 (8.45%) had a university degree. Sixty nine (69) participants were retired or received state pensions from the *Instituto Nacional de Seguridade Social* (INSS) and the mean family income was 3.22 (2-4) times the minimum wage; 2 (2.81%) participants were still working. The median distance from the dialysis center to the participants' residences was 19.25 (5-24.5) km.

The median time since the diagnosis of Diabetes Mellitus was 17.88 (10-25) years and, for CKD, 4 (2-8) years. Regarding the antidiabetic agents, 53 (74.64%) participants reported insulin application associated with dietary control as a form of DM treatment. None of the participants took oral hypoglycemic agents.

Regarding chronic complications, in addition to CKD, which affects 100% of the participants, diabetic retinopathy was diagnosed in 58 (81.69%), neuropathy in 48 (67.60%), diabetic foot in 10 (14.08%), vasculopathy in 23 (32.39%), cerebral vascular disease in 17 (23.94%), coronary heart disease in 29 (40.84%) and amputation due to diabetes mellitus in 9 (12.67%) participants.

In relation to comorbidities and risk factors, hypertension was identified in 59 (83.09%) participants, dyslipidemia in 32 (45.07%), obesity in 24 (33.8%), sedentary lifestyle in 41 (57.74%) and smoking in 6 (8.45%).

The median values for fasting glycaemia and glycated hemoglobin were 152 (124-228.5) mg/dl and 7.5 (6.42-8.27) mg/dl, respectively.

In relation to total cholesterol values, the mean was 167 \pm 43.9 mg/dl. Concerning triglycerides, the median was 157 (108.5-235) mg/dl; and, in 38 (53.52%) participants, these values were above 150 mg/dl.

Regarding knowledge about diabetes, Figure 2 shows a dispersion of the scores from the DKN-A questionnaire. The mean score was .84 \pm 2.55 points, and 24 (33.80%) participants had a score higher than 8.




FIGURE 2: Dispersion of the scores from the DKN-A* questionnaire. Porto Alegre, RS, Brazil, 2017.
*DKN-A: Diabetes Knowledge Scale Questionnaire

Table 1 shows the DKN-A questions, grouped according to the attributes, and the number and percentage of correct and incorrect answers for each question.

TABLE 1: Distribution of participants according to the number of answers and errors of the DKN-A*, grouped by attributes (Basic physiology, Food groups and substitutions, DM⁺ management in case of intercurrent and general principles of disease care). Porto Alegre, RS, Brazil, 2017.

nº	Question	Correct		Incorrect	
		n	%	n	%
Basic Physiology					
1	Indication of blood sugar variation when uncontrolled DM [†] (normal / high / low)	56	78.87	15	21.13
3	Indication of the NORMAL range of blood glucose	52	73.23	19	26.77
6	Identification of ketonuria	4	5.63	67	94.37
7	Identification of complications due to DM [†] (vision. kidneys. lungs)	51	71.83	20	28.17
12	Cause of hypoglycaemia	17	23.94	54	76.06
Food groups and substitutions					
4	Butter composition	41	57.74	30	42.26
5	Rice Composition	43	60.56	28	39.44
11	Food that you can eat at will	44	61.97	27	38.03
14	Correct substitutions (French bread / biscuit, egg / ground beef, milk / orange juice, noodles / vegetable soup)	23	32.39	48	67.61
15	Correct substitution of French bread (water and salt biscuit, cheese bread, slice of cheese, let it go)	20	28.16	51	71.84
DM [†] management in case of intercurrence and general principles of disease care					
2	Identification of complications related to DM [†] (coma, glycosuria, late complications)	45	63,38	26	36,32
8	Conduct in case of hyperglycemia	36	50,70	35	49,30
9	Conduct for insulin application in case of illness or inappetence	37	52,11	34	47,89
10	Conduct in case of hypoglycaemia	58	81,69	13	18,31
13	Reference to unit of measure - one kilo	30	42,26	41	57,74

*DKN-A: Diabetes Knowledge Scale Questionnaire; †DM: Diabetes Mellitus

The attitudes towards diabetes can be observed from the values obtained in the ATT-19 questionnaire and shown in Figure 3. The scores ranged from 28 to 77 points, with a mean of 50.26±11.7. Four (5.63%) participants had a score above 70 points.




FIGURE 3: Dispersion of the scores from the ATT-19* questionnaire. Porto Alegre, RS, Brazil, 2017.

Table 2 shows the distribution of the answers to ATT-19 based on a Likert Scale, which goes from Completely Disagree to Completely Agree.

TABLE 2: Distribution of participants according to the ATT-19* Likert Scale. Porto Alegre, RS, Brazil, 2017.

Questions	Disagree completely	Disagree	I don't know	Agree	Agree completely
Stress associated to Diabetes					
1. If I did not have DIABETES I think I would be quite a different person	24	37	1	5	4
3. Diabetes is the worst thing that has ever happened to me	29	18	0	17	7
4. Most people would find it difficult to adjust to having diabetes	13	37	5	16	0
5. I often feel embarrassed about having diabetes	2	5	0	35	29
8. The proper control of diabetes involves a lot of sacrifice and inconvenience	8	38	1	14	10
10. Being told you have diabetes is like being sentenced to a life time of illness	4	27	1	19	20
11. My diabetic diet does not really spoil my social life	4	15	0	39	13
19. I often think it is unfair that I should have diabetes when other people are so healthy	4	8	2	31	26
Receptivity to treatment					
7. There is little hope of leading a normal life with diabetes	10	18	2	21	20
15. Diabetes is not really a problem because it can be controlled	0	3	2	51	15
18. I believe I have adjusted well to having diabetes	2	8	1	39	21
Trust in treatment					
2. I dislike being referred to as a DIABETIC	4	20	1	32	14
6. There is not much I seem to be able to do to control my diabetes	2	19	1	34	15
9. I try not to let people know about my diabetes	2	4	0	41	24
17. There is really no-one I feel I can talk to openly about my diabetes	2	9	2	35	23
Personal effectiveness					
12. In general, doctors need to be a lot more sympathetic in their treatment of people with diabetes	8	36	7	15	5
13. Having diabetes for a long period changes the personality	11	28	6	15	11
Perception about the health					
14. I often find it difficult to decide whether I feel sick or well	4	12	2	37	16
Social acceptance					
16. There is really nothing you can do if you have diabetes	3	18	1	30	19

*ATT-19: Diabetes Attitude Questionnaire

DISCUSSION

This study presents results from patients with diabetes undergoing dialysis. Other investigations have been conducted with people with diabetes, but there is a gap related to people who also have CKD undergoing dialysis. In this context, it is important to look at the knowledge and attitude of the patients involved towards their disease.

In the present research, the participants presented a mean age similar to that of other studies that used the same DM knowledge and attitudes questionnaires²⁷⁻³⁰. However, unlike those, most of the participants in the present study are male. A plausible explanation for this is the fact that the study was carried out with diabetic patients on dialysis, where the number of males prevails in relation to the number of females (58% males)³¹.

The data related to the schooling level corroborate the findings of other studies, which also indicate low levels of schooling^{25,29,32,33}. There is evidence that the schooling level is not related to greater knowledge about DM or to a positive attitude to deal with it³². Nevertheless, schooling may interfere with the understanding of the instructions given, mechanisms related to the disease, and adherence to treatment. These facts constitute a challenge for health professionals, who need to devise strategies to strengthen the cognitive, motor and affective abilities of people with diabetes, so as to achieve satisfactory disease control²⁹.

In addition to low schooling, a low family income was found, even if higher than that reported by a systematic review (1.5 ± 2)³⁴. It is believed that this fact can hinder people's access to basic services, which directly impact on health, such as adequate food, transportation, education and housing.

Regarding employment, other surveys reported occupation rates of 47.7%³² and 30.5%¹⁷, although not conducted among dialysis patients. By contrast, in the present study few participants reported being employed. People on dialysis, in addition to presenting important comorbidities, need to adapt their routine to attend the dialysis center three times a week, which makes it difficult for them to enter the labor market. Moreover, the distance from the residence to the dialysis center would influence the chance to work. This fact points to the need for a flexible schedule, since people spend time traveling, in addition to that related to dialysis treatment.

Regarding the time with the pathology, published studies indicate a wide range, with 9.5 ± 7.9 ³², 11.18 ± 8.64 ²⁹, <10 ^{25,28} and 11 ± 8 years²⁷. The longer time elapsed from the diagnosis of DM presented in this study may be related to the sample of participants (patients with diabetes on dialysis). Diabetic nephropathy is the main microvascular complication of diabetes, as well as the major cause of CKD in the world³⁵, with 30% of dialysis patients having CKD due to diabetes³¹.

Regarding chronic complications, microvasculopathy is strongly associated with diabetes³⁶, when an imbalance occurs between pro-survival factors and inflammatory mediators. In this study, in addition to renal impairment, diabetic retinopathy had the highest prevalence, different from the results found in a study conducted at Primary Care, which found 15% of retinopathy³⁷. In addition to the impacts in terms of human suffering, the complications arising from diabetes directly impact on the cost to the health systems³⁶. The most frequent comorbidity was hypertension, which corroborates the indicator published by the Brazilian Society of Nephrology, which points to hypertension as the major cause of CKD in Brazil³⁸.

The present study showed less obesity compared to a study conducted with older adults, in which 51%²⁵ were overweight. On the other hand, the values were similar with those found in Brazilian Primary Care (31.1%)³⁷. Possibly, the participants of this study, being on dialysis treatment, present weight reduction, because end-stage kidney disease is associated with an increase of inflammatory cytokines and can result in cachexia, with the loss of muscle and fat reserves³⁹.

In relation to the glycaemia and glycated hemoglobin levels, they were found to denote poor glycemic control. A study conducted with 353 people detected a median of 157 mg/dL (58-391 mg/dL) and HbA1C of 8.67% (3.70 - 16.10%). A landmark study has shown that intense diabetes control prevents the progression of diabetes complications³. These data indicate the importance of establishing a multidisciplinary health education plan for better adherence to treatment and minimization of diabetes complications.

When investigating the knowledge about diabetes, the present study corroborates the findings which identified that 66.67% of the participants presented scores ≤ 8 in DKN-A¹⁶, and which found scores ≤ 8 in 64.6% of the participants⁴⁰. Scores below eight indicate insufficient knowledge about the disease.

On the other hand, a study carried out with users of a self-care education program in diabetes¹⁷ found that 78.05% of the participants scored above 8, indicating adequate knowledge and understanding of the disease. Also in a study of

diabetic patients in the Family Health Strategy, with a diabetes patient group²⁷, it was reported that 81.5% of the individuals scored over eight in the DKN-A questionnaire.

It is understood that the last two studies, with positive results regarding knowledge about the disease, are possibly related to the educational initiatives undertaken by the health teams to the diabetic groups. This perception is validated by the findings of studies^{33,41}, which obtained statistically significant differences in knowledge about diabetes, comparing the results before and after an educational program. The same was found by another research study⁴², which concluded that education presents itself as an effective strategy, which contributes for diabetics to live better with their condition. Finally, a systematic review suggests that health education is related to a positive response, when comparing the physiological, psychological and educational parameters, initial and final results of the studies³⁴.

In the DKN-A questionnaire, the questions that obtained the highest number of correct answers are in agreement with the findings of a research study, which found 83.8% of correct answers in the questions related to glucose variation when uncontrolled DM, and 86% in relation to the normal glucose range²⁷. Relative to complications due to DM and behavior towards hypoglycemia, the same study pointed out 67.6% and 60.3% of correct answers, respectively. Correct behavior in case of hypoglycemia is important, as it allows the individuals to return to their routine functions, overcoming the malaise installed due to the occurrence. Regarding the questions with the lowest number of correct answers, they were numbers 6 (on urinary ketones), 12 (causes of hypoglycemia) and 15 (substitution of French bread), with 5.63%, 23.94% and 28.16% of correct answers, respectively. When comparing with another studies, the percentages of correct answers were 39.2%, 27% and 39.9%/12.6%, respectively²⁷. In the same sense, a study conducted with older people found 19.8%, 11.4% and 1%, respectively²⁵. The findings of the present study are consistent with the Brazilian reality, since the urinary ketone test is not a common practice oriented in the health services. The low rate of correct answers regarding the causes of hypoglycemia indicates the lack of understanding that the participants have about their disease, which can impact on the management of the disease, as well as the complications inherent to the health status. Unsatisfactory indicators were also obtained in the group of questions related to DM management and the general principles of care for the disease.

The lack of knowledge about the food groups and the correct substitutions is evident in the results described in Table 1. These findings indicate the urgent need for initiatives in health education with this population. Due to their performance in different spaces, nurses have many opportunities to develop this work.

A number of authors pointed out that the identification of knowledge about DM is a relevant resource to direct the multi-professional team to make clinical decisions and propose a therapeutic plan²⁷. Other authors defend the need to seek new educational strategies, which allow diabetic patients to acquire knowledge for the management of the disease, incorporating it into their daily life, with possibilities to transform their attitudes towards the disease²⁹. A review on educational interventions and glycemic control concluded that the health education practices for the control and treatment of diabetes, around the world, indicate the involvement of health professionals in instrumentalising patients with a view to autonomy and the expansion of self-care, aiming to improve the clinical outcomes and quality of life⁴³.

Regarding attitudes towards the pathology, the present study found results that identified low readiness to face the disease, which corroborates the findings of other publications: 95.56%²⁹; 52.7% in women and 50.7% in men¹⁷, with 93.7%²⁸, 99.4%³² and 85.9% in older people²⁵. The results obtained in ATT-19 are in consonance with studies that identified a mean of 63.23 points³², 67.90 ± 8.0^{44} , 55.5 ± 8.0^{32} and 58.(IQR 52-65) points²⁵. There is evidence that knowledge about the pathology is directly proportional to the attitudes and self-care of diabetic patients on dialysis⁴⁵.

In the result of the ATT-19, it is observed that, although the participants consider that diabetes does not disturb social life and that it is not a problem, many report being ashamed of having the disease and not letting others know about the pathology. In addition, there seems to be some resignation with the disease, when they point out that there is nothing that can be done if you have diabetes, while many consider it unfair to have diabetes and other people have very good health. Although initiatives for patient education are necessary and emphasized by the Brazilian Diabetes Association and by the American Diabetes Association, it is required that health professionals draw strategies to increase patient empowerment, encouraging them to take the lead⁴³, since the expansion of knowledge does not necessarily imply the behavioral changes needed for the control of the disease²⁰.

In this sense, it is believed that the treatment of diabetes requires several self-care activities, and that emotional and cognitive aspects are directly related to the capacity for self-care and self-management of diabetes. However, the positive effects of health education are strongly impaired when the patient suffers from diabetes-related depression or distress. Thus, the health team needs to pay attention to the psychological aspects of the patients and, if necessary,

psychological support must be integrated into the care of these people⁴⁶. The involvement of the patients in their own care strongly contributes to a successful treatment but, often, the encouragement and support of health workers is necessary for the patients to find the motivation to take care of themselves⁴⁷.

Study limitations

Although this study has produced important findings related to the knowledge and attitudes towards diabetes in those who concurrently undergo dialysis, some limitations are recognized. The first concerns the number of participants. Although two dialysis services have been included, there is a need for studies that incorporate a greater number of subjects in order to increase the generalization ability. The second limitation is the study design, cross-sectional; in this sense, prospective studies are necessary, with health education interventions, so that the results can better guide the clinical practice.

CONCLUSION

The participants in this study were shown to have insufficient knowledge about diabetes. This fact has an impact on the ability to self-care, also influenced by the psychological aspects, herein evidenced by the attitudes towards diabetes, which also proved to be unsatisfactory in most of the sample.

Considering that there is a relationship between schooling, knowledge about the disease and health status, it is understandable that the individuals have diverse levels of comprehension about their pathology and, consequently, different coping strategies. Health education processes undertaken in the clinical practice, therefore, should consider the patients' knowledge about the pathology they are carrying, as well as the psychological and emotional aspects related to the disease.

Among the strategies used by nurses, validated questionnaires can support health education as a way of tailoring care, empowering patients and making them co-responsible for their treatment and condition.

REFERENCES

1. Cho NH, Shaw JE, Karuranga S, Huang Y, da Rocha Fernandes JD, Ohlrogge AW, et al. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res. Clin. Pract.* 2018 [cited 2020 Feb 20]; 138:271-81. DOI: <https://doi.org/10.1016/j.diabres.2018.02.023>.
2. Brazil. Vigitel Brazil 2017: surveillance of risk and protective factors for chronic diseases by telephone survey: estimates of frequency and sociodemographic distribution of risk and protective factors for chronic diseases in the capitals of the 26 Brazilian states and the Federal District in 2017. Ministry-of-Health. Brasília. [cited 2020 Feb 20] Available from http://bvsms.saude.gov.br/bvs/publicacoes/vigitel_brasil_2017_vigilancia_fatores_riscos.pdf.
3. Nathan DM, Genuth S, Lachin J, Cleary P, Crofford O, Davis M, et al. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N. Engl. J. Med.* 1993 [cited 2020 Mar 10]; 329(14):977-86. DOI: <https://doi.org/10.1056/NEJM199309303291401>.
4. Writing Group for the DCCT/EDIC Research Group. Coprogression of cardiovascular risk factors in type 1 diabetes during 30 years of follow-up in the DCCT/EDIC study. *Diabetes Care.* 2016 [cited 2020 Mar 7]; 39(9):1621-30. DOI: <https://doi.org/10.2337/dc16-0502>.
5. Nathan DM, Buse JB, Davidson MB, Ferrannini E, Holman RR, Sherwin R, et al. Medical management of hyperglycemia in type 2 diabetes: a consensus algorithm for the initiation and adjustment of therapy: a consensus statement of the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care.* 2009 [cited 2020 Mar 10]; 32(1):193-203. DOI: <https://doi.org/10.2337/dc08-9025>.
6. Bem AF, Kunde J. The importance of glycated hemoglobin determination in the management of chronic complications associated with diabetes mellitus. *J Bras. Patol. Med. Lab.* 2006 [cited 2020 Mar 10]; 42(3). DOI: <https://doi.org/10.1590/S1676-24442006000300007>.
7. Kirsztajn GM, Salgado Filho N, Draibe SA, Netto MVdP, Thome FS, Souza E, et al. Fast Reading of the KDIGO 2012: Guidelines for evaluation and management of chronic kidney disease in clinical practice. *J. Bras. Nefrol.* 2014 [cited 2020 Mar 8]; 63-73. DOI: <https://doi.org/10.5935/0101-2800.20140012>.
8. Eknoyan G, Lameire N, Eckardt K, Kasiske B, Wheeler D, Levin A, et al. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney International.* 2013 [cited 2019 Dez 10]; 3:5-14. DOI: <https://doi.org/10.1038/kisup.2012.75>.
9. Assunção TS, Ursine PGS. A study of factors associated to non-pharmacological treatment delivered by the Family Health Program in Ventosa, Belo Horizonte, to carriers of diabetes mellitus. *Ciênc. Saúde Coletiva.* 2008 [cited 2020 Mar 7]; 13(Suppl 2):2189-97. DOI: <https://doi.org/10.1590/S1413-81232008000900024>.
10. Davies MJ, Heller S, Skinner TC, Campbell MJ, Carey ME, Cradock S, et al. Effectiveness of the diabetes education and self management for ongoing and newly diagnosed (DESMOND) programme for people with newly diagnosed type 2 diabetes: cluster randomised controlled trial. *BMJ.* 2008 [cited 2020 Mar 7]; 336(7642):491-5. DOI: <https://doi.org/10.1136/bmj.39474.922025.BE>.

11. Mascarenhas NB, Pereira A, Silva RSd, Silva MGd. Systematization of nursing assistance to patients with diabetes mellitus and chronic renal. *Rev. Bras. Enferm.* 2011 [cited 2020 Mar 7]; 64:203-8. Available from: https://www.scielo.br/scielo.php?script=sci_abstract&pid=S0034-71672011000100031&lng=en&nrm=iso.
12. Beeney LJ, Dunn SM, Welch G. Measurement of diabetes knowledge: the development of the DKN scales. In: Harwood, Publishers A, eds. *Handbook of psychology and diabetes*. Amsterdam. 2001:159-189.
13. Torres HC, Hortale VA, Schall VT. Validation of Diabetes Mellitus knowledge (DKN-A) and attitude (ATT-19) questionnaires. *Rev. Saúde Pública.* 2005 [cited 2020 Feb 20]; 39(6):906-911. DOI: <https://doi.org/10.1590/S0034-89102005000600006>.
14. Welch G, Dunn SM, Beeney LJ. The ATT39: A measure of psychological adjustment to diabetes. Amsterdam. 2001.
15. Torres HC, Hortale VA, Schall VT. Validation of Diabetes Mellitus knowledge (DKN-A) and attitude (ATT-19) questionnaires. *Rev. Saúde Pública.* 2005 [cited 2020 Feb 20]; 39(6):906-911. DOI: <https://doi.org/10.1590/S0034-89102005000600006>.
16. Rodrigues FFL, Santos MAD, Teixeira CRdS, Gonela JT, Zanetti ML. Relationship between knowledge, attitude, education and duration of disease in individuals with diabetes mellitus. *Acta Paulista de Enfermagem.* 2012 [cited 2020 Feb 20]; 25(2):284-90. DOI: <https://doi.org/10.1590/S0103-21002012000200020>.
17. Rodrigues FFL, Zanetti ML, Santos MAD, Martins TA, Sousa VD, Teixeira CRdS. Knowledge and attitude: important components in diabetes education. *Revista Latino-Americana de Enfermagem.* 2009 [cited 2020 Feb 20]; 17(4):468-73. DOI: <https://doi.org/10.1590/S0104-11692009000400006>.
18. de Oliveira KCS, Zanetti ML. Knowledge and attitudes of patients with diabetes mellitus in a primary health care system. *Revista da Escola de Enfermagem da USP.* 2011 [cited 2020 Feb 20]; 45(4). DOI: <https://doi.org/10.1590/S0080-62342011000400010>.
19. de Oliveira Santos BM, Caixeta ACM, da Silva AA, de Souza Teixeira CR. Knowledge and attitudes in diabetes mellitus type 2: subsidies for self-care and health promotion. *Arquivos de Ciências da Saúde.* 2016 [cited 2020 Feb 20] (4):31-36%V 23. DOI: <https://doi.org/10.17696/2318-3691.23.4.2016.443>.
20. Pereira DA, Costa NMdSC, Sousa ALL, Jardim PCBV, Zanini CRdO. The effect of educational intervention on the disease knowledge of diabetes mellitus patients. *Revista Latino-Americana de Enfermagem.* 2012 [cited 2020 Feb 22]; 20(3):478-85. DOI: <https://doi.org/10.1590/S0104-11692012000300008>.
21. Gonçalves NEXM, Zanetti ML, Neiva CM, Vassimon HS. Knowledge of individuals with diabetes mellitus in the Family Health Strategy. *Revista de Enfermagem UFPE online.* 2017 [cited 2020 Feb 22]; 11(7):2779-2787. DOI: <https://doi.org/10.5205/reuol.10939-97553-1-RV.1107201718>.
22. Hussain Z, Yusoff ZM, Sulaiman SA. A study exploring the association of attitude and treatment satisfaction with glycaemic level among gestational diabetes mellitus patients. *Prim. Care Diabetes.* 2015 [cited 2020 Feb 22]; 9(4):275-82. DOI: <https://doi.org/10.5205/reuol.10939-97553-1-RV.1107201718>.
23. Campos SM, Aparecida DF, Santos Nascimento J, Dos Santos Tavares DM. Correlation of quality of life with knowledge and attitude of diabetic elderly. *Invest. educ. enferm.* 2016 [cited 2020 Feb 22]; 34(1):180-8. DOI: <https://doi.org/10.17533/udea.iee.v34n1a20>.
24. Figueira ALG, Boas LCGV, Coelho ACM, Freitas MCF, Pace AE. Educational interventions for knowledge on the disease, treatment adherence and control of diabetes mellitus. *Rev. Lat. Am. Enfermagem.* 2017 [cited 2020 Mar 9]; 25:e2863. DOI: <https://doi.org/10.17533/udea.iee.v34n1a20>.
25. Borba AKOT, Arruda IKG, Marques APO, Leal MCC, Diniz ADS. Knowledge and attitude about diabetes self-care of older adults in primary health care. *Cien. Saude Colet.* 2019 [cited 2020 Mar 9]; 24(1):125-36. DOI: <https://doi.org/10.17533/udea.iee.v34n1a20>.
26. Brucki SM, Nitrini R, Caramelli P, Bertolucci PH, Okamoto IH. Suggestions for using the mini-mental state exam in Brazil. *Arq. Neuropsiquiatr.* 2003 [cited 2020 Mar 8]; 61(3B):777-81. DOI: <https://doi.org/10.1590/S0004-282X2003000500014>.
27. Gonçalves NEXM, Zanetti ML, Neiva CM, Vassimon HS. *Revista de enfermagem UFPE.* 2017 [cited 2020 Mar 8]; 11(7):2779-2787. DOI: <https://doi.org/10.1590/S0004-282X2003000500014>.
28. Oliveira KCSd, Zanetti ML. Knowledge and attitudes of patients with diabetes mellitus in a primary health care system. *Revista da Escola de Enfermagem da USP.* 2011 [cited 2020 Mar 8]; 45(4):862-8. DOI: <https://dx.doi.org/10.1590/S0080-62342011000400010>.
29. Rodrigues FFL, Santos MAD, Teixeira CRdS, Gonela JT, Zanetti ML. Relationship between knowledge, attitude, education and duration of disease in individuals with diabetes mellitus. *Acta Paulista de Enfermagem.* 2012 [cited 2020 Mar 10]; 25(2):284-90. DOI: <https://doi.org/10.1590/S0103-21002012000200020>.
30. de Oliveira KCS, Zanetti ML. Knowledge and attitudes of patients with diabetes mellitus in a primary health care system. *Revista da Escola de Enfermagem da USP.* 2011 [cited 2020 Mar 10]; 45(4). DOI: <https://doi.org/10.1590/S0080-62342011000400010>.
31. Sesso RC, Lopes AA, Thomé FS, Lugon JR, Martins CT. Brazilian Chronic Dialysis Census 2014. *Jornal Brasileiro de Nefrologia.* 2016 [cited 2020 Mar 10]; 38(1):54-61. DOI: <https://doi.org/10.5935/0101-2800.20170049>.
32. de Oliveira Santos BM, Caixeta ACM, da Silva AA, de Souza Teixeira CR. Knowledge and attitudes towards diabetes mellitus type 2: contributions to self-care and health promotion. *Arquivos de Ciências da Saúde.* 2016 [cited 2020 Mar 10] (4):31-36%V 23. DOI: <https://doi.org/10.17696/2318-3691.23.4.2016.443>.
33. Otero LM, Zanetti ML, Ogrizio MD. Knowledge of diabetic patients about their disease before and after implementing a diabetes education program. *Revista Latino-Americana de Enfermagem.* 2008 [cited 2020 Mar 10]; 16(2):231-7. DOI: <https://doi.org/10.1590/S0104-11692008000200010>.

34. Iquize RCC, Theodoro FCET, Carvalho KA, Oliveira MdA, Barros JdF, Silva ARd. Educational practices in diabetic patient and perspective of health professional: a systematic review. *Jornal Brasileiro de Nefrologia*. 2017 [cited 2020 Mar 10]; 39(2):196-204. DOI: <https://doi.org/10.5935/0101-2800.20170034>.
35. Moreira H, Sette J, Keiralla L, et al. Diabetes mellitus, hypertension and chronic kidney disease: treatment strategies and their limitations. *Rev. Bras. Hipertens*. 2008 [cited 2020 Mar 10]; 15(2):111-6. Available from: <http://departamentos.cardiol.br/dha/revista/15-2/17-diabetes.pdf>.
36. Thomé FS, Sesso RC, Lopes AA, Lugon JR, Martins CT. Brazilian chronic dialysis survey 2017. *J. Bras. Nefrol*. 2019 [cited 2020 Mar 11]; 41(2): 208-14. DOI: <https://doi.org/10.1590/2175-8239-jbn-2018-0178>.
37. Muc R, Saracen A, Grabska-Liberek I. Associations of Diabetic Retinopathy with Retinal Neurodegeneration on the Background of Diabetes Mellitus. Overview of Recent Medical Studies with an Assessment of the Impact on Healthcare systems. *Open Med (Wars)*. 2018 [cited 2020 Mar 11]; 13:130-6. DOI: <https://doi.org/10.1515/med-2018-0008>.
38. Assunção SC, Fonseca AP, Silveira MF, Caldeira AP, Pinho Ld. Knowledge and attitude of patients with diabetes mellitus in Primary Health Care. *Escola Anna Nery*. 2017 [cited 2020 Mar 11]; 21. <http://dx.doi.org/10.1590/2177-9465-ean-2017-0208>.
39. Alvis Zibran M, Mohammadnezhad M. Management of Type 2 Diabetes and Chronic Kidney Disease in Fiji in 2018: Knowledge, Attitude, and Practice of Patients. *Rev. Diabet. Stud*. 2019 [cited 2020 Mar 11]; 15:26-34. DOI: <https://doi.org/10.1900/RDS.2019.15.26>.
40. Oliveira KCsD, Zanetti ML. Knowledge and attitudes of patients with diabetes mellitus in a primary health care system. *Revista da Escola de Enfermagem da USP*. 2011 [cited 2020 Mar 11]; 45(4):862-8. DOI: <https://doi.org/10.1590/S0080-62342011000400010>.
41. De Carvalhotorres H, Nogueiracortez D, Reis IA. Education assessment in diabetes group in primary health care. *Ciencia y Enfermería*. 2016 [cited 2020 Mar 11]; 22:35-45. Available from: <https://scielo.conicyt.cl/pdf/cienf/v22n3/0717-9553-cienf-22-03-00035.pdf>.
42. Vieira GdLC, Cecílio SG, Torres HdC. The perception of users with diabetes regarding a group education strategy for the promotion of self-care. *Escola Anna Nery*. 2017; 21(1). DOI: <https://doi.org/10.5935/1414-8145.20170017>.
43. Capellari C, Costa B, Larré A, Pasin D, Cardoso A, Vasconcelos L, et al. Educational Interventions and Glycemic Control: Integrative Review. *Journal of Diabetes & Metabolism*. 2016 [cited 2020 Mar 11]; 7(7). DOI: <https://doi.org/10.4172/2155-6156.1000675>.
44. Dempster M, McCarthy T, Davies M. Psychological adjustment to Type 2 diabetes and relationship quality. *Diabetic Medicine*. 2011 [cited 2020 Mar 11]; 28(4):487-92. DOI: <https://doi.org/10.1111/j.1464-5491.2010.03214.x>.
45. Ghannadi S, Amouzegar A, Amiri P, Karbalaieifar R, Tahmasebinejad Z, Kazempour-Ardebili S. Evaluating the Effect of Knowledge, Attitude, and Practice on Self-Management in Type 2 Diabetic Patients on Dialysis. *J Diabetes Res*. 2016 [cited 2020 Mar 11]; 2016:3730875. DOI: <https://doi.org/10.1155/2016/3730875>.
46. Schinckus L, Dangois F, Van den Broucke S, Mikolajczak M. When knowing is not enough: Emotional distress and depression reduce the positive effects of health literacy on diabetes self-management. *Patient Education and Counseling*. 2018 [cited 2020 Mar 11]. DOI: <https://doi.org/10.1016/j.pec.2017.08.006>.
47. Wong FKY, Chow SKY, Chan TMF. Evaluation of a nurse-led disease management programme for chronic kidney disease: A randomized controlled trial. *International Journal of Nursing Studies*. 2017 [cited 2020 Mar 11]; 47(3):268-278. DOI: <https://doi.org/10.1016/j.ijnurstu.2009.07.001>.