



Gomphonema Ehrenberg (Bacillariophyceae) in a lotic environment of the Upper Paraná River floodplain, Brazil

Nicoll Cristina Osório^{1*}, Priscila Izabel Tremarin², Thelma Veiga Ludwig² and Liliana Rodrigues¹

¹Laboratório de Algas Perifíticas, Departamento de Biologia Geral, Núcleo de Pesquisa em Limnologia, Ictiologia e Aquicultura, Programa de Pós-Graduação em Ecologia de Ambientes Aquáticos Continentais, Universidade Estadual de Maringá, Avenida Colombo, 5709, 87020-900, Maringá, Paraná, Brazil. ²Laboratório de Ficologia, Departamento de Botânica, Universidade Federal do Paraná, Curitiba, Paraná, Brazil. *Author for correspondence. E-mail: nicoll_cristina@hotmail.com

ABSTRACT. A taxonomic study of species of the genus *Gomphonema* Ehrenberg (Bacillariophyceae) was performed in a lotic environment (Ipoitá Channel) of the Upper Paraná River Floodplain. Samplings were performed in June, September and December 2013 and February 2014. Two petioles of *Eichhornia azurea* (Sw) Kunth were sampled at three different sites in the channel. Twelve species and two taxonomic varieties were identified. All species identified were recorded at the sampling site 3, located close to the Ivinhema River. The lowest number of taxa occurred at the sampling site 1 (9 specific and infra-specific taxa), in the confluence with the Paraná River. All taxa of *Gomphonema* were the first record for the Upper Paraná River Floodplain. Our results emphasize the lack of taxonomic studies for the region and the importance thereof to the knowledge of biodiversity.

Keywords: diatom, distribution area, periphyton, taxonomy.

Gomphonema Ehrenberg (Bacillariophyceae) em ambiente lótico da planície de inundação do Alto Rio Paraná

RESUMO. Realizou-se o estudo taxonômico das espécies do gênero *Gomphonema* Ehrenberg (Bacillariophyceae) em um ambiente lótico (canal Ipoitá) da planície de inundação do alto rio Paraná. As coletas foram realizadas em junho, setembro e dezembro de 2013 e fevereiro de 2014. Dois pecíolos de *Eichhornia azurea* (Sw) Kunth foram amostrados em três pontos distintos do canal. Foram identificadas 12 espécies e duas variedades taxonômicas, sendo que todas foram registradas no ponto 3, situado próximo ao rio Ivinhema. O menor número de táxons ocorreu no ponto 1 (9 táxons específicos e infra-específicos), na área de junção com o rio Paraná. Todos os táxons do gênero *Gomphonema* são primeiro registro para a planície de Inundação do Alto Rio Paraná. Os resultados ressaltam a carência de trabalhos taxonômicos para a região e a importância destes para o conhecimento da biodiversidade.

Palavras-chave: diatomácea, área de distribuição, perifítion, taxonomia.

Introduction

The Upper Paraná River floodplain is characterized by heterogeneous habitats that confer high biodiversity to this environment (Thomaz, Bini, & Bozelli, 2007, Agostinho, Pelicice, & Gomes, 2008). Periphytic algae are one of the aquatic communities found in this ecosystem, with a high proportion of diatoms (Bacillariophyceae), especially in lotic environments. These algae present morphological adaptations that favor their attachment to substrates (Wehr & Sheath, 2003).

The diatom genus *Gomphonema* Ehrenberg is well represented in aquatic environments, presenting high richness and abundance. *Gomphonema* species are characterized by cuneiform

cells in girdle view and heteropolar cells in valve view (Round, Crawford, & Mann, 1990). They generally attach to solid substrates through mucilage pads or stalks, secreted from the apical pore field located at the valve basis (Tremarin, Ludwig, Bertolli, Faria, & Costin, 2009a). In addition, cells possess a single H-shaped plastid with a central pyrenoid (Cox, 1996, Round et al., 1990). Several *Gomphonema* species are cosmopolitan. However, the morphological variability of frustules makes the taxonomy of this genus difficult (Krammer & Lange-Bertalot, 1986, 1991).

Approximately 90 *Gomphonema* species have been described for South America, 28 of which were recently proposed (Reichardt, 1995, Lange-Bertalot, Külbs, Lauser, Nörpel-Schempp, & Willmann, 1996,

Metzeltin & Lange-Bertalot, 1998, Rumrich, Lange-Bertalot, & Rumrich, 2000, Metzeltin, Lange-Bertalot, & Garcia-Rodrigues, 2005, Silva, Nogueira, & Souza, 2011). Approximately 40 of those species have been recorded in Brazil (Torgan, Becker, & Prates, 1999, Tremarin et al., 2009a, Santos, Tremarin, & Ludwig, 2011).

Almost 22 species of *Gomphonema* have been recorded for aquatic environments in Paraná (Tremarin, Freire, Bertolli, & Ludwig, 2009b), and only four species for Mato Grosso do Sul. The present study is the first floristic survey of *Gomphonema* in the Paraná River floodplain, contributing to the knowledge and characterization of aquatic biodiversity in this region, including ecology and biomonitoring studies, for which the correct identification of taxa is essential.

The goal of the present study was to perform a taxonomic analysis of species of the genus *Gomphonema* in a lotic environment of the Upper Paraná River

Floodplain, contributing to the knowledge regarding the distribution of these species.

Material and methods

The study site is the Ipoitá Channel ($22^{\circ} 50' S$; $53^{\circ} 33' W$), located in the Upper Paraná River floodplain, which connects the Ivinhema and Paraná rivers, with an average depth of 3.2 m. Samples were taken at three different sampling sites in this channel: site 1 at the connection with the Paraná River, site 2 in the middle of the channel, and site 3 at the connection with the Ivinhema River (Figure 1). This channel is inserted in the state of Mato Grosso do Sul, although it is practically on the border with the state of Paraná.

The substrates used were petioles of *Eichhornia azurea* (Sw) Kunth in the adult stage (Schwarzbald, 1990), selected from banks of this macrophyte of similar shape and size and under similar environmental conditions.




Figure 1. Upper Paraná River floodplain. The dashed line indicates the Ipoitá Channel.

Collections were performed in June, September and December 2013, and in February 2014. Periphytic algae were collected from the sixth or seventh plant internode of submerged mature petioles of *Eichhorniaazurea* (Sw.) Kunthas recommended by Schwarzbold (1990). This macrophyte was selected to this research because it was the most abundant and is used as a standard plant for the PELD- Long Term Ecological Research. Two petioles were collected randomly at each site, placed in 150 mL Wheaton bottles, sprayed with distilled water, and kept in boxes with ice. The periphytic material was subsequently removed using stainless steel blades wrapped in aluminum foil and jets of distilled water. The resulting material was fixed in 1:1 Transeau solution (Bicudo & Menezes, 2006).

Part of the sample was oxidized with potassium permanganate and hydrochloric acid, according to Simonsen (1974) as modified by Moreira-Filho and Valente-Moreira (1981). Due to the high amount of organic matter in the blades, the material was oxidized again according to Hasle and Fryxell (1970). The resins used to mount permanent slides were Hyrax for the first method and Nafrax for the

second. One slide was mounted per site sampled in certain periods, resulting in 12 slides, in total. These slides were not quantified, but all of them were analyzed for species richness.

Gomphonema species were analyzed using a binocular microscope with amicrometer ocular and 100x objective, and images were captured using an Olympus DP-071 camera. The terminology used for species description followed Round et al. (1990). Taxa identification was based, whenever possible, with the original literature referred to in the text. Samples were deposited in the Herbário of Universidade Estadual de Maringá (HUEM).

To record the occurrence of taxa distribution was conducted a review of literature with taxonomic approach.

Results and discussion

A total 11 species and 3 taxonomic varieties belonging to genus *Gomphonema* were identified (Table 1). These taxa were distributed in one channel of the Upper Paraná River floodplain, located between the states of Mato Grosso do Sul and Paraná (Table 2).

Table 1. Taxa of *Gomphonema* (Gomphonemataceae) recorded in Ipoitá Channel in the Upper Paraná River floodplain. It occurs in different regions (1: connection with Paraná River, 2: middle of the channel; 3: connection with Ivinhema River) and months (Jun: June, Sep: September, Nov: November, 2013; and Feb: February, 2014).




Table 2. Taxa distribution of *Gomphonema* recorded in Ipoitá Channel in the Upper Paraná River floodplain, located between the states of Mato Grosso do Sul and Paraná.

Taxa	Taxa Ocurrence											
	Paraná											
<i>G. affine</i> var. <i>affine</i> Kützing	Caetano (1984), Moreira (1990), Leandrinini (1999), Rodrigues and Bicudo (2001), Cetto, Leandrinini, Felisberto, and Rodrigues (2004), Bigunas (2005), Silva, Tavares, Aquino, and Wengrat (2007), Pires (2013)											
<i>G. angustatum</i> (Kützing) Rabenhorst	Contin (1983), Cecy (1986), Shirata (1986), Contin (1990), Lozovei and Shirata (1990), Cetto et al. (2004), Moro, Bicudo, Melo, and Schmitt (2004), Tremarin et al. (2009a and b), Bertolli, Tremarin, and Ludwig (2010), Bertolli (2010), Faria (2010)											
<i>G. brasiliense</i> Grunow	Ludwig (1987), Lozovei and Shirata (1990), Moreira (1990), Rodrigues (1991), Moro, Garcia, and Oliveira Júnior (1994), Brassac (1999), Atab (2000), Szawka (2001), Cetto et al. (2004), Bigunas (2005), Piccinini (2005), Aquino and Tavares (2006), Silva et al. (2007), Pires (2013), Marquardt, Furstenberger, Chaouiche, Caparica, and Carapunrala (2010)											
<i>G. costei</i> D. Metzeltin & H. Lange-Bertalot	Brassac (1999), Ferrari (2004), Costin (2007), Pires (2013)											
<i>G. lagenula</i> Kützing	Costin (2007), Santos (2007), Pavan (2008), Tremarin et al. (2009a and b), Bertolli et al. (2010), Bertolli (2010), Marquardt et al. (2010), Faria (2010), Silva, Ludwig, Tremarin, and Vercellino (2010), Moresco, Tremarin, Ludwig, and Rodrigues (2011), Pires (2013), Marra (2015)											
<i>G. laticollum</i> Reichardt	Silva et al. (2007), Tremarin et al. (2009a and b), Bertolli et al. (2010), Bertolli (2010), Marquardt et al. (2010), Silva et al. (2010), Bartozek et al. (2013)											

continuation...

... continuation

<i>G. mexicanum</i> Grunow	Piccinini (2005), Costin (2007), Silva et al. (2007)
<i>G. neonasutum</i> H. Lange-Bertalot & E. Reichardt	First record
	Moreira-Filho and Momoli (1963), Moreira-Filho et al. (1973), Cecy, Valente-Moreira, and Hohmann (1976), Lozovei and Luz (1976), Moreira-Filho, Cecy, and Valente-Moreira (1976), Lozovei and Hohmann (1977), Stankiewicz (1980), Contín (1983), Caetano (1984), Cecy (1986), Shirata (1986), Ludwig (1987), Contín (1990), Lozovei and Shirata (1990), Moreira (1990), Moro (1991), Rodrigues (1991), Train (1991), Moro and Fürsternberg (1993), Moro et al. (1994), Tavares (1994), Momoli (1997), Brassac (1999), Leandrin (1999), Atab (2000), Tavares and Valente-Moreira (2000), Rodrigues and Bicudo (2001), Szawka (2001), Bittencourt-Oliveira (2002), Borges, Rodrigues, Pagioto, and Train (2003), Cetto et al. (2004), Ferrari (2004), Train and Rodrigues (2004), Bigunas (2005), Felisberto and Rodrigues (2005), Ludwig, Bigunas, Neiva, Coquemala, and Piccinini (2005), Neiva (2005), Piccinini (2005), Rodrigues et al. (2005), Costin (2007), Santos (2007), Silva et al. (2007), Borges, Train, and Rodrigues (2008), Pavan (2008), Tremarin et al. (2009a and b), Bertolli et al. (2010), Bertolli (2010), Marquardt et al. (2010), Faria (2010), Silva et al. (2010), Fontana and Bicudo (2012), Pires (2013), Marra (2015)
<i>G. parvulum</i> (Kützing) Kützing	Moreira (1990), Brassac (1999), Leandrin (1999), Atab (2000), Ferrari (2004), Bigunas (2005), Piccinini (2005), Costin (2007), Silva et al. (2007), Pavan (2008), Bertolli et al. (2010), Bertolli (2010), Faria (2010), Moresco et al. (2011), Silva et al. (2010), Marra (2015)
<i>G. salae</i> Lange-Bertalot & Reichardt	Tremarin et al. (2009a and b)
<i>G. sphaerophorum</i> Ehrenberg	Faria (2010)
<i>G. subtile</i> Ehrenberg	Train (1990), Moro and Fürsternberg (1993), Tavares (1994), Brassac (1999), Tavares and Valente-Moreira (2000), Rodrigues and Bicudo (2001), Ferrari (2004), Train and Rodrigues (2004), Aquino and Tavares (2006), Santos (2007), Silva et al. (2007), Pavan (2008), Marquardt et al. (2010), Fontana and Bicudo (2012), Marra (2015)
<i>G. turris</i> var. <i>coarctata</i> (Frenguelli) Frenguelli	Tremarin et al. (2009a and b)

Theses and dissertations were included, since they register the species and this information was not published in scientific article.

Identification key for *Gomphonema* species and varieties included the following observations:

1. Presence of stigma 2
1. Absence of stigma *Gomphonema brasiliense*
2. Inconspicuous areolation 3
2. Conspicuous areolation 4
3. Valvelength 35 µm or longer 5
3. Valvelength 26 µm or shorter 6
4. Valve width 10.5 µm or lower 9
4. Valve width 12.5 µm or higher 10
5. Valves slightly lanceolate *Gomphonema affine* var. *affine*
5. Valves clavate *Gomphonema salae*
 6. Valves lanceolate-to-elliptic-lanceolate 7
 6. Othervalve shapes 8
7. Apices subcapitato subrostrate; bases subcapitate *Gomphonema laguna*
7. Apices subrostrate; bases attenuate-rounded *Gomphonema parvulum*
 8. Central area unilateral formed by a shortened median stria *Gomphonema angustatum*
 8. Central area rounded *Gomphonema pumilum*
9. Number of striae in 10 µm 13 or fewer 10
9. Number of striae in 10 µm 14 or more *Gomphonema laticollum*
10. Axial area linear 11
10. Axial area slightly undulate *Gomphonema neonasutum*
11. Valvethin and narrow, lanceolate, with expanded center *Gomphonema subtile*
11. Othervalve shapes 12
12. Raphe sternum wide *Gomphonema costei*
12. Raphe sternum narrow 13
13. Rapheliform *Gomphonema sphaerophorum*
13. Rapheslightly sinuous 14
14. Valvelength longer than 54 µm *Gomphonema turris* var. *coarctata*
14. Valvelength shorter than 35 µm *Gomphonema mexicanum*

Gomphonema affine var. *affine* Kützing, Bacill. Nordhausen. p. 86, pl. 30, fig. 54, 1844 (Figure 2/1-19).

Valves slightly heteropolar, lanceolate, with apices rounded and bases attenuate-rounded. Raphe sternum

narrow and linear. Raphes lightly sinuous with proximal ends slightly bent towards the stigma. Central area is asymmetrical, expanding until the valve margin is on one side of the valve due to the wider spacing of the median striae. Stigma located close to the medianstria. Striae straight to radiate, with one shortened stria on one side of the central nodule. Areola inconspicuous. Length: 35.8-51.7 µm; width: 7.6-10.5 µm; 9-13 striae in 10 µm.

Gomphonema affine is similar to *Gomphonema amoenum* Lange-Bertalot in its dimensions and valve outline. However, *G. amoenum* presents subrostrateapices and more radiate striae (Reichardt, 1999). The results of the analyzed population are in agreement with the morph metric data (length 36-88 µm, width 9-13.6 µm, 8-11 striae in 10 µm) of the tropical species material analyzed by Reichardt (1999).

Occurrence in samples: Huem-24354, 24355, 24356 collected on June 15th, 2013; Huem-24357, collected on August 30th, 2013.

Gomphonema angustatum (Kützing) Rabenhorst, Fl. Eur. Alg. I, p. 283, 1864.

Basionym: *Sphenella angustata* Kützing, Kies. Bacill. Diat., p. 83, pl. 8, fig. 4, 1844 (Figure 2/27-31).

Valves heteropolar, narrow lyclavate-lanceolate, with apices subrostrate and bases attenuate. Raphe sternum linear and narrow. Raphe straight or slightly sinuous, with proxima lents slightly bent towards the stigma. Central area unilateral formed by a shortened median stria. Stigma located close to the median stria. Striae straight, slightly radiate at the ends and with wider spacing at the median region of the valve. Areolae inconspicuous. Length: 16.4-21.1 µm; width: 3.5-4.7 µm; 11-15 striae in 10 µm.




Figure 2. 1-19: *Gomphonema affine* var. *affine*. 20: *Gomphonema brasiliense*. 21-26: *Gomphonema sphaerophorum*. 27-31: *Gomphonema angustum*. 32-38: *Gomphonema lagenula*. 39-42: *Gomphonema costei*. Scale: 10 µm.




Figure 3. *Gomphonema turris* var. *coarctata*. 45-48: *Gomphonema neonasutum*. 49-52: *Gomphonema laticollum*. 53: *Gomphonema pumilum*. 54-60: *Gomphonema parvulum*. 61-63: *Gomphonema subtile*. 64-65: *Gomphonema salae*. 66-73: *Gomphonema mexicanum*. Scale: 10 µm.

This species was found in others studies conducted in the state of Paraná. In Tremarin et al. (2009a), this species was found in the macrophyte *Potamogeton polygonus* and the individuals were larger than observed in our study.

Occurrence in samples: Huem-24354, 24355, 24356 collected on June 15th, 2013; Huem-24361, collected on November 29th, 2013.

Gomphonema brasiliense Grunow in Schneider, Naturw. Beitr. Kenntn. Kauk., p. 110, 1878 (Figure 2/ 20).

Valves slightly heteropolar, lanceolate, with apices cuneate-rounded and bases narrowly rounded. Raphe sternum wide and lanceolate. Raphes lightly sinuous with proxima lents bent towards the stigma. Central area rounded. Stigma absent. Striae short, straight toradiate along the length of the valve. Areolae inconspicuous. Length: 36.4 µm; width: 6.4 µm; 15 striae in 10 µm.

Gomphonema brasiliense ssp. *pacificum* Moser, Lange-Bertalot and Metzeltin can be differentiated from *Gomphonema brasiliense* by its narrower valves (3.6-5.2 µm) (Moser, Lange-Bertalot, & Metzeltin, 1998).

Occurrence in samples: Huem-24354, collected on June 15th, 2013; Huem-24363, collected on February 18th, 2014.

Gomphonema costei Metzeltin and Lange-Bertalot in Lange-Bertalot, Iconogr. Diatomol. 5: 115, pl. 154, fig. 7-12, 1998 (Figure 2/ 39-42).

Valves heteropolar, lanceolate, slightly swollen at the median region, with rounded apices and bases. Raphe sternum lanceolate. Raphe sinuous with proxima lents bent towards the stigma. Stigma rounded located close to the central nodule. Striae straight toradiate at the ends, coarse and shortened long the full length of the valve. Areolae conspicuous, difficult to observe. Length: 41.7-48.8 µm; width: 7.0-8.2 µm; 9-10 striae in 10 µm.

Gomphonema costei and *Gomphonema evexus* Hohn has very similar valve outline as well as identical dimensions and density of striae, suggesting that the species may be synonymized in the future (Patrick et al., 1966).

Occurrence in samples: Huem-24354, 24355, 24356 collected on June 15th, 2013; Huem-24357, collected on August 30th, 2013; Huem-24360, collected on November 29th, 2013; Huem-24365, collected on February 18th, 2014.

Gomphonema lagenula Kützing, Kies. Bacill. Diat., p. 85, pl. 30, fig. 60, 1844 (Figure 2/ 32-38).

Valves heteropolar, lanceolate to elliptic-lanceolate, with apices subcapitate to subrostrate and bases subcapitate. Raphe sternum narrow and linear.

Raphe straight with proxima lents bent towards the stigma. Central area unilaterally expanded, delimited by shortened median striae. Stigma located at the end of the median stria. Striae uniseriate, straight to slightly radiate, with central striae more widely spaced than the others. Areola inconspicuous. Length: 21.1-25.8 µm; width: 5.8-6.4 µm; 14-16 striae in 10 µm.

Occurrence in samples: Huem-24354, 24355, 24356 collected on June 15th, 2013; Huem-24357, 24358, 24359 collected on August 30th, 2013; Huem-24360, 24361, 24362, collected on November 29th, 2013; HUEM-24363, 24364, 24365, collected on February 18th, 2014.

Gomphonema laticollum Reichardt (2001) in Jahn et al., Studies on Diatoms, p. 199, pl. 5, fig. 1-14, 2001 (Figure 3/ 49-52).

Valves clavate, swollen at the median region and with little pronounced constriction between the median region and the apex. Apices widely rounded and bases attenuate-rounded. Raphe sternum narrow and linear. Raphe sinuous with proxima lents dilated in a pore shape, bent towards the stigma. Central area irregular delimited by shortened median striae. Stigma present. Striae uniseriate and radiate, formed by conspicuous areolae. Length: 21.1-25.8 µm; width: 5.8-6.4 µm; 14-16 striae in 10 µm.

Reichardt (2001) reviewed the species *Gomphonema truncatum* Ehrenberg and *G. capitatum* Ehrenberg and proposed *G. laticollum* Reichardt because it presents striae less pronounced constriction close to the valve apices.

Occurrence in samples: Huem-24354, 24355, 24356 collected on June 15th, 2013; Huem-24357, collected on August 30th, 2013.

Gomphonema mexicanum Grunow in Van Heurck, Syn. Diat. Belg., pl. 24; fig. 3, 1880. (Figure 3/ 66-73).

Valves heteropolar, lanceolate, with apices broadly rostrate and bases attenuate-rounded to subcapitate. Raphe sternum narrow and linear. Raphe weakly undulate, with proxima lents dilated in a pore shape. Central area unilaterally expanded, delimited by a shortened median stria. Stigma punctiform located close to the median stria. Striae straight to radiate close to the valve ends. Areola conspicuous, difficult to observe. Length: 20-34.7 µm; width: 8.8-10.5 µm; 12-13 striae in 10 µm.

Gomphonema mexicanum is very similar to *G. affinopsis* Metzeltin, Lange-Bertalot and García-Rodríguez, differentiated by the punctiform shape of the stigma, whereas in *G. affinopsis*, the stigma is elongated (Metzeltin & Lange-Bertalot, 1998).

Occurrence in samples: Huem-24354, collected on June 15th, 2013; Huem-24357, collected on August 30th, 2013.

Gomphonema neonasutum Lange-Bertalot and Reichardt in Lange-Bertalot, Iconogr. Diatomol. 5: 121, pl. 156, figs. 1-4, 1998 (Figure 3/ 45-48).

Valves heteropolar, lanceolate, slightly swollen at the median region, with apices cuneate-apiculate to cuneate-subrostrate and bases attenuate-rounded. Raphe sternum narrow and linear. Raphe sinuous with proxima lents dilated in a pore shape, bent towards the stigma. Central area irregular delimited by shortened median striae. Stigma located close to the median stria. Striae uniserial, radiate. Areola conspicuous. Length: 67-89.4 μm ; width: 14.7-17 μm ; 9-10 striae in 10 μm .

Gomphonema neonasutum is similar to *G. turris* in valve morphology and size but differs in the apiculate shape of the apices (Metzeltin & Lange-Bertalot, 1998).

Occurrence at the samples: Huem-24354, collected on June 15th, 2013; Huem-24357, 24358, collected on August 30th, 2013; Huem- 24361, 24362, collected on November 29th, 2013; Huem-24363, collected on February 18th, 2014.

Gomphonema parvulum (Kützing) Kützing, Spec. Alg., p. 65, 1849. Basionym: *Sphenella parvula* Kützing, Kies. Bacill. Diat., p. 83, pl. 30, fig. 63, 1844 (Figure 3).

Valves heteropolar, lanceolate to elliptic-lanceolate, with apices subrostrate and bases attenuate-rounded. Raphe sternum linear, narrow. Raphe straight to slightly sinuous with proxima lents bent towards the stigma. Central area irregular and narrow, delimited by a shortened median stria. Stigma present. Striae uniserial, straight to slightly radiate at the ends. Areola inconspicuous. Length: 14.1-19.4 μm ; width: 4.7 μm ; 14-17 striae in 10 μm .

Gomphonema micropus Kützing is morphologically similar to *Gomphonema parvulum* var. *parvulum*. However, the two species can be differentiated according to their differently organized apices and striae (Krammer & Lange-Bertalot, 1986, Reichardt, 1999). *Gomphonema micropus* also differs by being longer (19-44 μm) and wider (6.3-9 μm), and by its lower density of striae (11-14 in 10 μm) (Reichardt, 1999).

Occurrence in samples: Huem-24355, 24356, collected on June 15th, 2013; Huem-24357, collected on August 30th, 2013; Huem- 24361, 24362, collected on November 29th, 2013.

Gomphonema pumilum (Grunow) Reichardt and Lange-Bertalot, Nova Hedwigia 53(3-4): 528, pl. 6, figs. 4-11, 1991.

Basionym: *Gomphonema intricatum* Kützing var. *pumila* Grunow in Van Heurck, Syn. Diat. Belg., pl.24, figs. 35-36, 1880 (Figure 3/ 53).

Valves clavate, with apices rounded and bases attenuate. Raphe sternum linear, narrow. Raphe filiform with proxima lents dilated in a pore shape, bent towards the stigma. Central area rounded. Stigma present. Striae straight to radiate. Areola inconspicuous. Length: 19.4 μm ; width: 4.1 μm ; 10 striae in 10 μm .

Occurrence in samples: Huem-24354, 24355, 24356, collected on June 15th, 2013; Huem-24357, 24359, collected on August 30th, 2013; Huem-24360, 24361, 24362, collected on November 29th, 2013; Huem-24363, 24364, 24365, collected on February 18th, 2013.

Gomphonema salae Lange-Bertalot and Reichardt in Lange-Bertalot, Iconogr. Diatomol. 5: 548, pl. 157, figs. 3-5, 1998 (Figure 3/ 64-65).

Valves clavate, swollen at the median region and with slightly pronounced constriction between the median region and the apex. Apices cuneate-subrostrate and bases attenuate-rounded. Raphe sternum slightly linear, narrow. Raphe weakly sinuous with proxima lents bent towards the stigma. Central area unilateral, delimited by a shortened median stria. Stigma located close to the median stria. Striae uniserial, straight to radiate close to the apices, with wider spacing at the median region. Areola inconspicuous. Length: 42.3-48.2 μm ; width: 10-10.5 μm ; 12-13 striae in 10 μm .

Gomphonema neonasutum differs from *G. salae* in size, and the apiculate shape of the apices. Furthermore, the number of striae in *G. salae* is greater than in *G. neonasutum* and *G. turris* (Metzeltin & Lange-Bertalot, 1998).

Occurrence in samples: Huem-24354, 24356, collected on June 15th, 2013.

Gomphonema sphaerophorum Ehrenberg, Ber. Bek. Verh. Königl. Preuss. Akad. Wiss. Berl., p. 78, 1845 (Figure 2/ 21-26).

Valves heteropolar, lanceolate, with apices rostrate-capitate to subrostrate and bases capitate to subcapitate. Raphe sternum linear and narrow. Raphe filiform with proximal ends simple and rounded. Central area unilateral formed by a shortened median stria. Stigma located close to the median stria. Striae slightly radiate. Areolae rounded. Length: 43.5-56.4 μm ; width: 12.9-13.5 μm ; 9-12 striae in 10 μm .

Occurrence in samples: Huem-24354, 24355, 24356 collected on June 15th, 2013; Huem-24363, 24364, 24365, collected on February 18th, 2014.

Gomphonema subtile Ehrenberg, Abh. Königl. Akad. Wiss. Berl. 1841: 416, 1843 (Figure 3/ 61-63).

Valves heteropolar, lanceolate, with apices capitate and bases rounded. Raphe sterno narrow, linear. Raphe straight with proxima lend dilated in a pore shape. Central area unilateral formed by a shortened median stria. Stigma located close to the median stria. Areolae conspicuous but difficult to observe. Length: 46.4-49.4 μm ; width: 7.6-8.2 μm ; 9-10 striae in 10 μm .

Occurrence in samples: Huem-24354, collected on June 15th, 2013; Huem-24360, collected on November 29th, 2013; Huem-24363, 24365, collected on February 18th, 2014.

Gomphonema turris var. *coarctata* (Frenguelli) Frenguelli, Rev. Mus. La Plata, Sec. Bot. 3: 275, 1941.

Basionym: *Gomphonema turris* f. *coarctata* Frenguelli, An. Mus. Nac. Hist. Nat. 4: 423, pl. 4, figs. 35-36, 1933 (Figure 3/ 43-44).

Valves clavate, lanceolate, with apices cuneate-subrostrate and bases attenuate-rounded. Raphe sterno linear, narrow. Raphe slightly sinuous with proximal ends dilated in a pore shape and bent towards the stigma. Central areae elliptic or asymmetric, delimited by irregular shortening of the median striae. Stigma located close to the median stria. Striae uniserial, straight to radiate near the ends. Areolae conspicuous. Length: 54.7-58.2 μm ; width: 15.2-15.8 μm ; 10 striae in 10 μm ; 18-22 areolae in 10 μm .

Occurrence in samples: Huem-24354, 24356 collected on June 15th, 2013.

In this study, only *Gomphonema neonasutum* is the first record to the state of Paraná (Table 1). The species that have been widely distributed was *G. lagunula* and *G. parvulum*. Only *Gomphonema lagunula* Kützing, a species with cosmopolitan distribution, occurred in all sampling sites and in all months.

The Ipoitá Channel is influenced hydrologically by Paraná and Ivinhema rivers. The sampling site 3, located close to the Ivinhema River, was the most species-rich area (Table 2). This can be explained by the characteristics of the Ivinhema River, which has natural hydrodynamics, and because it is located in a protected area where there is still marginal vegetation and many flood areas (Souza Filho & Stevaux, 1997).

The lowest number of taxa occurred in the sampling site 1, at the connection with the Paraná River (Table 2). Paraná River is strongly impacted by dams and has higher water transparency and lower nutrient concentrations (Agostinho et al., 2008; Roberto, Santana, & Thomaz, 2009). The differences between these rivers may help to explain the distribution of species in the floodplain, and even the success of some species in certain

habitats. Fluctuations in the water level of the Ivinhema River are independent from levels in the Paraná River (Souza Filho, Comunello, & Rocha, 2005), and this pattern also contributes to increase habitat heterogeneity among different riverine habitats in the floodplain (Roberto et al., 2009), elevating the biodiversity in this area. Higher periphyton biomasses, for example, are usually found in the Ivinhema River, which carries more phosphate than the Paraná River (Leandrini, Fonseca, & Rodrigues, 2008).

Conclusion

The present study contributes to the knowledge regarding diatom biodiversity in this region, and it provides support to future ecological and biomonitoring studies in the Upper Paraná River Floodplain. The results emphasize the lack of taxonomic studies for the region and the importance thereof to the knowledge of biodiversity.

Acknowledgements

The authors wish to thank the Long Term Ecological Research (*Programa Ecológico de Longa Duração - PELD*) and the National Council for Scientific and Technological Development (*Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq*) for financial support, and the Center for Research in Limnology, Ichthyology and Aquaculture (*Núcleo de Pesquisas em Limnologia, Ictiologia e Aquicultura - Nupélia*) for logistic support during the field work.

References

- Agostinho, A. A., Pelicice, F. M., & Gomes, L. C. (2008). Damns and the fish fauna of the neotropical region: impacts and management related to diversity and fisheries. *Brasilian Journal of Biology*, 68(6), 1119-1132.
- Aquino, N. F., & Tavares, B. (2006). Levantamento das diatomáceas de um lago marginal do reservatório de Itaipu, rio São Francisco Falso, município de Santa Helena, Paraná, Brasil. In *Encontro Anual de Iniciação Científica*. Ponta Grossa, PR: UEPG.
- Atab, D. R. (2000). *Diatomáceas (Bacillariophyta) de rios das bacias do Tibagi e do Itararé, centro-leste do estado do Paraná, Brasil* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Bartozek, E., Bueno, N. C., Ludwig, T. A. V., Tremarin, P., Nardelli, M., & Rocha, A. (2013). Diatoms (Bacillariophyceae) of Iguaçu National Park, Foz do Iguaçu, Brazil. *Acta Botânica Brasílica*, 27(1), 108-123.
- Bertolli, L. M. (2010). *Diatomáceas perifíticas em substratos natural e artificial, reservatório do rio Passáuá, região metropolitana de Curitiba, Paraná* (Dissertação de

- Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Bertolli, L. M., Tremarin, P. I., & Ludwig, T. A. V. (2010). Diatomáceas perifíticas em *Polygonum hydropiperoides* Michaux, reservatório do Passaúna, Região Metropolitana de Curitiba, Paraná, Brasil. *Acta Botânica Brasílica*, 24(4), 1065-1081.
- Bicudo, C. E. M., & Menezes, M. (2006). Técnicas para coleta, fixação, preservação e estudo. In C. E. M. Bicudo, & M. Menezes (Eds.). *Gênero de algas de águas continentais do Brasil. Chave para identificação e descrição* (2 ed., p. 7-14). São Carlos, SP: Rima.
- Bigunas, P. I. T. (2005). *Diatomáceas (Ochrophyta) do rio Guaraguá, litoral do Paraná, Brasil* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Bittencourt-Oliveira, M. C. (2002). A comunidade fitoplanctônica do rio Tibagi: uma abordagem preliminar de sua diversidade. In M. E. Medri, E. Bianchini, O. A. Shibatta, & J. A. Pimenta (Eds.), *A bacia do rio Tibagi* (p. 373-402). Londrina, PR.
- Borges, P. A. F., Rodrigues, L. C., Pagioro, T. A., & Train, S. (2003). Spatial variation of phytoplankton and some abiotic variables in the Pirapó River – PR (Brasil) in August 1999: a preliminary study. *Acta Scientiarum. Biological Sciences*, 25(1), 1-8.
- Borges, P. A. F., Train, S., & Rodrigues, L. C. (2008). Estrutura do fitoplâncton, em um curto período de tempo, em um braço do reservatório de Rosana (ribeirão do Corvo, Paraná, Brasil). *Acta Scientiarum. Biological Sciences*, 30(1), 57-65.
- Brassac, N. M. (1999). *Diatomoflórlula dos rios da área de abrangência da usina hidrelétrica de Salto Caxias, Bacia do Iguaçu, Paraná* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Caetano, Z. (1984). *Diatomáceas (Bacillariophyta) dos lagos do colégio Santa Maria, município de Almirante Tamandaré, estado do Paraná, Brasil* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Cecy, I. I. T. (1986). Estudo das algas microscópicas (Nostocophyta, Euglenophyta, Chrysophyta e Chlorophyta) do Lago do Parque Barigüi, em Curitiba, estado do Paraná, Brasil. *Arquivos de Biologia e Tecnologia*, 29(2), 383-405.
- Cecy, I. I. T., Valente-Moreira, I. M., & Hohmann, E. (1976). Estudo ficológico e químico da água do tanque do Passeio Público de Curitiba, estado do Paraná – Brasil. *Boletim do Museu Botânico Municipal*, (25), 1-37.
- Cetto, J. M., Leandrini, J. A., Felisberto, S. A., & Rodrigues, L. (2004). Comunidade de algas perifíticas no reservatório de Irai, Estado do Paraná, Brasil. *Acta Scientiarum. Biological Sciences*, 26(1), 1-7.
- Contin, L. F. (1983). *Contribuição ao estudo das diatomáceas (Chrysophyta, Bacillariophyceae) na região da barragem de captação d'água do rio Iguaçu (Sanepar), em Curitiba, Estado do Paraná, Brasil* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Contin, L. F. (1990). Contribuição ao estudo das diatomáceas (Chrysophyta, Bacillariophyceae) na região da barragem de captação d'água do rio Iguaçu (Sanepar), em Curitiba, estado do Paraná, Brasil. *Estudos de Biologia*, (24), 5-95.
- Costin, J. C. (2007). *Diatomáceas (Ochrophyta) epilíticas do rio Negro, Paraná: estrutura da comunidade antes e após um derramamento de óleo diesel e vegetal* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Cox, E. J. (1996). *Identification of freshwater diatoms from live material*. London, UK: Chapman & Hall.
- Faria, D. M. (2010). *Diatomáceas perifíticas de um reservatório eutrófico do rio Itaqui: aspectos qualitativos e quantitativos* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Felisberto, S. A., & Rodrigues, L. C. (2005). Comunidade de algas perifíticas em reservatórios de diferentes latitudes. In L. Rodrigues, S. M. T. Thomaz, A. A. Agostinho, & L. C. Gomes (Orgs.), *Biocenoses em reservatórios – padrões espaciais e temporais* (p. 97-114). São Carlos, SP: Rima.
- Ferrari, F. (2004). *Diatomoflórlula (Ochrophyta) dos rios Ivaí, São João e dos Patos, Bacia Hidrográfica do rio Ivaí (alto curso), Prudentópolis, Paraná* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Fontana, L., & Bicudo, D. C. (2012). Biodiversidade e distribuição das diatomáceas (Bacillariophyceae) de sedimentos superficiais nos reservatórios em cascata do rio Paranapanema, SP/PR, Brasil. *Hoehnea*, 39(4), 587-614.
- Hasle, G. R., & Fryxell, G. A. (1970). Diatoms: cleaning and mounting for light and electron microscopy. *Transactions of the American Microscopical Society*, 89(4), 469-474.
- Krammer, K., & Lange-Bertalot, H. (1986). Bacillariophyceae: Naviculaceae. In H. Ettl, J. Gerloff, H. Heyning, & D. Mollenhauer (Eds.), *Süßwasserflora von Mitteleuropa, Band 2/1* (p. 1-876). Stuttgart, GER: Gustav Fischer Verlag.
- Krammer, K., & Lange-Bertalot, H. (1991). Bacillariophyceae: Achnanthaceae. Kritische Ergänzung zu *Navicula* (Lineolatae) und *Gomphonema*. In H. Ettl, G. Gärtner, J. Gerloff, H. Heyning, & D. Mollenhauer (Eds.), *Süßwasserflora von Mitteleuropa* (p. 12-437). Stuttgart, GE: Gustav Fischer Verlag.
- Lange-Bertalot, H., Külbs, K., Lauser, T., Nörpel-Schempp, M., & Willmann, M. (1996). Diatom taxa introduced by Georg Krasske – documentation and revision. In H. Lange-Bertalot (Ed.), *Iconographia Diatomologica. Annotated diatoms micrographs* (p. 3-358). Königstein, GE: Koeltz Scientific Books.
- Leandrini, J. A. (1999). *Diatomáceas perifíticas da região de captação de água da SANEPAR, Maringá, Paraná – córrego Sarandi e rio Pirapó* (Dissertação de Mestrado). Setor de

- Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Leandrin, J. A., Fonseca, I. A., & Rodrigues, L. (2008). Characterization of habitats based on algal periphyton biomass in the upper Paraná River floodplain, Brazil. *Brazilian Journal of Biology*, 68(3), 503-509.
- Lozovei, A. L., & Luz, E. (1976). Diptera Culicidae em Curitiba e arredores. *Arquivos de Biologia e Tecnologia*, 19(1), 43-83.
- Lozovei, A. L., & Hohmann, E. (1977). Principais gêneros de microalgas em biótopos de larvas de mosquitos de Curitiba, estado do Paraná, Brasil: III – levantamento e constatação de ecologia. *Arquivos de Biologia e Tecnologia*, 19(1-4), 123-151.
- Lozovei, A. L., & Shirata, M. T. (1990). Diatomáceas (Chrysophyta, Bacillariophyceae) no rio Passaúna, Curitiba, Paraná, Brasil - Levantamento qualitativo da diatomoflórida em segmento manancial. *Estudos de Biologia*, 27(1), 5-56.
- Ludwig, T. A. V. (1987). *Diatomoflórida do parque regional do Iguaçu, Curitiba, Paraná* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Ludwig, T. A. V., Bigunas, P. I. T., Neiva, T. F., Coquemala, V., & Piccinini, C. (2005). Diatomáceas (Ochrophyta) dos lagos do Jardim Botânico. In R.C., Pereira (Ed.), *Anais da 10ª Reunião Brasileira de Ficologia* (10 ed., p. 301-323). Rio de Janeiro, RJ: Série Livros.
- Marquardt, G. C., Furstenberger, C. B., Chaouiche, T. E., Caparica, R., & Carapunarla, L. (2010). Diatomáceas (Bacillariophyceae) perifíticas em substratos naturais do rio das Pedras, município de Guarapuava, Paraná, Brasil. *Terr@ Plural*, 4(2), 217-240.
- Marra, R. C. (2015). *Diatomáceas bioindicadoras do estado trófico de reservatórios paranaenses: Piraquara II e Iraí* (Dissertação de Mestrado). Universidade Federal do Paraná, Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Curitiba.
- Metzeltin, D., & Lange-Bertalot, H. (1998). Tropical diatoms of South America. Diversity, taxonomy, geobotany. In H. Lange-Bertalot (Ed.), *Iconographia Diatomologica* (5th ed., p. 1-220). Königstein, GE: Koeltz Scientific Books.
- Metzeltin, D., Lange-Bertalot, H., & Garcia-Rodrigues, F. (2005). Diatoms of Uruguay. In H. Lange-Bertalot (Ed.), *Iconographia Diatomologica* (15th ed., p. 1-173). Königstein, GE: Koeltz Scientific Books.
- Momoli, D. M. M. (1997). Contribuição ao estudo das diatomáceas do Tanque de Senegaglia, São José dos Pinhais, estado do Paraná, Brasil. In *Congresso da Sociedade de Botânica do Brasil* (p. 33-46). Porto Alegre, RS: UFRGS.
- Moreira, A. L. O. R. (1990). Estudo taxonômico de *Cymbella* Agardh e *Gomphonema* Ehrenberg da região de captação de água do rio Pirapó, Maringá, Paraná, Brasil (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Moreira-Filho, H., & Momoli, D. M. M. (1963). Diatomáceas no trato digestivo do *Australorbis glabratus* (Say, 1818). *Boletim da Universidade Federal do Paraná*, 9, 1-7.
- Moreira-Filho, H., & Valente-Moreira, I. M. (1981). Avaliação taxonômica e ecológica das diatomáceas (Bacillariophyceae) epífitas em algas pluricelulares obtidas nos litorais dos estados do Paraná, Santa Catarina e São Paulo. *Boletim do Museu Botânico Municipal*, 47(47), 1-17.
- Moreira Filho, H., Valente-Moreira, I. M., & Cecy, I. T. (1973). Diatomáceas na barragem de captação d'água (Sanepar) do Rio Iguaçu, em Curitiba, Estado do Paraná. *Acta Biologica Paranaense*, 2(1-4), 133-145.
- Moreira-Filho, H., Cecy, I. I. T., & Valente-Moreira, I. M. V. (1976). Diatomáceas da lagoa Dourada, estado do Paraná, Brasil. *Tribuna Farmacéutica*, 44(1), 1-14.
- Moresco, C., Tremarin, P. I., Ludwig, T. A. V., & Rodrigues, L. (2011). Diatomáceas perifíticas abundantes em três córregos com diferentes ações antrópicas em Maringá, PR, Brasil. *Brazilian Journal of Botany*, 34(1), 359-373.
- Moro, R. S., & Fürsternberg, C. B. (1993). Diatomáceas (Bacillariophyceae) da Lagoa Dourada (Parque Estadual de Vila Velha), Paraná, Brasil. *Acta Biológica Paranaense*, 22(1-4), 15-30.
- Moro, R. S., Bicudo, C. E. M., Melo, M. S., & Schmitt, J. (2004). Paleoclimate of the late pleistocene and holocene at Lagoa Dourada, Paraná State, Southern Brazil. *Quaternary International*, 114(1), 87-99.
- Moro, R. S., Garcia, E., & Oliveira Júnior, H. F. (1994). Diatomáceas (Bacillariophyceae) da represa Alagados, Ponta Grossa, Paraná, Brasil. *Iheringia. Série Botânica*, 45(1), 5-19.
- Moro, R. S. (1991). Morphology of *Aulacoseira granulata* (Ehr.) Simonsen var. *australiensis* (Grunow) nov. comb. under light microscopy. *Arquivos de Biologia e Tecnologia*, 34(2), 353-359.
- Moser, G., Lange-Bertalot, H., & Metzeltin, D. (1998). Insel der endemiten. Geobotanisches phänomen neukaledonien. *Bibliotheca Diatomologica*, 38, 1-464.
- Neiva, T. F. (2005). *Diatomáceas briofíticas em Sphagnum L. spp. e Rhacocarpus inermis (C. Muell.) Lingb* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Patrick, R., Aldrich, F. A., Cairns, Jr., J., Drouet, F., Hohn, M. H., Roback, S. S., ... Whitford, L. A. (1966). *The Catherwood Foundation Peruvian-Amazon expedition: limnological and systematic studies*. Monographs of the Academy of Natural Sciences of Philadelphia, Philadelphia.
- Pavan, G. (2008). *Diatomáceas perifíticas em Potamogeton montevidensis Arth. Bennett (Potamogetonaceae) em ambientes lótico e lêntico* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Piccinini, C. (2005). *Diatomáceas perifíticas (Ochrophyta) do rio Gonçalves Dias, Parque Nacional do Iguaçu, Paraná*,

- Brasil (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Pires, E. C. C. (2013). Diatomáceas de rios da bacia hidrográfica litorânea do Paraná, Brasil: *Cymbellaceae* e *Gomphonemataceae* (Dissertação de Mestrado). Universidade Federal do Paraná, Conselho Nacional de Desenvolvimento Científico e Tecnológico, Curitiba.
- Reichardt, E. (1995). Die diatomeen (Bacillariophyceae). *Iconographia Diatomologica*, 1, 1-49.
- Reichardt, E. (1999). Zur revision der gattung *Gomphonema*. *Iconographia Diatomologica*, 8, 1-203.
- Reichardt, E. (2001). Revision der Arten um *Gomphonema truncatum* und *G. capitatum*. In R. Jahn, J. P. Kocielek, A. Witkowski, & P. Compère (Eds.), *Studies on diatoms* (p. 187-224). Koenigstein, GER: Koeltz Scientific Books.
- Roberto, M. C., Santana, N. F., & Thomaz, S. M. (2009). Limnology in the Upper Paraná River floodplain: large-scale spatial and temporal patterns, and the influence of reservoirs. *Brazilian Journal of Biology*, 69(2), 717-725.
- Rodrigues, L. (1991). Naviculaceae (Bacillariophyceae) nas lagoas do horto florestal Dr. Luiz Teixeira Mendes, município de Maringá, Paraná, Brasil. *Unimar*, 13(2), 273-298.
- Rodrigues, L. C., Train, S., Pivato, B. M., Bovo, V. M., Borges, P. A. F., & Jati, S. (2005). Assemblées fitoplanctônicas de trinta reservatórios do Estado do Paraná. In L. Rodrigues, S. M. Thomaz, A. A. Agostinho, & L. C. Gomes (Orgs.), *Biocenose sem reservatórios – padrões espaciais e temporais* (p. 57-72). São Carlos, SP: Rima.
- Rodrigues, L., & Bicudo, D. C. (2001). Similarity among periphyton algal communities in a lentic-lotic gradient of the upper Paraná River floodplain, Brazil. *Revista Brasileira de Botânica*, 24(3), 235-248.
- Round, F. E., Crawford, R. M., & Mann, D. G. (1990). *The Diatoms: biology and morphology of the Genera*. Cambridge, UK: University Press.
- Rumrich, U., Lange-Bertalot, H., & Rumrich, M. (2000). Diatomeen der Anden. Von Venezuela bis Patagonien/Tierra del Fuego. In H. Lange-Bertalot (Ed.). *Iconographia Diatomologica*, Koeltz Scientific Books (p. 9-673). Königstein, GER: Band.
- Santos, E. M. (2007). Diatomáceas perifíticas (*Ochrophyta*) associadas à *Potamogeton polygonus* Chamess. & Schltl. (*Potamogetonaceae*): taxonomia e formas de fixação (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Santos, E. M., Tremarin, P. I., & Ludwig, T. A. V. (2011). Periphytic diatoms on *Potamogeton polygonus* Cham. & Schltl.: first records from Paraná State. *Biota Neotropica*, 11(3), 303-315.
- Schwarzbold, A. (1990). Métodos ecológicos aplicados ao estudo do perifiton. *Acta Limnologica Brasiliensis*, 3(1), 545-592.
- Shirata, M. T. (1986). *Contribuição ao estudo das diatomáceas (Bacillariophyceae) no lago do Parque São Lourenço, Curitiba, estado do Paraná, Brasil* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Silva, A. M., Ludwig, T. A. V., Tremarin, P. I., & Vercellino, I. S. (2010). Diatomáceas perifíticas em um sistema eutrófico brasileiro (Reservatório do Iraí, estado do Paraná). *Acta Botanica Brasilica*, 24(4), 997-1016.
- Silva, A. M., Tavares, B., Aquino, N. F., & Wengrat, S. (2007). Gomphonemaceae (Bacillariophyceae) do rio São Francisco Falso, Estado do Paraná, Brasil. *Revista Brasileira de Biociências*, 5(2), 306-308.
- Silva, W. J., Nogueira, I. S., & Souza, M. G. M. (2011). Catálogo de diatomáceas da região Centro-Oeste brasileira. *Iheringia, Série Botânica*, 66(1), 61-86.
- Simonsen, R. (1974). The diatom plankton of the Indian Ocean Expedition of R/V -Meteor, Meteor Forschungsergebnisse. Reihe D-Biologie, 19, 1-66.
- Souza Filho, E. E., & Stevaux, J. C. (1997). Geologia e geomorfologia do complexo rio Baia, Curutuba, Ivinheima. In A. E. A. M. Vazzoler, A. A. Agostinho, & N. S. Hahn (Eds.), *A planície de inundação do Alto Rio Paraná* (p. 1-46). Maringá, PR: Eduem/Nupélia.
- Souza Filho, E. L., Comunello, E., & Rocha, P. C. (2005). Flood extension in the Baía-Curutuba-Ivinheima complex of the Paraná River floodplain. In A. A. Agostinho, L. Rodrigues, L. C. Gomes, S. M. Thomaz, & S. L. Miranda (Eds.). *Structure and functioning of the upper Paraná River and its floodplain* (p. 19-24). Maringá, PR: Eduem.
- Stankiewicz, E. H. (1980). *Flórula no conteúdo estomacal do Pseudocurimata gilberti (Quoy & Gaimard, 1824)* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Szawka, C. M. (2001). Estrutura e dinâmica espacial e temporal da comunidade fitoplanctônica de reservatório da usina hidrelétrica de Salto Caxias, Paraná, Brasil (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Tavares, B. (1994). *Diatomoflórula no lago artificial de Cascavel, município de Cascavel, estado do Paraná, Brasil* (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Tavares, B., & Valente-Moreira, I. M. (2000). Diatomoflórula do lago de Cascavel, município de Cascavel, estado do Paraná, Brasil. *Hoehnea*, 27(1), 1-24.
- Thomaz, S. M., Bini, L. M., & Bozelli, L. R. (2007). Floods increase similarity among aquatic habitats in river-floodplain systems. *Hydrobiologia*, 579, 1-13.
- Torgan, L. C., Becker, V., & Prates, H. M. (1999). Checklist das diatomáceas (Bacillariophyceae) de ambientes de águas continentais e costeiras do estado do Rio Grande do Sul. *Iheringia*, 52(2), 21-162.
- Train, S. (1990). *Diatomoflórula do córrego dos Moscados, município de Maringá, estado do Paraná, Brasil*

- (Dissertação de Mestrado). Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba.
- Train, S. (1991). Diatomáceas (Bacillariophyceae) do Córrego Moscados, Maringá, Paraná: 1. Bacillariaceae. *Unimar*, 13(2), 313-326.
- Train, S., & Rodrigues, L. C. (2004). Phytoplanktonic assemblages. In S. M. Thomaz, A. A. Agostinho, & N. S. Hahn (Eds.). *The upper Paraná river and its floodplain: physical aspects, ecology and conservation* (p. 103-124). Leiden, NED: Backhuys Publishers.
- Tremarin, P. I., Freire, E. G., Bertolli, L. M., & Ludwig, T. A. V. (2009b). Catálogo das diatomáceas (Ochrophyta-Diatomeae) continentais do estado do Paraná. *Iheringia: Série Botânica*, 64(2), 79-107.
- Tremarin, P. I., Ludwig, T. A. V., Bertolli, L. M., Faria, D. M., & Costin, J. C. (2009a). *Gomphonema Ehrenberg* and *Gomphosphenia Lange-Bertalot* (Bacillariophyceae) from Maurício river, Paraná, Brazil. *Biota Neotropica*, 4(4), 1-9.
- Wehr, J. D., & Sheath, R. G. (2003). Freshwater habitats of algae. In J. D. Wehr, & R. G. Sheath (Eds.). *Freshwater algae of North America: Ecology and classification* (p. 1-917). California: Academic Press.

Received on June 1, 2016.

Accepted on October 28, 2016.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.