


Journal Pre-proof

ESCMID COVID-19 guidelines: diagnostic testing for SARS-CoV-2

Paraskevi C. Fragkou, Giulia De Angelis, Giulia Menchinelli, Fusun Can, Federico Garcia, Florence Morfin-Sherpa, Dimitra Dimopoulou, Elisabeth Mack, Adolfo de Salazar, Adriano Grossi, Theodore Lytras, Chrysanthi Skevaki



PII: S1198-743X(22)00084-2

DOI: <https://doi.org/10.1016/j.cmi.2022.02.011>

Reference: CMI 2837

To appear in: *Clinical Microbiology and Infection*

Received Date: 4 November 2021

Revised Date: 3 February 2022

Accepted Date: 6 February 2022

Please cite this article as: Fragkou PC, De Angelis G, Menchinelli G, Can F, Garcia F, Morfin-Sherpa F, Dimopoulou D, Mack E, de Salazar A, Grossi A, Lytras T, Skevaki C, ESCMID COVID-19 guidelines: diagnostic testing for SARS-CoV-2, *Clinical Microbiology and Infection*, <https://doi.org/10.1016/j.cmi.2022.02.011>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved.

ESCMID COVID-19 Guidelines: Diagnostic testing for SARS-CoV-2

Paraskevi C. Fragkou^{1,2}, Giulia De Angelis^{3,4*}, Giulia Menchinelli^{3,4*}, Fusun Can^{5,6#}, Federico Garcia^{7,8#}, Florence Morfin-Sherpa^{9#}, Dimitra Dimopoulou^{2,10}, Elisabeth Mack¹¹, Adolfo de Salazar^{7,8}, Adriano Grossi¹², Theodore Lytras¹³, Chrysanthi Skevaki^{2,14}

* *equal contribution (in alphabetical order)*

equal contribution (in alphabetical order)

Affiliations

¹ First Department of Critical Care Medicine & Pulmonary Services, Evangelismos General Hospital, National and Kapodistrian University of Athens, Athens, Greece.

² European Society of Clinical Microbiology and Infectious Diseases (ESCMID) Study Group for Respiratory Viruses (ESGREV).

³ Dipartimento di Scienze Biotechnologiche di Base, Cliniche Intensivologiche e Perioperatorie, Università Cattolica del Sacro Cuore, Rome, Italy.

⁴ Dipartimento di Scienze di Laboratorio e Infettivologiche, Fondazione Policlinico Universitario A. Gemelli IRCCS – 00168 Rome, Italy..

⁵ Department of Medical Microbiology, Koc University School of Medicine, Istanbul, Turkey

⁶ Koc University IsBank Research Center for Infectious Diseases (KUISCID), Istanbul, Turkey.

⁷ Servicio de Microbiología Clínica. Hospital Universitario Clínico San Cecilio. Instituto de Investigación Biosanitaria, Granada, Spain.

⁸ CIBER de Enfermedades Infecciosas, CIBERINFEC, ISCIII, Madrid, Spain

⁹ Laboratory of Virology, Institut des Agents Infectieux, National Reference Centre for respiratory viruses, Hospices Civils de Lyon, Université Claude Bernard Lyon1, Lyon, France¹⁰

Second Department of Paediatrics, “P. and A. Kyriakou” Children's Hospital, National and Kapodistrian University of Athens, Athens, Greece.

¹¹Department of Hematology, Oncology and Immunology, University Hospital Giessen and Marburg Campus Marburg and Faculty of Medicine, Philipps University Marburg, Marburg, Germany.

¹²Sezione di Igiene, Istituto di SanitàPubblica, Università Cattolica del Sacro Cuore, Rome, Italy.

¹³School of Medicine, European University Cyprus, Nicosia, Cyprus.

¹⁴ Institute of Laboratory Medicine, Universities of Giessen and Marburg Lung Center (UGMLC), Philipps University Marburg, German Center for Lung Research (DZL), Marburg, Germany.

Corresponding Author

PD Dr. Chrysanthi Skevaki, Institute of Laboratory Medicine and Pathobiochemistry, Molecular Diagnostics, Philipps University Marburg Baldingerstr, 35043 Marburg

Tel: +49 6421 5863850, Fax: +49 6421 5865594

Chrysanthi.Skevaki@uk-gm.de

1 **Abstract**

2 **Scope:** The objective of these guidelines is to identify the most appropriate diagnostic test
3 and/or diagnostic approach for SARS-CoV-2. The recommendations are intended to provide
4 guidance to clinicians, clinical microbiologists, other health care personnel, and decision
5 makers.

6 **Methods:** An ESCMID COVID-19 guidelines task force was established by the ESCMID
7 Executive Committee. A small group was established, half appointed by the chair, and the
8 remaining selected with an open call. Each panel met virtually once a week. For all decisions,
9 a simple majority vote was used. A list of clinical questions using the PICO (population,
10 intervention, comparison, outcome) format was developed at the beginning of the process. For
11 each PICO, two panel members performed a literature search focusing on systematic reviews
12 with a third panelist involved in case of inconsistent results. Quality of evidence assessment
13 was based on the GRADE-ADOLOPMENT approach.

14 **Questions addressed by the guideline and recommendations:** A total of 43 PICO questions
15 were selected that involve the following types of populations: 1) patients with signs and
16 symptoms of COVID-19; 2) travelers, healthcare workers, and other individuals at risk for
17 exposure to SARS-CoV-2; 3) asymptomatic individuals and 4) close contacts of patients
18 infected with SARS-CoV-2. The type of diagnostic test (commercial rapid nucleic acid
19 amplification tests, and rapid antigen detection), biomaterial, time since onset of
20 symptoms/contact with an infectious case, age, disease severity, and risk of developing severe
21 disease are also taken into consideration.

22 **Keywords:** SARS-CoV-2, COVID-19, diagnosis, guidelines, testing.

23

Scope

The present guideline has the objective of identifying the most appropriate diagnostic test and/or diagnostic/screening approach for 1) patients with signs and symptoms of COVID-19; 2) travelers from areas with low and high COVID-19 prevalence, healthcare workers, and other individuals at risk for exposure to SARS-CoV-2; 3) asymptomatic individuals (including general population) and 4) those with close contact with a person infected with SARS-CoV-2; and 5) symptomatic individuals following re-infection and/or vaccination. However, evidence for re-infection and post-vaccination testing approach was scarce when the literature search for the index guidelines was performed. Hence, associated PICOs could not be addressed. Additional considerations include the type of biomaterial, time since onset of symptoms/contact with infectious case, age, disease severity, and risk of developing severe disease. The document is intended to provide guidance to clinicians, clinical microbiologists, other health care personnel, and decision makers.

Context

The ongoing COVID-19 pandemic has severely disrupted human life worldwide and represents an unprecedented challenge to public health [1]. As stated in the first paper of the newly-developed ESCMID guidelines for COVID-19 [2], ESCMID did not develop recommendations at the start of the pandemic. However, given the time passed and the rapid growth in evidence upon which to base recommendations, in January 2021 the ESCMID Executive Committee (EC) decided to launch a new initiative to develop ESCMID guidelines on several COVID-19-related issues. The present guideline provides recommendations for diagnostic testing for SARS-CoV-2, which is of particular relevance given the high incidence of both infections and deaths. This is further complicated by circulating mutations of SARS-

CoV-2 and potential implications for diagnostic testing. Thus, a smart and effective approach to testing and screening is required to minimize spread of the virus.

Consensus guideline development

The general principles and methodology adopted have been described in the first paper in the ESCMID guidelines for COVID-19 related clinical topics [2]. Herein, the panel members proposed a list of diagnostic PICO questions to the panel Chair, who in turn selected the most clinically relevant questions and compiled a set of 55 priority PICOs that reached consensus within the panel. Considering the available evidence, a total of 43 PICOs were used for the development of the present guidelines. The PICOs that were excluded are listed in Supplementary Appendix 1 along with additional details on the methods used.

Quality of evidence assessment

The quality of the body of evidence was evaluated following the GRADE approach. Accordingly, five factors for rating down the quality of evidence (risk of bias, inconsistency of results, indirectness of evidence, imprecision, and publication bias) and three for rating it up (large magnitude of effect, direction of plausible confounding, dose-response gradient) were assessed. Adaptation of the quality assessment factors was applied by the panel for the Diagnostic Guidelines, as described below.

Risk of bias

The risk of bias of each study was assessed using the QUADAS-2 tool for diagnostic accuracy studies. The QUADAS-2 tool includes four key domains that discuss patient selection, index test, reference standard, and flow of patients through the study and timing of the index tests and reference standard. The QUADAS-2 assessment of studies included in the original systematic review was adopted, whereas two panel members independently assessed the

QUADAS-2 for any new study retrieved during the evidence syntheses update, if performed (see Supplementary Methods). The risk of bias was judged very serious or serious if all or more/equal than a half of studies, respectively, have high or unclear concern regarding all or one to three QUADAS-2 domains. The risk of bias was judged not serious in any other case.

Indirectness

Indirectness was judged for all studies based on QUADAS-2 tool concerns for applicability for patient selection, index test, and reference standard. The QUADAS-2 applicability assessment of studies included in the original systematic review was adopted; two panel members independently assessed the QUADAS-2 applicability for any new study retrieved during the systematic review update, if performed. The indirectness was judged very serious or serious if all or more/equal than half of studies, respectively, have high or unclear concern regarding all or one to two QUADAS-2 domains for applicability concern. The indirectness was judged not serious in any other case. Furthermore, evidence was judged indirect if marked differences in population respect to the review question were suspected.

Inconsistency

Heterogeneity of results was assessed statistically by the I^2 test. We searched for a plausible source of heterogeneity by subgroup analyses according to the most plausible reason of heterogeneity, i.e. the type of samples, the type of test and the quality of the studies. These sub-analyses were performed only if subgroups included at least 4 or more studies. If a plausible explanation for heterogeneity was not identified, the quality of evidence was downgraded.. Inconsistency was judged very serious if high and unexplained heterogeneity ($I^2 > 90\%$) was detected. Inconsistency was judged not serious if either low ($I^2 < 50\%$) or otherwise explained through subgroup analyses. The inconsistency was judged serious in any other case.

Imprecision

We downgraded the evidence for imprecision if the boundaries of the 95% confidence intervals (CI) of all or $\geq 50\%$ of studies included a threshold of sensitivity or specificity for which the panel agreed that the estimate was adequate to support the decision. The panel agreed to set the threshold of sensitivity or specificity for all the diagnostic tests encompassed by the guidelines at 80% or 90%, respectively.

Publication bias

Linear regression of log odds ratios on inverse root of effective sample sizes test was used to assess the presence of publication bias when 3 or more studies were available [3, 4]. Publication bias was strongly suspected if the p value of the test was < 0.10 . Publication bias was considered as “strongly suspected” when less than 3 studies informed the outcome and could not be excluded.

Definitions of tests

Rapid nucleic acid amplification tests (NAAT) were considered as those having the capacity to be performed at the ‘point of care’ or in a ‘near patient’ setting. This means decentralized testing requiring minimal equipment, sample preparation, and biosafety considerations to be performed near a patient and outside of central laboratory testing [5]. Rapid antigen tests were considered all commercially available tests including both point of care testing, defined as above, and ‘in lab’ testing [6]. However, only one study provided information about in-lab testing [7].

Questions addressed by the guideline

In developing the guidelines for diagnosis of SARS-CoV-2, focus has been placed on testing by using commercial NAAT and rapid antigen detection. The performance characteristics of these tests depends on the population being examined, viral prevalence in the community, and biomaterial used. Furthermore, the resources required to apply any of these tests and approaches

depends on the local infrastructure of the individual setting and healthcare system facilities. This includes trained personnel, properly equipped diagnostic laboratories, and the types of tests being covered or reimbursed by the local healthcare system. Regarding particular recommendations for (or against) a specific diagnostic approach, the main beneficial outcomes considered were: i) reduction of mortality, facilitation of early treatment (if deemed to play a critical role), ii) reduction of viral transmission, iii) minimizing unnecessary treatment and/or isolation, iv) reduction of anxiety of patients/potentially exposed persons, and v) burden to healthcare systems with consequences for health inequality.

Recommendations

The 43 PICO questions selected were divided according to the target population into: A, patients with signs and symptoms of COVID-19 (Table 1); B, travelers from areas with low COVID-19 prevalence, healthcare workers, and asymptomatic individuals at risk for exposure (5 questions; Table 2); C, asymptomatic individuals and those with close contact with an infected person (Table 3). PICO questions in group A were further divided according to the type of test: A1, commercial rapid NAAT vs. standard NAAT (9 questions); A2, rapid antigen testing vs. standard NAAT (10 questions); A3, saliva sampling vs. nasopharyngeal swabs for NAAT (7 questions). PICO questions in group C were further divided into: C1, rapid antigen testing vs. NAAT (6 questions); C2, NAAT in saliva samples vs. nasopharyngeal swabs (6 questions). A full summary of the evidence for each PICO question is presented in Supplementary Appendices 2-4.

A: Patients with signs and symptoms of COVID-19

Rapid NAAT

The quality of the available evidence for all PICOs prioritized for patients with signs and symptoms compatible with COVID-19 was very low. Along these lines, the strength of recommendation was almost always weak except for cases where disease severity and/or risk of developing severe disease was considered of utmost importance. Diagnosis of COVID-19 with rapid NAAT will reduce the required time from sample acquisition to results. Extrapolation from studies looking into rapid NAATs for the detection of other respiratory viruses would suggest that early isolation is beneficial [8]. In general, rapid NAAT is suggested due to the high accuracy of the test, the low risk of anticipated harm, and its feasible implementation. However, the anticipated benefits are likely to depend on the different risk factors for hospitalization, mortality, and associated morbidity.

Rapid antigen testing

Diagnosis of COVID-19 with rapid antigen testing will reduce the required time from sample acquisition to results as well as reduce potential health inequities as antigen tests are (usually) readily available, easy to perform, and feasible to implement in various settings. This in turn, will reduce the potential consequences of a delayed COVID-19 diagnosis or even the ability to perform COVID-19 testing when access is limited due to restricted resources. However, current data suggest that antigen tests are not as accurate as the reference standard NAAT tests.

Saliva sampling

Laboratory-based NAAT in nasopharyngeal samples is the reference standard test for the diagnosis of COVID-19. The accuracy of other than nasopharyngeal samples, like saliva, is being investigated. Saliva sampling is easier to perform with essentially no adverse events. However, saliva processing may be challenging unless collected in appropriate media, since it has also been associated with aerosol generation [9, 10]. Especially in some subgroups of patients, like children, saliva might be preferred over nasopharyngeal swabs. Nevertheless,

children may not always be able to produce saliva on demand, and in such cases nasopharyngeal samples are still preferred. Notwithstanding, the available data suggest that the accuracy of saliva NAAT is not as high as nasopharyngeal NAAT.

B: Travelers from areas with low and high COVID-19 prevalence, healthcare workers, and asymptomatic individuals at risk for exposure

The quality of the available evidence for all PICOs prioritized for individuals at high risk of exposure to COVID-19 was very low. Therefore, the strength of recommendation for all PICOs in this section is weak. NAAT driven by questioning for contact history or high-risk exposure is very inaccurate, with very low sensitivity. Despite the very low quality of evidence, the panel recommended against the use of NAAT driven by questioning for contact history or exposure in returning travelers from areas of low and high prevalence of COVID-19, healthcare workers, and asymptomatic individuals at risk for exposure instead of universal NAAT testing.

C: Asymptomatic individuals and close contact of an infected person

As for all previous PICOs, quality of available evidence for asymptomatic individuals and close contacts was very low; thus all recommendations are weak.

Rapid antigen testing

In asymptomatic individuals and those with close contact with an infected person, the use of laboratory-based NAAT should be preferred over rapid antigen testing for diagnosis of COVID-19. The panel based its recommendations regarding rapid antigen testing on evidence from studies examining mainly the accuracy of first- and second-generation antigen tests that are not as accurate as the third-generation ones [11].

1 *NAAT in saliva samples*

2 In asymptomatic children <12 years old (with or without close contact), NAAT in saliva
3 samples was recommended over NAAT on nasopharyngeal swab samples for diagnosis of
4 COVID-19 considering the accuracy of the test, the large anticipated benefits and small harm,
5 low amount of resources, small incremental cost relative to net benefits, acceptability, and
6 feasibility of the test. In patients ≥ 12 years old, NAAT on nasopharyngeal samples should be
7 preferred, also considering that young children may not produce saliva on command.

8 **Future considerations**

10 It is worthwhile noting that several gaps remain regarding the evidence for diagnostic
11 testing in several areas. These include infection following vaccination or previous infection (re-
12 infection); performance of newer generation antigen tests in different patient populations and
13 samples; correlation of viral nucleic acid or antigen detection with contagiousness; accuracy of
14 different diagnostic tests in different populations and non-nasopharyngeal samples; the
15 accuracy of different diagnostic tests in asymptomatic individuals and the general population.
16 Moreover, there is a paucity of data regarding costs and/or resource modelling studies, patient
17 values, preferences and beliefs, definition of ‘feasibility’ of various tests, stakeholders’ opinion
18 regarding acceptability, and assessment criteria on the potential implications of interventions
19 on health inequities. The lack of objective criteria to judge the priority of clinically-relevant
20 questions should also be highlighted, and very little data is available regarding the clinical
21 impact (treatment, isolation, hospitalization) of these tests.

22 It is further noted that several methodological challenges were encountered that could be
23 improved upon. Firstly, literature databases specific for COVID-19 are not user friendly and
24 are difficult to navigate. Especially for diagnostic testing, there was a lack of exportable
25 literature. The ongoing pandemic also posed considerable time constraints for development of

guidelines given the large amount of new information, need for regular updates, and revision of recommendations, all within a context with limited resources for guideline development.

Among the limitations of the present guidelines is the very low quality of evidence and lack of dedicated resources to update specific evidence syntheses. Furthermore, poorly disclosed methodological details in relevant meta-analyses did not allow reproducibility of literature searches. We also included reagents for SARS-CoV-2 testing independently of the biomaterial for which they were optimized. We may thus not exclude the possibility that any unfavorable performance of a group of tests (e.g. pooled sensitivity of antigen tests) is due to diluting the favorable performance of tests, which are indeed optimized for one or another biomaterial. We also did not distinguish between different saliva samples, which may perform differently, and thus cannot draw more specific conclusions in this regard. In addition, it should be mentioned that this is not a systematic review, but rather guideline recommendations based on GRADE ADOLOPMENT methodology, which by definition is based on updating (if applicable) existing moderate to high quality already (or preprint) published evidence syntheses. Lastly, consensus was reached by a simple majority vote and not through consensus software/application such as the Delphi technique, as suggested by the ESCMID Guidelines Committee. However, its strengths include the multidisciplinary expertise of the writing group and the transparent, structured, thorough, and sound methodological approach adopted.

Contribution of individual authors:

Panel members: Chrysanthi Skevaki (clinical microbiology, virology, infection epidemiology and laboratory medicine), Paraskevi C. Fragkou (internal medicine, infectious diseases), Giulia De Angelis (clinical microbiology, infection epidemiology, infectious diseases), Giulia Menchinelli (clinical microbiology), Florence Morfin (clinical virology), Federico Garcia (clinical microbiology, infectious diseases), Fusun Can (clinical microbiology).

Conceptualization and co-ordination of the overall scope of the guidelines: CS, PCF. First manuscript draft (scope/context, methodology, discussion, etc.): CS, PCF. Initial screening for available evidence synthesis: PCF, DD, EM. Information extraction/summary of existing relevant literature on methodological aspects: PCF, GDA, GM. Literature search for update of existing evidence syntheses: PCF, GDA, GM, DD, EM, AG. Data extraction: PCF, GDA, GM, FC, FG, FMS, DD, AdS. Additional meta-analyses: GDA, GM. Creation of SOFs: GDA, GM. First draft of final recommendations (completed summary templates of panel consensus): PCF, DD. PICO formulation and final selection, recommendations, finalization of associated recommendations summary templates: CS, PCF, GDA, GM, FMS, FG, FC. Methodological consultation: TL, LS.

Updating: the panel will meet once a month to assess the need for further updates.

Acknowledgements: We acknowledge ESCMID for supporting medical writing services and ESGREV. Medical writing assistance by Patrick Moore and methodology direction and overview by Luigia Scudeller are acknowledged. We also acknowledge the technical assistance of Chiara Speziale (ESCMID).

Funding:

CS is supported by the Universities Giessen and Marburg Lung Center (UGMLC), the German Center for Lung Research (DZL), University Hospital Giessen and Marburg (UKGM) research funding according to article 2, section 3 cooperation agreement, and the Deutsche Forschungsgemeinschaft (DFG)-funded SFB 1021 (Project ID 197785619), KFO 309 (P10), and SK 317/1-1 (Project ID 428518790) as well as by the Foundation for Pathobiochemistry

and Molecular Diagnostics. FG is supported in part by Fundación Progreso y Salud, Consejería de Salud, Junta de Andalucía, Acciones de Intensificación del personal investigador. For EM is supported by the Foundation of the Faculty of Medicine, Philipps University Marburg, Stiftung P. E. Kempkes and Deutsche José Carreras Leukämie-Stiftung. PCF, DD, GDA, GM, FM, FC, ADS, AG, and TL declare no sources of funding.

Conflicts of interest:

For CS: Consultancy and research funding, Hycor Biomedical, BencardAllergie and Thermo Fisher Scientific; Research Funding, Mead Johnson Nutrition (MJN).

PCF, GDA, GM, FM, FC, ADS, AG, TL, DD, FG, and EM declare no conflicts of interest.

References

- [1] Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. 2020.
- [2] Bartoletti M, Azap O, Barac A, Bussini L, Ergonul O, Krause R, et al. ESCMID COVID-19 Living Guidelines: Drug Treatment and Clinical Management. *Clin Microbiol Infect*. 2022;In press.
- [3] Macaskill P, Walter SD, Irwig L. A comparison of methods to detect publication bias in meta-analysis. *Stat Med*. 2001;20(4):641-654.
- [4] Peters JL, Sutton AJ, Jones DR, Abrams KR, Rushton L. Comparison of two methods to detect publication bias in meta-analysis. *JAMA*. 2006;295(6):676-680.
- [5] Dinnes J, Deeks JJ, Berhane S, Taylor M, Adriano A, Davenport C, et al. Rapid, point-of-care antigen and molecular-based tests for diagnosis of SARS-CoV-2 infection. *Cochrane Database Syst Rev*. 2021;3CD013705.
- [6] Brummer LE, Katzenschlager S, Gaeddert M, Erdmann C, Schmitz S, Bota M, et al. Accuracy of novel antigen rapid diagnostics for SARS-CoV-2: A living systematic review and meta-analysis. *PLoS Med*. 2021;18(8):e1003735.
- [7] Yokota I, Sakurazawa T, Sugita J, Iwasaki S, Yasuda K, Yamashita N, et al. Performance of qualitative and quantitative antigen tests for SARS-CoV-2 in early symptomatic patients using saliva. *medRxiv*. 20202020.2011.2006.20227363.
- [8] Hinson JS, Rothman RE, Carroll K, Mostafa HH, Ghobadi K, Smith A, et al. Targeted rapid testing for SARS-CoV-2 in the emergency department is associated with large reductions in uninfected patient exposure time. *J Hosp Infect*. 2021;10735-39.
- [9] Berenger BM, Conly JM, Fonseca K, Hu J, Louie T, Schneider AR, et al. Saliva collected in universal transport media is an effective, simple and high-volume amenable method to detect SARS-CoV-2. *Clin Microbiol Infect*. 2021;27(4):656-657.

- [10] Lee RA, Herigon JC, Benedetti A, Pollock NR, Denkinger CM. Performance of Saliva, Oropharyngeal Swabs, and Nasal Swabs for SARS-CoV-2 Molecular Detection: a Systematic Review and Meta-analysis. *J Clin Microbiol.* 2021;59(5).
- [11] Cento V, Renica S, Matarazzo E, Antonello M, Colagrossi L, Di Ruscio F, et al. Frontline Screening for SARS-CoV-2 Infection at Emergency Department Admission by Third Generation Rapid Antigen Test: Can We Spare RT-qPCR? *Viruses.* 2021;13(5).
- [12] Assennato SM, Ritchie AV, Nadala C, Goel N, Tie C, Nadala LM, et al. Performance Evaluation of the SAMBA II SARS-CoV-2 Test for Point-of-Care Detection of SARS-CoV-2. *J Clin Microbiol.* 2020;59(1).
- [13] Chen JH, Yip CC, Poon RW, Chan KH, Cheng VC, Hung IF, et al. Evaluating the use of posterior oropharyngeal saliva in a point-of-care assay for the detection of SARS-CoV-2. *Emerg Microbes Infect.* 2020;9(1):1356-1359.
- [14] Collier DA, Assennato SM, Warne B, Sithole N, Sharrocks K, Ritchie A, et al. Point of Care Nucleic Acid Testing for SARS-CoV-2 in Hospitalized Patients: A Clinical Validation Trial and Implementation Study. *Cell Rep Med.* 2020;1(5):100062.
- [15] Cradic K, Lockhart M, Ozbolt P, Fatica L, Landon L, Lieber M, et al. Clinical Evaluation and Utilization of Multiple Molecular In Vitro Diagnostic Assays for the Detection of SARS-CoV-2. *Am J Clin Pathol.* 2020;154(2):201-207.
- [16] Gibani MM, Toumazou C, Sohbati M, Sahoo R, Karvela M, Hon TK, et al. Assessing a novel, lab-free, point-of-care test for SARS-CoV-2 (CovidNudge): a diagnostic accuracy study. *Lancet Microbe.* 2020;1(7):e300-e307.
- [17] Harrington A, Cox B, Snowden J, Bakst J, Ley E, Grajales P, et al. Comparison of Abbott ID Now and Abbott m2000 Methods for the Detection of SARS-CoV-2 from Nasopharyngeal and Nasal Swabs from Symptomatic Patients. *J Clin Microbiol.* 2020;58(8).
- [18] Moore NM, Li H, Schejbal D, Lindsley J, Hayden MK. Comparison of Two Commercial Molecular Tests and a Laboratory-Developed Modification of the CDC 2019-

- 1 nCoV Reverse Transcriptase PCR Assay for the Detection of SARS-CoV-2. *J Clin Microbiol.*
- 2 2020;58(8).
- 3 [19] SoRelle JA, Mahimainathan L, McCormick-Baw C, Cavuoti D, Lee F, Thomas A, et al.
- 4 Saliva for use with a point of care assay for the rapid diagnosis of COVID-19. *Clin Chim Acta.*
- 5 2020;510685-686.
- 6 [20] Thwe PM, Ren P. How many are we missing with ID NOW COVID-19 assay using
- 7 direct nasopharyngeal swabs? Findings from a mid-sized academic hospital clinical
- 8 microbiology laboratory. *Diagn Microbiol Infect Dis.* 2020;98(2):115123.
- 9 [21] Wong RC, Wong AH, Ho YI, Leung EC, Lai RW. Evaluation on testing of deep throat
- 10 saliva and lower respiratory tract specimens with Xpert Xpress SARS-CoV-2 assay. *J Clin*
- 11 *Virol.* 2020;131104593.
- 12 [22] Zhen W, Smith E, Manji R, Schron D, Berry GJ. Clinical Evaluation of Three Sample-
- 13 to-Answer Platforms for Detection of SARS-CoV-2. *J Clin Microbiol.* 2020;58(8).
- 14 [23] Dinnes J, Deeks JJ, Adriano A, Berhane S, Davenport C, Dittrich S, et al. Rapid, point-
- 15 of-care antigen and molecular-based tests for diagnosis of SARS-CoV-2 infection. *Cochrane*
- 16 *Database Syst Rev.* 2020;8CD013705.
- 17 [24] Filgueiras, Priscilla; Corsini, Camila; Almeida, Nathalie BF; Assis, Jessica; Pedrosa,
- 18 Maria Luysa; de Oliveira, Alana; Amorim, Raquel NH; de Miranda, Daniel AP; COVID-19
- 19 Rapid Antigen Test at hospital admission associated to the knowledge of individual risk factors
- 20 allow overcoming the difficulty of managing suspected patients in hospitals COVID-19 Rapid
- 21 Antigen Test facilitates the management of suspected patients on hospital admission medRxiv
- 22 2021.01.06.21249282.
- 23 [25] Abdulrahman A, Mustafa F, AlAwadhi AI, Alansari Q, AlAlawi B, AlQahtani M.
- 24 Comparison of SARS-COV-2 nasal antigen test to nasopharyngeal RT-PCR in mildly
- 25 symptomatic patients. *medRxiv.* 20202020.2011.2010.20228973.

- 1 [26] Bulilete O, Lorente P, Leiva A, Carandell E, Oliver A, Rojo E, et al. Evaluation of the
2 Panbio™ rapid antigen test for SARS-CoV-2 in primary health care centers and test sites.
3 medRxiv. 20202020.2011.2013.20231316.
- 4 [27] Gupta A, Khurana S, Das R, Srighyan D, Singh A, Mittal A, et al. Rapid chromatographic
5 immunoassay-based evaluation of COVID-19: A cross-sectional, diagnostic test accuracy study
6 & its implications for COVID-19 management in India. Indian J Med Res. 2020.
- 7 [28] Linares M, Perez-Tanoira R, Carrero A, Romanyk J, Perez-Garcia F, Gomez-Herruz P,
8 et al. Panbio antigen rapid test is reliable to diagnose SARS-CoV-2 infection in the first 7 days
9 after the onset of symptoms. J Clin Virol. 2020;133104659.
- 10 [29] Lv Y, Ma Y, Si Y, Zhu X, Zhang L, Feng H, et al. Rapid SARS-CoV-2 antigen detection
11 potentiates early diagnosis of COVID-19 disease. Biosci Trends. 2021;15(2):93-99.
- 12 [30] Nagura-Ikeda M, Imai K, Tabata S, Miyoshi K, Murahara N, Mizuno T, et al. Clinical
13 Evaluation of Self-Collected Saliva by Quantitative Reverse Transcription-PCR (RT-qPCR),
14 Direct RT-qPCR, Reverse Transcription-Loop-Mediated Isothermal Amplification, and a
15 Rapid Antigen Test To Diagnose COVID-19. J Clin Microbiol. 2020;58(9).
- 16 [31] Pray IW, Ford L, Cole D, Lee C, Bigouette JP, Abedi GR, et al. Performance of an
17 Antigen-Based Test for Asymptomatic and Symptomatic SARS-CoV-2 Testing at Two
18 University Campuses - Wisconsin, September-October 2020. MMWR Morb Mortal Wkly Rep.
19 2021;69(5152):1642-1647.
- 20 [32] Ristic M, Nikolic N, Cabarkapa V, Turkulov V, Petrovic V. Validation of the
21 STANDARD Q COVID-19 antigen test in Vojvodina, Serbia. PLoS One.
22 2021;16(2):e0247606.
- 23 [33] Takeuchi Y, Akashi Y, Kato D, Kuwahara M, Muramatsu S, Ueda A, et al. The
24 evaluation of a newly developed antigen test (QuickNavi-COVID19 Ag) for SARS-CoV-2: A
25 prospective observational study in Japan. J Infect Chemother. 2021;27(6):890-894.

- 1 [34] Yamamoto K, Suzuki M, Yamada G, Sudo T, Nomoto H, Kinoshita N, et al. Utility of
2 the antigen test for coronavirus disease 2019: Factors influencing the prediction of the
3 possibility of disease transmission. *Int J Infect Dis.* 2021;10465-72.
- 4 [35] Foundation for Innovative New Diagnostics (FIND). Evaluation of Bionote, Inc.
5 NowCheck COVID-19 Ag Test - External Report. Switzerland: FIND, 2020.
- 6 [36] Foundation for Innovative New Diagnostics (FIND). Evaluation of SD Biosensor, Inc.
7 STANDARD Q COVID-19 Ag Test - External Report. Switzerland: FIND, 2020.
- 8 [37] Foundation for Innovative New Diagnostics (FIND). Evaluation of Abbott Panbio
9 COVID-19 Ag Rapid Test Device - External Report. Switzerland: FIND, 2020.
- 10 [38] Foundation for Innovative New Diagnostics (FIND). Evaluation of RapiGEN Inc.
11 BIOCREDIT COVID-19 Ag - External Report. Switzerland: FIND, 2020.
- 12 [39] Fourati S, Audureau E, Chevaliez S, Pawlotsky JM. Évaluation de la performance
13 diagnostique des tests rapides d'orientation diagnostique antigéniques Covid-19. Laboratoire
14 de virologie, département de prévention, diagnostic et traitement des Infections ; service de
15 santé publique, hôpitaux universitaires Henri-Mondor, AP-HP ; institut Mondor de recherche
16 biomédicale (Inserm U955); université Paris Est-Créteil, 29 septembre 2020.
- 17 [40] Agullo V, Fernandez-Gonzalez M, Ortiz de la Tabla V, Gonzalo-Jimenez N, Garcia JA,
18 Masia M, et al. Evaluation of the rapid antigen test Panbio COVID-19 in saliva and nasal swabs
19 in a population-based point-of-care study. *J Infect.* 2021;82(5):186-230.
- 20 [41] Albert E, Torres I, Bueno F, Huntley D, Molla E, Fernandez-Fuentes MA, et al. Field
21 evaluation of a rapid antigen test (Panbio COVID-19 Ag Rapid Test Device) for COVID-19
22 diagnosis in primary healthcare centres. *Clin Microbiol Infect.* 2021;27(3):472 e477-472 e410.
- 23 [42] Alemany A, Baro B, Ouchi D, Rodo P, Ubals M, Corbacho-Monne M, et al. Analytical
24 and clinical performance of the panbio COVID-19 antigen-detecting rapid diagnostic test. *J*
25 *Infect.* 2021;82(5):186-230.

- 1 [43] Aoki K, Nagasawa T, Ishii Y, Yagi S, Kashiwagi K, Miyazaki T, et al. Evaluation of
2 clinical utility of novel coronavirus antigen detection reagent, Espline(R) SARS-CoV-2. J
3 Infect Chemother. 2021;27(2):319-322.
- 4 [44] Beck ET, Paar W, Fojut L, Serwe J, Jahnke RR. Comparison of the Quidel Sofia SARS
5 FIA Test to the Hologic Aptima SARS-CoV-2 TMA Test for Diagnosis of COVID-19 in
6 Symptomatic Outpatients. J Clin Microbiol. 2021;59(2).
- 7 [45] Boum Y, Fai KN, Nikolay B, Mboringong AB, Bebell LM, Ndifon M, et al.
8 Performance and operational feasibility of antigen and antibody rapid diagnostic tests for
9 COVID-19 in symptomatic and asymptomatic patients in Cameroon: a clinical, prospective,
10 diagnostic accuracy study. Lancet Infect Dis. 2021;21(8):1089-1096.
- 11 [46] Caruana G, Croxatto A, Kampouri E, Kritikos A, Opota O, Foerster M, et al.
12 Implementing SARS-CoV-2 Rapid Antigen Testing in the Emergency Ward of a Swiss
13 University Hospital: The INCREASE Study. Microorganisms. 2021;9(4).
- 14 [47] Cerutti F, Burdino E, Milia MG, Alice T, Gregori G, Bruzzone B, et al. Urgent need of
15 rapid tests for SARS CoV-2 antigen detection: Evaluation of the SD-Biosensor antigen test for
16 SARS-CoV-2. J Clin Virol. 2020;132:104654.
- 17 [48] Courtellemont L, Guinard J, Guillaume C, Giache S, Rzepecki V, Seve A, et al. High
18 performance of a novel antigen detection test on nasopharyngeal specimens for diagnosing
19 SARS-CoV-2 infection. J Med Virol. 2021;93(5):3152-3157.
- 20 [49] Drain PK, Ampajwala M, Chappel C, Gvozden AB, Hoppers M, Wang M, et al. A
21 Rapid, High-Sensitivity SARS-CoV-2 Nucleocapsid Immunoassay to Aid Diagnosis of Acute
22 COVID-19 at the Point of Care: A Clinical Performance Study. Infect Dis Ther.
23 2021;10(2):753-761.
- 24 [50] Drevinek P, Hurych J, Kepka Z, Briksi A, Kulich M, Zajac M, et al. The sensitivity of
25 SARS-CoV-2 antigen tests in the view of large-scale testing. medRxiv.
26 20202020.2011.2023.20237198.

- 1 [51] Fenollar F, Bouam A, Ballouche M, Fuster L, Prudent E, Colson P, et al. Evaluation of
2 the Panbio COVID-19 Rapid Antigen Detection Test Device for the Screening of Patients with
3 COVID-19. *J Clin Microbiol.* 2021;59(2).
- 4 [52] Gonzalez-Donapetry P, Garcia-Clemente P, Bloise I, Garcia-Sanchez C, Sanchez
5 Castellano MA, Romero MP, et al. Think of the Children: Evaluation of SARS-CoV-2 Rapid
6 Antigen Test in Pediatric Population. *Pediatr Infect Dis J.* 2021;40(5):385-388.
- 7 [53] Herrera V, Hsu V, Adewale A, Johnson L, Hendrix T, Kuhlman J, et al. Testing
8 Healthcare Workers Exposed to COVID19 using Rapid Antigen Detection. *medRxiv.*
9 20202020.2008.2012.20172726.
- 10 [54] Jakobsen KK, Jensen JS, Todsen T, Lippert F, Martel CJ-M, Klokke M, et al. Detection
11 of SARS-CoV-2 infection by rapid antigen test in comparison with RT-PCR in a public setting.
12 *medRxiv.* 20212021.2001.2022.21250042.
- 13 [55] Korenkov M, Poopalasingam N, Madler M, Vanshylla K, Eggeling R, Wirtz M, et al.
14 Assessment of SARS-CoV-2 infectivity by a Rapid Antigen Detection Test. *medRxiv.*
15 20212021.2003.2030.21254624.
- 16 [56] Kruger LJ, Gaeddert M, Tobian F, Lainati F, Gottschalk C, Klein JAF, et al. The Abbott
17 PanBio WHO emergency use listed, rapid, antigen-detecting point-of-care diagnostic test for
18 SARS-CoV-2-Evaluation of the accuracy and ease-of-use. *PLoS One.* 2021;16(5):e0247918.
- 19 [57] Landaas ET, Storm ML, Tollanes MC, Barlinn R, Kran AB, Bragstad K, et al.
20 Diagnostic performance of a SARS-CoV-2 rapid antigen test in a large, Norwegian cohort. *J*
21 *Clin Virol.* 2021;137:104789.
- 22 [58] Lindner AK, Nikolai O, Kausch F, Wintel M, Hommes F, Gertler M, et al. Head-to-
23 head comparison of SARS-CoV-2 antigen-detecting rapid test with self-collected nasal swab
24 versus professional-collected nasopharyngeal swab. *Eur Respir J.* 2021;57(4).

- [59] Lindner AK, Nikolai O, Rohardt C, Burock S, Hulso C, Bolke A, et al. Head-to-head comparison of SARS-CoV-2 antigen-detecting rapid test with professional-collected nasal versus nasopharyngeal swab. *Eur Respir J*. 2021;57(5).
- [60] Masia M, Fernandez-Gonzalez M, Sanchez M, Carvajal M, Garcia JA, Gonzalo-Jimenez N, et al. Nasopharyngeal Panbio COVID-19 Antigen Performed at Point-of-Care Has a High Sensitivity in Symptomatic and Asymptomatic Patients With Higher Risk for Transmission and Older Age. *Open Forum Infect Dis*. 2021;8(3):ofab059.
- [61] Mboumba Bouassa RS, Veyer D, Pere H, Belec L. Analytical performances of the point-of-care SIENNA COVID-19 Antigen Rapid Test for the detection of SARS-CoV-2 nucleocapsid protein in nasopharyngeal swabs: A prospective evaluation during the COVID-19 second wave in France. *Int J Infect Dis*. 2021;1068-12.
- [62] Merino P, Guinea J, Munoz-Gallego I, Gonzalez-Donapetry P, Galan JC, Antona N, et al. Multicenter evaluation of the Panbio COVID-19 rapid antigen-detection test for the diagnosis of SARS-CoV-2 infection. *Clin Microbiol Infect*. 2021.
- [63] Mertens P, De Vos N, Martiny D, Jassoy C, Mirazimi A, Cuypers L, et al. Development and Potential Usefulness of the COVID-19 Ag Respi-Strip Diagnostic Assay in a Pandemic Context. *Front Med (Lausanne)*. 2020;7225.
- [64] Mockel M, Corman VM, Stegemann MS, Hofmann J, Stein A, Jones TC, et al. SARS-CoV-2 antigen rapid immunoassay for diagnosis of COVID-19 in the emergency department. *Biomarkers*. 2021;26(3):213-220.
- [65] Nikolai O, Rohardt C, Tobian F, Junge A, Corman VM, Jones TC, et al. Anterior nasal versus nasal mid-turbinate sampling for a SARS-CoV-2 antigen-detecting rapid test: does localisation or professional collection matter? *Infect Dis (Lond)*. 2021;1-6.
- [66] Olearo F, Norz D, Heinrich F, Sutter JP, Roedl K, Schultze A, et al. Handling and accuracy of four rapid antigen tests for the diagnosis of SARS-CoV-2 compared to RT-qPCR. *J Clin Virol*. 2021;137104782.

- [67] Parada-Ricart E, Gomez-Bertomeu F, Pico-Plana E, Olona-Cabases M. Usefulness of the antigen for diagnosing SARS-CoV-2 infection in patients with and without symptoms. *Enferm Infecc Microbiol Clin*. 2021;39(7):357-358.
- [68] Pekosz A, Parvu V, Li M, Andrews JC, Manabe YC, Kodsí S, et al. Antigen-Based Testing but Not Real-Time Polymerase Chain Reaction Correlates With Severe Acute Respiratory Syndrome Coronavirus 2 Viral Culture. *Clin Infect Dis*. 2021.
- [69] Perez-Garcia F, Romanyk J, Gomez-Herruz P, Arroyo T, Perez-Tanoira R, Linares M, et al. Diagnostic performance of CerTest and Panbio antigen rapid diagnostic tests to diagnose SARS-CoV-2 infection. *J Clin Virol*. 2021;137104781.
- [70] Peto T, Team UC-LFO. COVID-19: Rapid antigen detection for SARS-CoV-2 by lateral flow assay: A national systematic evaluation of sensitivity and specificity for mass-testing. *EClinicalMedicine*. 2021;36100924.
- [71] Pollock NR, Jacobs JR, Tran K, Cranston AE, Smith S, O'Kane CY, et al. Performance and Implementation Evaluation of the Abbott BinaxNOW Rapid Antigen Test in a High-Throughput Drive-Through Community Testing Site in Massachusetts. *J Clin Microbiol*. 2021;59(5).
- [72] Porte L, Legarraga P, Iruretagoyena M, Vollrath V, Pizarro G, Munita JM, et al. Rapid SARS-CoV-2 antigen detection by immunofluorescence – a new tool to detect infectivity. *medRxiv*. 20202020.2010.2004.20206466.
- [73] Porte L, Legarraga P, Vollrath V, Aguilera X, Munita JM, Araos R, et al. Evaluation of a novel antigen-based rapid detection test for the diagnosis of SARS-CoV-2 in respiratory samples. *Int J Infect Dis*. 2020;99328-333.
- [74] Prince-Guerra JL, Almendares O, Nolen LD, Gunn JKL, Dale AP, Buono SA, et al. Evaluation of Abbott BinaxNOW Rapid Antigen Test for SARS-CoV-2 Infection at Two Community-Based Testing Sites - Pima County, Arizona, November 3-17, 2020. *MMWR Morb Mortal Wkly Rep*. 2021;70(3):100-105.

- [75] Regev-Yochay G, Kriger O, Beni S, Rubin C, Mina MJ, Mechnik B, et al. Real World Performance of SARS-CoV-2 Antigen Rapid Diagnostic Tests in Various Clinical Settings. medRxiv. 20212021.2003.2002.21252400.
- [76] Schwob JM, Miauton A, Petrovic D, Perdrix J, Senn N, Jatton K, et al. Antigen rapid tests, nasopharyngeal PCR and saliva PCR to detect SARS-CoV-2: a prospective comparative clinical trial. medRxiv. 20202020.2011.2023.20237057.
- [77] Scohy A, Anantharajah A, Bodeus M, Kabamba-Mukadi B, Verroken A, Rodriguez-Villalobos H. Low performance of rapid antigen detection test as frontline testing for COVID-19 diagnosis. J Clin Virol. 2020;129104455.
- [78] Shah MM, Salvatore PP, Ford L, Kamitani E, Whaley MJ, Mitchell K, et al. Performance of Repeat BinaxNOW Severe Acute Respiratory Syndrome Coronavirus 2 Antigen Testing in a Community Setting, Wisconsin, November 2020-December 2020. Clin Infect Dis. 2021;73(Suppl 1):S54-S57.
- [79] Sood N, Shetgiri R, Rodriguez A, Jimenez D, Treminino S, Daflos A, et al. Evaluation of the Abbott BinaxNOW rapid antigen test for SARS-CoV-2 infection in children: Implications for screening in a school setting. PLoS One. 2021;16(4):e0249710.
- [80] Stokes W, Berenger BM, Portnoy D, Scott B, Szelewicki J, Singh T, et al. Clinical performance of the Abbott Panbio with nasopharyngeal, throat, and saliva swabs among symptomatic individuals with COVID-19. Eur J Clin Microbiol Infect Dis. 2021;40(8):1721-1726.
- [81] Thommes L, Burkert FR, Ottl KW, Goldin D, Loacker L, Lanser L, et al. Comparative evaluation of four SARS-CoV-2 antigen tests in hospitalized patients. Int J Infect Dis. 2021;105144-146.
- [82] Torres I, Poujois S, Albert E, Alvarez G, Colomina J, Navarro D. Point-of-care evaluation of a rapid antigen test (CLINITEST() Rapid COVID-19 Antigen Test) for diagnosis

of SARS-CoV-2 infection in symptomatic and asymptomatic individuals. *J Infect.* 2021;82(5):e11-e12.

[83] Turcato G, Zaboli A, Pfeifer N, Ciccariello L, Sibilio S, Tezza G, et al. Clinical application of a rapid antigen test for the detection of SARS-CoV-2 infection in symptomatic and asymptomatic patients evaluated in the emergency department: A preliminary report. *J Infect.* 2021;82(3):e14-e16.

[84] Van der Moeren N, Zwart VF, Lodder EB, Van den Bijllaardt W, Van Esch HRJM, Stohr JJM, et al. PERFORMANCE EVALUATION OF A SARS-COV-2 RAPID ANTIGENTEST: TEST PERFORMANCE IN THE COMMUNITY IN THE NETHERLANDS. medRxiv. 20202020.2010.2019.20215202.

[85] Veyrenche N, Bollore K, Pisoni A, Bedin AS, Mondain AM, Ducos J, et al. Diagnosis value of SARS-CoV-2 antigen/antibody combined testing using rapid diagnostic tests at hospital admission. *J Med Virol.* 2021;93(5):3069-3076.

[86] Villaverde S, Dominguez-Rodriguez S, Sabrido G, Perez-Jorge C, Plata M, Romero MP, et al. Diagnostic Accuracy of the Panbio Severe Acute Respiratory Syndrome Coronavirus 2 Antigen Rapid Test Compared with Reverse-Transcriptase Polymerase Chain Reaction Testing of Nasopharyngeal Samples in the Pediatric Population. *J Pediatr.* 2021;232:287-289 e284.

[87] Weitzel T, Legarraga P, Iruretagoyena M, Pizarro G, Vollrath V, Araos R, et al. Comparative evaluation of four rapid SARS-CoV-2 antigen detection tests using universal transport medium. *Travel Med Infect Dis.* 2021;39:101942.

[88] Young S, Taylor SN, Cammarata CL, Varnado KG, Roger-Dalbert C, Montano A, et al. Clinical Evaluation of BD Veritor SARS-CoV-2 Point-of-Care Test Performance Compared to PCR-Based Testing and versus the Sofia 2 SARS Antigen Point-of-Care Test. *J Clin Microbiol.* 2020;59(1).

- 1 [89] Torres I, Poujois S, Albert E, Colomina J, Navarro D. Evaluation of a rapid antigen test
2 (Panbio COVID-19 Ag rapid test device) for SARS-CoV-2 detection in asymptomatic close
3 contacts of COVID-19 patients. *Clin Microbiol Infect*. 2021;27(4):636 e631-636 e634.
- 4 [90] Bhattacharya D, Parai D, Rout UK, Dash P, Nanda RR, Dash GC, et al. Saliva for
5 diagnosis of SARS-CoV-2: First report from India. *J Med Virol*. 2021;93(4):2529-2533.
- 6 [91] Braz-Silva PH, Mamana AC, Romano CM, Felix AC, de Paula AV, Fereira NE, et al.
7 Performance of at-home self-collected saliva and nasal-oro-pharyngeal swabs in the surveillance
8 of COVID-19. *J Oral Microbiol*. 2020;13(1):1858002.
- 9 [92] Guclu E, Koroglu M, Yurumez Y, Toptan H, Kose E, Guneyasu F, et al. Comparison of
10 saliva and oro-nasopharyngeal swab sample in the molecular diagnosis of COVID-19. *Rev*
11 *Assoc Med Bras* (1992). 2020;66(8):1116-1121.
- 12 [93] Hanson KE, Barker AP, Hillyard DR, Gilmore N, Barrett JW, Orlandi RR, et al. Self-
13 Collected Anterior Nasal and Saliva Specimens versus Health Care Worker-Collected
14 Nasopharyngeal Swabs for the Molecular Detection of SARS-CoV-2. *J Clin Microbiol*.
15 2020;58(11).
- 16 [94] Landry ML, Criscuolo J, Peaper DR. Challenges in use of saliva for detection of SARS
17 CoV-2 RNA in symptomatic outpatients. *J Clin Virol*. 2020;130104567.
- 18 [95] Sakanashi D, Asai N, Nakamura A, Miyazaki N, Kawamoto Y, Ohno T, et al.
19 Comparative evaluation of nasopharyngeal swab and saliva specimens for the molecular
20 detection of SARS-CoV-2 RNA in Japanese patients with COVID-19. *J Infect Chemother*.
21 2021;27(1):126-129.
- 22 [96] Senok A, Alsuwaidi H, Atrah Y, Al Ayedi O, Al Zahid J, Han A, et al. Saliva as an
23 Alternative Specimen for Molecular COVID-19 Testing in Community Settings and
24 Population-Based Screening. *Infect Drug Resist*. 2020;133393-3399.

- 1 [97] Teo AKJ, Choudhury Y, Tan IB, Cher CY, Chew SH, Wan ZY, et al. Saliva is more
2 sensitive than nasopharyngeal or nasal swabs for diagnosis of asymptomatic and mild COVID-
3 19 infection. *Sci Rep*. 2021;11(1):3134.
- 4 [98] Aita A, Basso D, Cattelan AM, Fioretto P, Navaglia F, Barbaro F, et al. SARS-CoV-2
5 identification and IgA antibodies in saliva: One sample two tests approach for diagnosis. *Clin*
6 *Chim Acta*. 2020;510717-722.
- 7 [99] Becker D, Sandoval E, Amin A, De Hoff P, Diets A, Leonetti N, et al. Saliva is less
8 sensitive than nasopharyngeal swabs for COVID-19 detection in the community setting.
9 *medRxiv*. 20202020.2005.2011.20092338.
- 10 [100] Byrne RL, Kay GA, Kontogianni K, Aljayyousi G, Brown L, Collins AM, et al. Saliva
11 Alternative to Upper Respiratory Swabs for SARS-CoV-2 Diagnosis. *Emerg Infect Dis*.
12 2020;26(11):2770-2771.
- 13 [101] Dogan OA, Kose B, Agaoglu NB, Yildiz J, Alkurt G, Demirkol YK, et al. Does
14 sampling saliva increase detection of SARS-CoV-2 by RT-PCR? Comparing saliva with oro-
15 nasopharyngeal swabs. *J Virol Methods*. 2021;290114049.
- 16 [102] Echavarria M, Reyes NS, Rodriguez PE, Ypas M, Ricarte C, Rodriguez MP, et al. Self-
17 collected saliva for SARS-CoV-2 detection: A prospective study in the emergency room. *J Med*
18 *Virol*. 2021;93(5):3268-3272.
- 19 [103] Goldfarb DM, Tilley P, Al-Rawahi GN, Srigley JA, Ford G, Pedersen H, et al. Self-
20 Collected Saline Gargle Samples as an Alternative to Health Care Worker-Collected
21 Nasopharyngeal Swabs for COVID-19 Diagnosis in Outpatients. *J Clin Microbiol*. 2021;59(4).
- 22 [104] Jamal AJ, Mozafarihashjin M, Coomes E, Powis J, Li AX, Paterson A, et al. Sensitivity
23 of Nasopharyngeal Swabs and Saliva for the Detection of Severe Acute Respiratory Syndrome
24 Coronavirus 2. *Clin Infect Dis*. 2021;72(6):1064-1066.

- 1 [105] Ku CW, Shivani D, Kwan JQT, Loy SL, Erwin C, Ko KKK, et al. Validation of self-
2 collected buccal swab and saliva as a diagnostic tool for COVID-19. *Int J Infect Dis.*
3 2021;104:255-261.
- 4 [106] Leung EC, Chow VC, Lee MK, Lai RW. Deep throat saliva as an alternative diagnostic
5 specimen type for the detection of SARS-CoV-2. *J Med Virol.* 2021;93(1):533-536.
- 6 [107] McCormick-Baw C, Morgan K, Gaffney D, Cazares Y, Jaworski K, Byrd A, et al. Saliva
7 as an Alternate Specimen Source for Detection of SARS-CoV-2 in Symptomatic Patients Using
8 Cepheid Xpert Xpress SARS-CoV-2. *J Clin Microbiol.* 2020;58(8).
- 9 [108] Miguères M, Mengelle C, Dimeglio C, Didier A, Alvarez M, Delobel P, et al. Saliva
10 sampling for diagnosing SARS-CoV-2 infections in symptomatic patients and asymptomatic
11 carriers. *J Clin Virol.* 2020;130:104580.
- 12 [109] Pasomsub E, Watcharananan SP, Boonyawat K, Janchompoo P, Wongtabtim G,
13 Sukswan W, et al. Saliva sample as a non-invasive specimen for the diagnosis of coronavirus
14 disease 2019: a cross-sectional study. *Clin Microbiol Infect.* 2021;27(2):285 e281-285 e284.
- 15 [110] Procop GW, Shrestha NK, Vogel S, Van Sickle K, Harrington S, Rhoads DD, et al. A
16 Direct Comparison of Enhanced Saliva to Nasopharyngeal Swab for the Detection of SARS-
17 CoV-2 in Symptomatic Patients. *J Clin Microbiol.* 2020;58(11).
- 18 [111] Tsujimoto Y, Terada J, Kimura M, Moriya A, Motohashi A, Izumi S, et al. Diagnostic
19 accuracy of nasopharyngeal swab, nasal swab and saliva swab samples for the detection of
20 SARS-CoV-2 using RT-PCR. *Infect Dis (Lond).* 2021;53(8):581-589.
- 21 [112] Uwamino Y, Nagata M, Aoki W, Fujimori Y, Nakagawa T, Yokota H, et al. Accuracy
22 and stability of saliva as a sample for reverse transcription PCR detection of SARS-CoV-2. *J*
23 *Clin Pathol.* 2021;74(1):67-68.
- 24 [113] Vaz SN, Santana DS, Netto EM, Wang WK, Brites C. Validation of the GeneXpert
25 Xpress SARS-CoV-2 PCR assay using saliva as biological specimen. *Braz J Infect Dis.*
26 2021;25(2):101543.

- 1 [114] Villar LM, da Costa VD, Marques BCL, da Silva LL, Santos AC, Mendonca A, et al.
2 USEFULNESS OF SALIVA SAMPLES FOR DETECTING SARS-CoV-2 RNA AMONG
3 LIVER DISEASE PATIENTS. *J Infect.* 2021;82(1):e4-e5.
- 4 [115] Yee R, Truong TT, Pannaraj PS, Eubanks N, Gai E, Jumarang J, et al. Saliva Is a
5 Promising Alternative Specimen for the Detection of SARS-CoV-2 in Children and Adults. *J*
6 *Clin Microbiol.* 2021;59(2).
- 7 [116] Yokota I, Hattori T, Shane PY, Konno S, Nagasaka A, Takeyabu K, et al. Equivalent
8 SARS-CoV-2 viral loads by PCR between nasopharyngeal swab and saliva in symptomatic
9 patients. *Sci Rep.* 2021;11(1):4500.
- 10 [117] Altawalah H, AlHuraish F, Alkandari WA, Ezzikouri S. Saliva specimens for detection
11 of severe acute respiratory syndrome coronavirus 2 in Kuwait: A cross-sectional study. *J Clin*
12 *Virol.* 2020;132104652.
- 13 [118] Moreno-Contreras J, Espinoza MA, Sandoval-Jaime C, Cantu-Cuevas MA, Baron-
14 Olivares H, Ortiz-Orozco OD, et al. Saliva Sampling and Its Direct Lysis, an Excellent Option
15 To Increase the Number of SARS-CoV-2 Diagnostic Tests in Settings with Supply Shortages.
16 *J Clin Microbiol.* 2020;58(10).
- 17 [119] Barrett ES, Horton DB, Roy J, Gennaro ML, Brooks A, Tischfield J, et al. Prevalence
18 of SARS-CoV-2 infection in previously undiagnosed health care workers at the onset of the
19 U.S. COVID-19 epidemic. *medRxiv.* 2020.
- 20 [120] Gudbjartsson DF, Helgason A, Jonsson H, Magnusson OT, Melsted P, Norddahl GL, et
21 al. Spread of SARS-CoV-2 in the Icelandic Population. *N Engl J Med.* 2020;382(24):2302-
22 2315.
- 23 [121] Viswanathan M, Kahwati L, Jahn B, Giger K, Dobrescu AI, Hill C, et al. Universal
24 screening for SARS-CoV-2 infection: a rapid review. *Cochrane Database Syst Rev.*
25 2020;9CD013718.

[122] Hoehl S, Rabenau H, Berger A, Kortenbusch M, Cinatl J, Bojkova D, et al. Evidence of SARS-CoV-2 Infection in Returning Travelers from Wuhan, China. *N Engl J Med*. 2020;382(13):1278-1280.

[123] Hasanoglu I, Korukluoglu G, Asilturk D, Cosgun Y, Kalem AK, Altas AB, et al. Higher viral loads in asymptomatic COVID-19 patients might be the invisible part of the iceberg. *Infection*. 2021;49(1):117-126.

[124] Yokota I, Shane PY, Okada K, Unoki Y, Yang Y, Inao T, et al. Mass Screening of Asymptomatic Persons for Severe Acute Respiratory Syndrome Coronavirus 2 Using Saliva. *Clin Infect Dis*. 2021;73(3):e559-e565.

[125] Rao M, Rashid FA, Sabri F, Jamil NN, Zain R, Hashim R, et al. Comparing Nasopharyngeal Swab and Early Morning Saliva for the Identification of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). *Clin Infect Dis*. 2021;72(9):e352-e356.

[126] Moreira VM, Mascarenhas P, Machado V, Botelho J, Mendes JJ, Taveira N, et al. Diagnosis of SARS-Cov-2 Infection by RT-PCR Using Specimens Other Than Naso- and Oropharyngeal Swabs: A Systematic Review and Meta-Analysis. *Diagnostics (Basel)*. 2021;11(2).

1 Table 1. PICO questions and recommendations in patients with signs and symptoms of COVID-
2 19.

A	PICO question	Recommendation	Strength of recommendation*	Overall certainty of the evidence**	References
Rapid NAAT					
1	In patients with signs and symptoms compatible with mild or moderate COVID-19, should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19?	In patients with signs and symptoms compatible with mild or moderate COVID-19 we suggest the use of rapid NAAT versus laboratory based NAAT testing for the diagnosis of COVID-19.	Weak for	Very low	[12-23].
2	In patients with signs and symptoms compatible with severe or critical COVID-19, should commercial rapid NAAT testing be used, compared with standard NAAT (commercial and/or in house) testing for diagnosis of COVID-19?	In patients with signs and symptoms compatible with severe or critical COVID-19 we suggest the use of rapid NAAT versus laboratory based NAAT for the diagnosis of COVID-19.	Strong for	Very low	[12-23]
3	In patients with signs and symptoms compatible with COVID-19, should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19 in nasopharyngeal samples?	In patients with signs and symptoms compatible with COVID-19 we suggest the use of rapid NAAT in nasopharyngeal samples versus laboratory based NAAT in nasopharyngeal samples for the diagnosis of COVID-19.	Weak for	Very low	[13, 15, 16, 18, 20, 22, 23]
4	In patients with signs and symptoms compatible with COVID-19, should commercial rapid NAAT be used, compared with the standard NAAT (commercial and/or in	In patients with signs and symptoms compatible with COVID-19 we suggest the use of rapid NAAT in samples other than nasopharyngeal swab versus laboratory based NAAT in samples	Weak for	Very low	[12, 14, 15, 17, 21, 23]

	house) for diagnosis of COVID-19 in samples other than nasopharyngeal swab?	other than nasopharyngeal swab for the diagnosis of COVID-19, when allowed by regulatory boards and manufacturer's instructions.			
5	In patients with signs and symptoms compatible with COVID-19 of equal or less than 7 days-onset , should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19?	In patients with signs and symptoms compatible with COVID-19 of equal or less than 7 days-onset we suggest the use of rapid NAAT versus laboratory based NAAT for the diagnosis of COVID-19.	Weak for	Very low	[12-22]
6	In patients with signs and symptoms compatible with COVID-19 of more than 7 days-onset , should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19?	In patients with signs and symptoms compatible with COVID-19 of more than 7 days-onset we suggest the use of rapid NAAT versus laboratory based NAAT for the diagnosis of COVID-19.	Weak for	Very low	[12-23]
7	In children <12 years old with signs and symptoms compatible with COVID-19, should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19?	In children <12 years old with signs and symptoms compatible with COVID-19 we suggest the use of rapid NAAT versus laboratory based NAAT for diagnosis of COVID-19.	Weak for	Very low	[12, 15, 17-20, 22, 23]
8	In patients ≥12 years old with signs and symptoms compatible with COVID-19, should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19?	In patients ≥12 years old with signs and symptoms compatible with COVID-19, we suggest the use of rapid NAAT versus laboratory based NAAT for diagnosis of COVID-19.	Weak for	Very low	[13, 14, 16, 21, 23]
9	In patients with signs and symptoms compatible with	In patients with signs and symptoms compatible with	Weak for	Very low	[12-23]

COVID-19 **at risk for severe illness**, should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19?

Rapid antigen testing

COVID-19 at risk for severe disease, we suggest the use of rapid NAAT versus laboratory-based NAAT for diagnosis of COVID-19.

- | | | | | | |
|----|---|---|----------------|----------|--|
| 10 | In patients with signs and symptoms compatible with mild or moderate COVID-19, should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19? | In patients with mild and moderate COVID-19, we suggest the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19. | Weak against | Very low | [6, 23-34] |
| 11 | In patients with signs and symptoms compatible with severe or critical COVID-19, should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in house) testing for diagnosis of COVID-19? | In patients with severe or critical COVID-19, we recommend the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19. | Strong against | Very low | [6, 7, 23, 30, 35-88] |
| 12 | In patients with signs and symptoms compatible with COVID-19, should rapid antigen detection testing be used, compared with the standard NAAT (commercial and/or in house) for diagnosis of COVID-19 in nasopharyngeal samples ? | In patients with signs and symptoms compatible with COVID-19, we suggest the use of laboratory based NAAT in nasopharyngeal samples versus rapid antigen detection testing in nasopharyngeal samples for diagnosis of COVID-19. | Weak against | Very low | [6, 7, 23, 24, 26-29, 32-36, 38, 39, 41-43, 45-52, 55-64, 69, 72, 73, 75-77, 80, 81, 85, 86, 89] |
| 13 | In patients with signs and symptoms compatible with COVID-19, should rapid antigen detection testing be used, | In patients with signs and symptoms compatible with COVID-19, we suggest the use of laboratory-based NAAT in saliva | Weak against | Very low | [6, 7, 23, 30, 40] |

	compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19 in saliva samples?	samples versus rapid antigen detection testing in saliva samples for diagnosis of COVID-19.			
14	In patients with signs and symptoms compatible with COVID-19, should rapid antigen detection testing be used, with standard NAAT (commercial and/or in house) testing for diagnosis of COVID-19 in samples other than nasopharyngeal sample and saliva?	In patients with signs and symptoms compatible with COVID-19, we suggest the use of laboratory-based NAAT in samples other than nasopharyngeal and saliva samples versus rapid antigen detection testing in samples other than nasopharyngeal and saliva samples for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 25, 31, 40, 44, 49, 54, 65-68, 71, 74, 78, 79, 84, 88]
15	In patients with signs and symptoms compatible with COVID-19 of equal or less than 7 days-onset , should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19?	In patients with signs and symptoms compatible with COVID-19 of equal or less than 7 days-onset, we suggest the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19.	Weak against	Very low	[6, 7, 23, 25, 28, 30, 33, 41, 49, 52, 59, 60, 68, 71-73, 78, 82, 84-86, 88]
16	In patients with signs and symptoms compatible with COVID-19 of more than 7 days-onset , should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19?	In patients with signs and symptoms compatible with COVID-19 of more than 7 days-onset, we suggest the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19.	Weak against	Very low	[6, 7, 23, 25, 28, 30, 49, 59, 60, 71, 73, 78, 84, 85]
17	In children <12 years with signs and symptoms compatible with COVID-19, should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in	In children <12 years old with signs and symptoms compatible with COVID-19, we suggest the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 52, 71, 79, 86]

house) for diagnosis of COVID-19?

- | | | | | | |
|----|---|--|----------------|----------|---|
| 18 | In patients ≥ 12 years old with signs and symptoms compatible with COVID-19, should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19? | In patients ≥ 12 years old with signs and symptoms compatible with COVID-19, we suggest the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19. | Weak against | Very low | [6, 23, 26, 29, 30, 32, 36, 45-47, 54-56, 68, 71, 76, 84, 88] |
| 19 | In patients with signs and symptoms compatible with COVID-19 at risk for severe illness , should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19? | In patients with signs and symptoms compatible with COVID-19 at risk for severe illness, we recommend the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19. | Strong against | Very low | [6, 23, 46, 60] |

Saliva sampling

- | | | | | | |
|----|---|---|--------------|----------|--------------------|
| 20 | In patients with signs and symptoms compatible with mild or moderate COVID-19, should saliva sampling be used, compared with nasopharyngeal swab sampling for diagnosis of COVID-19 with NAAT? | In patients with signs and symptoms compatible with mild or moderate COVID-19, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19. | Weak against | Very low | [90-97] |
| 21 | In patients with signs and symptoms compatible with severe or critical COVID-19, should saliva sampling be used, compared with nasopharyngeal swab sampling for diagnosis of COVID-19 with NAAT? | In patients with signs and symptoms compatible with severe or critical COVID-19 for the diagnosis of COVID-19 infection, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19. | Weak against | Very low | [13, 92, 98-116] |
| 22 | In patients with signs and symptoms compatible with | In patients with signs and symptoms compatible with | Weak for | Very low | [90, 95, 104, 108, |

	COVID-19 of equal or less than 7 days onset, should saliva sampling be used, compared with nasopharyngeal swab sampling for diagnosis of COVID-19 with NAAT?	COVID-19 of equal or less than 7 days-onset, we suggest the use of NAAT in saliva samples versus NAAT in nasopharyngeal swab samples for diagnosis of COVID-19.			109, 113, 114]
23	In patients with signs and symptoms compatible with COVID-19 of more than 7 days onset, should saliva sampling be used, compared with nasopharyngeal swab for diagnosis of COVID-19 with NAAT?	In patients with signs and symptoms compatible with COVID-19 of more than 7 days-onset, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[95, 98, 104, 108]
24	In children <12 years old with signs and symptoms compatible with COVID-19, should saliva sampling be used, compared with nasopharyngeal swab for diagnosis of COVID-19 with NAAT?	In children <12 years old with signs and symptoms compatible with COVID-19, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[90, 94, 95, 99, 103, 107, 108, 114, 115, 117]
25	In patients ≥12 years old with signs and symptoms compatible with COVID-19, should saliva sampling be used, compared with nasopharyngeal swab sampling for diagnosis of COVID-19 with NAAT?	In patients ≥12 years old with signs and symptoms compatible with COVID-19, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[13, 91-93, 96-98, 100-102, 104-106, 109-113, 115, 116, 118]
26	In patients with signs and symptoms compatible with COVID-19 at risk for severe illness, should saliva sampling be used, compared with nasopharyngeal swab sampling for diagnosis of COVID-19 with NAAT?	In patients with signs and symptoms compatible with COVID-19 at risk for severe illness, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[13, 90-118]

1 *Strength of recommendation (strong against, weak against, in research only, weak for, strong for).

2 **Overall certainty of the evidence (high, moderate, low, very low).

Journal Pre-proof

- 1 Table 2. PICO questions and recommendations in travellers, healthcare workers, and
 2 asymptomatic individuals at risk for exposure.

B	PICO question	Recommendation	Strength of recommendation*	Overall certainty of the evidence**	References
1	In travelers from areas with low prevalence , should surveys for contact history with known or suspected exposures to infected people followed by NAAT be used compared to universal NAAT to diagnose COVID-19?	In returning travelers from areas with low prevalence of COVID-19, we suggest the use of universal NAAT versus survey for contact history with known or suspected exposures in addition to NAAT for diagnosis of COVID-19.	Weak against	Very low	[119-121]
2	In travelers from areas with high prevalence , should surveys for contact history with known or suspected exposures to infected people followed by NAAT be used compared to universal NAAT to diagnose COVID-19?	In returning travelers from areas with high prevalence of COVID-19, we suggest the use of universal NAAT versus survey for contact history with known or suspected exposures in addition to NAAT for diagnosis of COVID-19.	Weak against	Very low	[121, 122]
3	In healthcare workers , should surveys for contact history with known or suspected exposures to infected people followed by NAAT be used compared to universal NAAT to diagnose COVID-19?	In healthcare workers, we recommend the use of universal NAAT versus survey for contact history with known or suspected exposures in addition to NAAT for diagnosis of COVID-19.	Strong against	Very low	[119, 121]
4	In asymptomatic populations at risk for exposure , should surveys for contact history with known or suspected exposures within less than 7 days to infected people followed by NAAT be used compared to universal NAAT to diagnose COVID-19?	In asymptomatic populations at risk for exposure, we suggest the use of universal NAAT versus survey for contact history with known or suspected exposures within less than 7 days in addition to NAAT for diagnosis of COVID-19.	Weak against	Very low	[119-122]
5	In asymptomatic populations at risk for exposure , should surveys for contact history with known or suspected exposures above 7 days to	In asymptomatic populations at risk for exposure, we suggest the use of universal NAAT versus survey for contact history with	Weak against	Very low	[119-122]

infected people followed known or suspected
by NAAT be used exposures above 7 days
compared to universal in addition to NAAT
NAAT to diagnose for diagnosis of
COVID-19? COVID-19.

- 1 *Strength of recommendation (strong against, weak against, in research only, weak for, strong for).
- 2 **Overall certainty of the evidence (high, moderate, low, very low).
- 3

- 1 Table 3. PICO questions and recommendations in asymptomatic individuals and those with
 2 close contact with an infected person.

C	PICO question	Recommendation	Strength of recommendation*	Overall certainty of the evidence**	References
Rapid antigen testing					
1	In asymptomatic children <12 years old without risk factors for severe COVID-19 should rapid antigen tests be used, as compared to laboratory-based NAAT to diagnose COVID-19?	In asymptomatic children <12 years old without risk factors for severe COVID-19, we suggest the use of laboratory-based NAAT versus rapid antigen testing for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 30, 33, 40, 51, 62, 67, 79, 83]
2	In asymptomatic patients ≥12 years old without risk factors for severe COVID-19 should rapid antigen tests be used, as compared to laboratory-based NAAT to diagnose COVID-19?	In asymptomatic patients ≥12 years old without risk factors for severe COVID-19, we suggest the use of laboratory-based NAAT versus rapid antigen testing for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 26, 27, 31, 45, 46, 48, 54-57, 60, 74, 75, 82]
3	In asymptomatic people of any age with any risk factor(s) for severe COVID-19 (including age <3 or ≥65 years) should rapid antigen tests be used, as compared to laboratory-based NAAT to diagnose COVID-19?	In asymptomatic people of any age with any risk factor(s) for severe COVID-19 (including age <3 months old or ≥65 years old), we suggest the use of laboratory based NAAT versus antigen testing for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 26, 27, 30, 31, 33, 40, 45, 46, 48, 51, 54-57, 60, 62, 67, 74, 75, 77-79, 82, 83]
4	In asymptomatic people should rapid antigen test be used as compared to laboratory-based NAAT in nasopharyngeal samples to diagnose COVID-19?	In asymptomatic people, we suggest the use of laboratory-based NAAT in nasopharyngeal samples versus rapid antigen testing in nasopharyngeal samples for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 26, 27, 33, 45, 46, 48, 51, 54-57, 60, 62, 75, 77, 82, 89]
5	In asymptomatic people should rapid antigen	In asymptomatic people, we suggest the use of	Weak against	Very low	[6, 23, 31, 40,

	test be used as compared to laboratory based NAAT in non-nasopharyngeal/non-saliva samples to diagnose COVID-19?	laboratory-based NAAT in non-nasopharyngeal/non-saliva samples versus rapid antigen testing in non-nasopharyngeal/non-saliva for diagnosis of COVID-19.			67, 74, 78, 79, 83]
6	In asymptomatic people should rapid antigen tests be used in saliva samples as compared laboratory based NAAT to diagnose COVID-19?	In asymptomatic people, we suggest the use of laboratory-based NAAT in saliva samples versus rapid antigen testing in saliva for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 30, 40]
NAAT in saliva samples					
7	In asymptomatic children <12 years , should NAAT in saliva samples be used, as compared to nasopharyngeal samples to diagnose COVID-19?	In asymptomatic children <12 years, we suggest the use of NAAT in saliva samples versus NAAT in nasopharyngeal swab samples for diagnosis of COVID-19.	Weak for	Very low	[108, 115, 123, 124]
8	In asymptomatic patients ≥12 years , should NAAT test in saliva samples be used, as compared to nasopharyngeal samples to diagnose COVID-19?	In asymptomatic patients ≥12 years, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[96, 97, 105, 115, 125]
9	In close contacts asymptomatic children <12 years , should NAAT in saliva samples be used, as compared to nasopharyngeal samples to diagnose COVID-19?	In close contact asymptomatic children <12 years, we suggest NAAT in saliva samples be used compared to NAAT testing in nasopharyngeal swab samples for diagnosis of COVID-19.	Weak for	Very low	[126]
10	In close contacts asymptomatic patients ≥12 years , should NAAT in saliva samples be used, as compared to nasopharyngeal samples to diagnose COVID-19?	In close contact asymptomatic patients ≥12 years, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[96, 105, 115, 124, 125]

- | | | | | | |
|----|--|---|--------------|----------|----------------------|
| 11 | In close contacts asymptomatic children <12 years with less than 7 days since contact , should NAAT in saliva samples be used, as compared to nasopharyngeal samples to diagnose COVID-19? | In close contacts asymptomatic children <12 years with less than 7 days since contact, we suggest the use of NAAT in saliva samples versus NAAT in nasopharyngeal swab samples for diagnosis of COVID-19. | Weak for | Very low | [108, 115, 123, 124] |
| 12 | In close contacts asymptomatic patients ≥12 years with less than 7 days since contact , should NAAT in saliva samples be used, as compared to nasopharyngeal samples to diagnose COVID-19? | In close contact asymptomatic patients ≥12 years with less than 7 days since contact, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19. | Weak against | Very low | [96, 97, 105, 115] |

1 *Strength of recommendation (strong against, weak against, in research only, weak for, strong for).

2 **Overall certainty of the evidence (high, moderate, low, very low).

3

4

1 Table 1. PICO questions and recommendations in patients with signs and symptoms of COVID-19.

A	PICO question	Recommendation	Strength of recommendation*	Overall certainty of the evidence**	References
Rapid NAAT					
1	In patients with signs and symptoms compatible with mild or moderate COVID-19, should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19?	In patients with signs and symptoms compatible with mild or moderate COVID-19 we suggest the use of rapid NAAT versus laboratory based NAAT testing for the diagnosis of COVID-19.	Weak for	Very low	[12-23].
2	In patients with signs and symptoms compatible with severe or critical COVID-19, should commercial rapid NAAT testing be used, compared with standard NAAT (commercial and/or in house) testing for diagnosis of COVID-19?	In patients with signs and symptoms compatible with severe or critical COVID-19 we suggest the use of rapid NAAT versus laboratory based NAAT for the diagnosis of COVID-19.	Strong for	Very low	[12-23]
3	In patients with signs and symptoms compatible with COVID-19, should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19 in nasopharyngeal samples?	In patients with signs and symptoms compatible with COVID-19 we suggest the use of rapid NAAT in nasopharyngeal samples versus laboratory based NAAT in nasopharyngeal samples for the diagnosis of COVID-19.	Weak for	Very low	[13, 15, 16, 18, 20, 22, 23]
4	In patients with signs and symptoms compatible with COVID-19, should commercial rapid NAAT be used, compared with the standard NAAT (commercial and/or in house) for diagnosis of COVID-19 in samples other than nasopharyngeal swab?	In patients with signs and symptoms compatible with COVID-19 we suggest the use of rapid NAAT in samples other than nasopharyngeal swab versus laboratory based NAAT in samples other than nasopharyngeal swab for the diagnosis of COVID-19, when allowed by regulatory boards and manufacturer's instructions.	Weak for	Very low	[12, 14, 15, 17, 21, 23]

- | | | | | | |
|------------------------------|--|--|--------------|----------|-------------------------|
| 5 | In patients with signs and symptoms compatible with COVID-19 of equal or less than 7 days-onset , should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19? | In patients with signs and symptoms compatible with COVID-19 of equal or less than 7 days-onset we suggest the use of rapid NAAT versus laboratory based NAAT for the diagnosis of COVID-19. | Weak for | Very low | [12-22] |
| 6 | In patients with signs and symptoms compatible with COVID-19 of more than 7 days-onset , should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19? | In patients with signs and symptoms compatible with COVID-19 of more than 7 days-onset we suggest the use of rapid NAAT versus laboratory based NAAT for the diagnosis of COVID-19. | Weak for | Very low | [12-23] |
| 7 | In children <12 years old with signs and symptoms compatible with COVID-19, should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19? | In children <12 years old with signs and symptoms compatible with COVID-19 we suggest the use of rapid NAAT versus laboratory based NAAT for diagnosis of COVID-19. | Weak for | Very low | [12, 15, 17-20, 22, 23] |
| 8 | In patients ≥12 years old with signs and symptoms compatible with COVID-19, should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19? | In patients ≥12 years old with signs and symptoms compatible with COVID-19, we suggest the use of rapid NAAT versus laboratory based NAAT for diagnosis of COVID-19. | Weak for | Very low | [13, 14, 16, 21, 23] |
| 9 | In patients with signs and symptoms compatible with COVID-19 at risk for severe illness , should commercial rapid NAAT be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19? | In patients with signs and symptoms compatible with COVID-19 at risk for severe disease, we suggest the use of rapid NAAT versus laboratory-based NAAT for diagnosis of COVID-19. | Weak for | Very low | [12-23] |
| Rapid antigen testing | | | | | |
| 10 | In patients with signs and symptoms compatible with mild or moderate COVID-19, should rapid antigen detection testing be used, compared with standard NAAT | In patients with mild and moderate COVID-19, we suggest the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19. | Weak against | Very low | [6, 23-34] |

(commercial and/or in house) for diagnosis of COVID-19?

- | | | | | | |
|----|---|---|----------------|----------|--|
| 11 | In patients with signs and symptoms compatible with severe or critical COVID-19, should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in house) testing for diagnosis of COVID-19? | In patients with severe or critical COVID-19, we recommend the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19. | Strong against | Very low | [6, 7, 23, 30, 35-88] |
| 12 | In patients with signs and symptoms compatible with COVID-19, should rapid antigen detection testing be used, compared with the standard NAAT (commercial and/or in house) for diagnosis of COVID-19 in nasopharyngeal samples ? | In patients with signs and symptoms compatible with COVID-19, we suggest the use of laboratory based NAAT in nasopharyngeal samples versus rapid antigen detection testing in nasopharyngeal samples for diagnosis of COVID-19. | Weak against | Very low | [6, 7, 23, 24, 26-29, 32-36, 38, 39, 41-43, 45-52, 55-64, 69, 72, 73, 75-77, 80, 81, 85, 86, 89] |
| 13 | In patients with signs and symptoms compatible with COVID-19, should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19 in saliva samples ? | In patients with signs and symptoms compatible with COVID-19, we suggest the use of laboratory-based NAAT in saliva samples versus rapid antigen detection testing in saliva samples for diagnosis of COVID-19. | Weak against | Very low | [6, 7, 23, 30, 40] |
| 14 | In patients with signs and symptoms compatible with COVID-19, should rapid antigen detection testing be used, with standard NAAT (commercial and/or in house) testing for diagnosis of COVID-19 in samples other than nasopharyngeal sample and saliva ? | In patients with signs and symptoms compatible with COVID-19, we suggest the use of laboratory-based NAAT in samples other than nasopharyngeal and saliva samples versus rapid antigen detection testing in samples other than nasopharyngeal and saliva samples for diagnosis of COVID-19. | Weak against | Very low | [6, 23, 25, 31, 40, 44, 49, 54, 65-68, 71, 74, 78, 79, 84, 88] |
| 15 | In patients with signs and symptoms compatible with COVID-19 of equal or less than 7 days-onset , should rapid antigen detection testing be used, compared with | In patients with signs and symptoms compatible with COVID-19 of equal or less than 7 days-onset, we suggest the use of laboratory-based NAAT | Weak against | Very low | [6, 7, 23, 25, 28, 30, 33, 41, 49, 52, 59, 60, 68, 71-73, 78, 82, 84-86, 88] |

standard NAAT (commercial and/or in house) for diagnosis of COVID-19? versus rapid antigen detection testing for diagnosis of COVID-19.

- | | | | | | |
|----|--|--|----------------|----------|---|
| 16 | In patients with signs and symptoms compatible with COVID-19 of more than 7 days-onset , should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19? | In patients with signs and symptoms compatible with COVID-19 of more than 7 days-onset, we suggest the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19. | Weak against | Very low | [6, 7, 23, 25, 28, 30, 49, 59, 60, 71, 73, 78, 84, 85] |
| 17 | In children <12 years with signs and symptoms compatible with COVID-19, should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19? | In children <12 years old with signs and symptoms compatible with COVID-19, we suggest the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19. | Weak against | Very low | [6, 23, 52, 71, 79, 86] |
| 18 | In patients ≥12 years old with signs and symptoms compatible with COVID-19, should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19? | In patients ≥ 12 years old with signs and symptoms compatible with COVID-19, we suggest the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19. | Weak against | Very low | [6, 23, 26, 29, 30, 32, 36, 45-47, 54-56, 68, 71, 76, 84, 88] |
| 19 | In patients with signs and symptoms compatible with COVID-19 at risk for severe illness , should rapid antigen detection testing be used, compared with standard NAAT (commercial and/or in house) for diagnosis of COVID-19? | In patients with signs and symptoms compatible with COVID-19 at risk for severe illness, we recommend the use of laboratory-based NAAT versus rapid antigen detection testing for diagnosis of COVID-19. | Strong against | Very low | [6, 23, 46, 60] |

Saliva sampling

- | | | | | | |
|----|---|---|--------------|----------|---------|
| 20 | In patients with signs and symptoms compatible with mild or moderate COVID-19, should saliva sampling be used, compared with nasopharyngeal swab sampling for diagnosis of COVID-19 with NAAT? | In patients with signs and symptoms compatible with mild or moderate COVID-19, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19. | Weak against | Very low | [90-97] |
|----|---|---|--------------|----------|---------|

21	In patients with signs and symptoms compatible with severe or critical COVID-19, should saliva sampling be used, compared with nasopharyngeal swab sampling for diagnosis of COVID-19 with NAAT?	In patients with signs and symptoms compatible with severe or critical COVID-19 for the diagnosis of COVID-19 infection, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[13, 92, 98-116]
22	In patients with signs and symptoms compatible with COVID-19 of equal or less than 7 days onset , should saliva sampling be used, compared with nasopharyngeal swab sampling for diagnosis of COVID-19 with NAAT?	In patients with signs and symptoms compatible with COVID-19 of equal or less than 7 days-onset, we suggest the use of NAAT in saliva samples versus NAAT in nasopharyngeal swab samples for diagnosis of COVID-19.	Weak for	Very low	[90, 95, 104, 108, 109, 113, 114]
23	In patients with signs and symptoms compatible with COVID-19 of more than 7 days onset , should saliva sampling be used, compared with nasopharyngeal swab for diagnosis of COVID-19 with NAAT?	In patients with signs and symptoms compatible with COVID-19 of more than 7 days-onset, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[95, 98, 104, 108]
24	In children <12 years old with signs and symptoms compatible with COVID-19, should saliva sampling be used, compared with nasopharyngeal swab for diagnosis of COVID-19 with NAAT?	In children <12 years old with signs and symptoms compatible with COVID-19, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[90, 94, 95, 99, 103, 107, 108, 114, 115, 117]
25	In patients ≥12 years old with signs and symptoms compatible with COVID-19, should saliva sampling be used, compared with nasopharyngeal swab sampling for diagnosis of COVID-19 with NAAT?	In patients ≥12 years old with signs and symptoms compatible with COVID-19, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[13, 91-93, 96-98, 100-102, 104-106, 109-113, 115, 116, 118]
26	In patients with signs and symptoms compatible with COVID-19 at risk for severe illness , should saliva sampling be used, compared with nasopharyngeal swab sampling for diagnosis of COVID-19 with NAAT?	In patients with signs and symptoms compatible with COVID-19 at risk for severe illness, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[13, 90-118]

1 *Strength of recommendation (strong against, weak against, in research only, weak for, strong for).

1 **Overall certainty of the evidence (high, moderate, low, very low).

2

Journal Pre-proof

1 Table 2. PICO questions and recommendations in travellers, healthcare workers, and asymptomatic individuals at risk for exposure.

B	PICO question	Recommendation	Strength of recommendation*	Overall certainty of the evidence**	References
1	In travelers from areas with low prevalence , should surveys for contact history with known or suspected exposures to infected people followed by NAAT be used compared to universal NAAT to diagnose COVID-19?	In returning travelers from areas with low prevalence of COVID-19, we suggest the use of universal NAAT versus survey for contact history with known or suspected exposures in addition to NAAT for diagnosis of COVID-19.	Weak against	Very low	[119-121]
2	In travelers from areas with high prevalence , should surveys for contact history with known or suspected exposures to infected people followed by NAAT be used compared to universal NAAT to diagnose COVID-19?	In returning travelers from areas with high prevalence of COVID-19, we suggest the use of universal NAAT versus survey for contact history with known or suspected exposures in addition to NAAT for diagnosis of COVID-19.	Weak against	Very low	[121, 122]
3	In healthcare workers , should surveys for contact history with known or suspected exposures to infected people followed by NAAT be used compared to universal NAAT to diagnose COVID-19?	In healthcare workers, we recommend the use of universal NAAT versus survey for contact history with known or suspected exposures in addition to NAAT for diagnosis of COVID-19.	Strong against	Very low	[119, 121]
4	In asymptomatic populations at risk for exposure , should surveys for contact history with known or suspected exposures within less than 7 days to infected people followed by NAAT be used compared to universal NAAT to diagnose COVID-19?	In asymptomatic populations at risk for exposure, we suggest the use of universal NAAT versus survey for contact history with known or suspected exposures within less than 7 days in addition to NAAT for diagnosis of COVID-19.	Weak against	Very low	[119-122]
5	In asymptomatic populations at risk for exposure , should surveys for contact history with known or suspected exposures above 7 days to infected people followed by NAAT be used compared to universal NAAT to diagnose COVID-19?	In asymptomatic populations at risk for exposure, we suggest the use of universal NAAT versus survey for contact history with known or suspected exposures above 7 days in	Weak against	Very low	[119-122]

addition to NAAT for diagnosis of COVID-19.

1 *Strength of recommendation (strong against, weak against, in research only, weak for, strong for).

2 **Overall certainty of the evidence (high, moderate, low, very low).

3

1 Table 3. PICO questions and recommendations in asymptomatic individuals and those with close contact with an infected person.

C	PICO question	Recommendation	Strength of recommendation*	Overall certainty of the evidence**	References
Rapid antigen testing					
1	In asymptomatic children <12 years old without risk factors for severe COVID-19 should rapid antigen tests be used, as compared to laboratory-based NAAT to diagnose COVID-19?	In asymptomatic children <12 years old without risk factors for severe COVID-19, we suggest the use of laboratory-based NAAT versus rapid antigen testing for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 30, 33, 40, 51, 62, 67, 79, 83]
2	In asymptomatic patients ≥12 years old without risk factors for severe COVID-19 should rapid antigen tests be used, as compared to laboratory-based NAAT to diagnose COVID-19?	In asymptomatic patients ≥12 years old without risk factors for severe COVID-19, we suggest the use of laboratory-based NAAT versus rapid antigen testing for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 26, 27, 31, 45, 46, 48, 54-57, 60, 74, 75, 82]
3	In asymptomatic people of any age with any risk factor(s) for severe COVID-19 (including age <3 or ≥65 years) should rapid antigen tests be used, as compared to laboratory-based NAAT to diagnose COVID-19?	In asymptomatic people of any age with any risk factor(s) for severe COVID-19 (including age <3 months old or ≥65 years old), we suggest the use of laboratory based NAAT versus antigen testing for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 26, 27, 30, 31, 33, 40, 45, 46, 48, 51, 54-57, 60, 62, 67, 74, 75, 77-79, 82, 83]
4	In asymptomatic people should rapid antigen test be used as compared to laboratory-based NAAT in nasopharyngeal samples to diagnose COVID-19?	In asymptomatic people, we suggest the use of laboratory-based NAAT in nasopharyngeal samples versus rapid antigen testing in nasopharyngeal samples for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 26, 27, 33, 45, 46, 48, 51, 54-57, 60, 62, 75, 77, 82, 89]

5	In asymptomatic people should rapid antigen test be used as compared to laboratory based NAAT in non-nasopharyngeal/non-saliva samples to diagnose COVID-19?	In asymptomatic people, we suggest the use of laboratory-based NAAT in non-nasopharyngeal/non-saliva samples versus rapid antigen testing in non-nasopharyngeal/non-saliva for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 31, 40, 67, 74, 78, 79, 83]
6	In asymptomatic people should rapid antigen tests be used in saliva samples as compared laboratory based NAAT to diagnose COVID-19?	In asymptomatic people, we suggest the use of laboratory-based NAAT in saliva samples versus rapid antigen testing in saliva for diagnosis of COVID-19.	Weak against	Very low	[6, 23, 30, 40]
NAAT in saliva samples					
7	In asymptomatic children <12 years , should NAAT in saliva samples be used, as compared to nasopharyngeal samples to diagnose COVID-19?	In asymptomatic children <12 years, we suggest the use of NAAT in saliva samples versus NAAT in nasopharyngeal swab samples for diagnosis of COVID-19.	Weak for	Very low	[108, 115, 123, 124]
8	In asymptomatic patients ≥12 years , should NAAT test in saliva samples be used, as compared to nasopharyngeal samples to diagnose COVID-19?	In asymptomatic patients ≥12 years, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[96, 97, 105, 115, 125]
9	In close contacts asymptomatic children <12 years , should NAAT in saliva samples be used, as compared to nasopharyngeal samples to diagnose COVID-19?	In close contact asymptomatic children <12 years, we suggest NAAT in saliva samples be used compared to NAAT testing in nasopharyngeal swab samples for diagnosis of COVID-19.	Weak for	Very low	[126]
10	In close contacts asymptomatic patients ≥12 years , should NAAT in saliva samples be used, as compared to nasopharyngeal samples to diagnose COVID-19?	In close contact asymptomatic patients ≥12 years, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19.	Weak against	Very low	[96, 105, 115, 124, 125]
11	In close contacts asymptomatic children <12 years with less than 7 days since contact , should NAAT in	In close contacts asymptomatic children <12 years with less than 7 days since contact, we suggest the use of NAAT in saliva samples versus	Weak for	Very low	[108, 115, 123, 124]

saliva samples be used, as compared to NAAT in nasopharyngeal swab samples for nasopharyngeal samples to diagnose COVID-19? diagnosis of COVID-19.

- | | | | | | |
|----|--|--|--------------|----------|--------------------|
| 12 | In close contacts asymptomatic patients ≥ 12 years with less than 7 days since contact , should NAAT in saliva samples be used, as compared to nasopharyngeal samples to diagnose COVID-19? | In close contact asymptomatic patients ≥ 12 years with less than 7 days since contact, we suggest the use of NAAT in nasopharyngeal swab samples versus NAAT in saliva samples for diagnosis of COVID-19. | Weak against | Very low | [96, 97, 105, 115] |
|----|--|--|--------------|----------|--------------------|

1 *Strength of recommendation (strong against, weak against, in research only, weak for, strong for).

2 **Overall certainty of the evidence (high, moderate, low, very low).

3

4