



Screening for neurocognitive deficits in adult populations reliant on toxic cassava as the main source of food

Dépistage des déficits neurocognitifs dans les populations adultes dépendant du manioc toxique comme principale source de denrées alimentaires

Victor Hutu Kabamba¹, Daniel Okitundu Luwa¹, Jean-Pierre Mayambu Banea^{2†}, Bruno Giordani³, Michael Joseph Boivin⁴, Dieudonné Mumba-Ngoyi⁵, Désiré Tshala-Katumbay^{1,5,6}

Corresponding Author

Victor Hutu Kabamba, MD
Department of Neurology, University of Kinshasa, Democratic Republic of Congo.
Phone : + 243 (0)998264873
E-mail : docteurhkabamba@yahoo.fr

Résumé

Contexte & objectif. La dépendance alimentaire chronique au manioc cyanogène mal détoxifié est largement répandue en Afrique subsaharienne. L'objectif de la présente étude était de dépister les troubles neurocognitifs et le fonctionnement de la vie quotidienne chez les adultes avec dépendance alimentaire au manioc cyanogène comme principale source de nourriture. **Méthodes.** C'était une étude transversale analytique incluant les chefs de ménage vivant dans le district rural de Kahemba, en République démocratique du Congo. Les participants ont été dépistés pour les troubles neurocognitifs à l'aide de Community Screening Interview for Dementia (CSID). L'exposition au manioc cyanogène a été déterminée par les concentrations urinaires de thiocyanate (SCN). Des modèles de régression ont été utilisés pour identifier les prédicteurs de mauvaises performances au CSID. **Résultats.** Quatre cent et six chefs de ménage (203 couples, âge moyen $38,4 \pm 11,4$ ans) ont été enrôlés. Cent trente-six (33,5 %) [69 femmes et 67 hommes, âge moyen $39 \pm 14,4$ ans et 13 (3,2 %) [7 femmes et 6 hommes, âge moyen $32 \pm 2,6$ ans,) remplissaient respectivement les critères de trouble neurocognitif léger (TNL) et de trouble neurocognitif majeur (TNM). La concentration urinaire Moyenne globale de SCN était de $949,5 \pm 518,3 \mu\text{mol/L}$. Après ajustement pour l'âge, le sexe, l'état nutritionnel et les antécédents de konzo, les déficits spécifiques aux différents domaines de la neuro-cognition étaient significativement associés de manière

Summary

Context and objective. Chronic dietary reliance on improperly processed cyanogenic toxic cassava is widespread in sub-Saharan Africa. The objective of the present study was to screen for neurocognition impairments and daily-life functioning in adults with dietary dependency on cyanogenic cassava as the main source of food. **Methods.** A cross-sectional design enrolled heads of households (in couples) in the rural district of Kahemba, Democratic Republic of Congo. Participants were screened for neurocognitive impairments using the Community Screening Interview for Dementia (CSID). Detailed neuropsychiatric evaluations were performed and disease entities classified according to the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) criteria when applicable. Cassava cyanogenic exposure was ascertained by urinary concentrations of thiocyanate (SCN). Regression models were used to identify predictors of CSID performance at the 0.05 significance level. **Results.** For hundred and six households (203 couples, mean age 38.4 ± 11.4 years) were involved. One hundred thirty-six subjects (33.5 %) [69 women and 67 men, mean age 39 ± 14.4 years] and 13 (3.2 %) [7 women and 6 men, mean age: 32 ± 2.6 years] fulfilled the criteria for mild cognitive impairment (MCI) and Major Neurocognitive disorder (MNCD), respectively. The overall mean urinary concentration of SCN was $949.5 \pm 518.3 \mu\text{mol/L}$. After adjusting for age, gender, nutritional status, and history of konzo, neurocognition domain-specific deficits were independently associated with either hypertension or USCN ($350 \mu\text{mol/l}$ incremental increase in excretion). Functional impairments in daily-life activities increased as subjects poorly performed at the CSID screening (Spearman $r = -.2$, $p < 0.01$). **Conclusion.** Neurocognitive deficits in adults are



indépendante à l'hypertension ou à l'USCN (350 $\mu\text{mol/L}$, augmentation incrémentale de l'excrétion). Les déficiences fonctionnelles dans les activités de la vie quotidienne ont été directement proportionnelles à la mauvaise performance au CSID (Spearman $r = -0,2$, $p < 0,01$). Conclusion. Les déficits neurocognitifs sont courants chez les adultes congolais qui dépendent du manioc cyanogène comme principale source de nourriture. Ce qui justifie la réalisation d'études ultérieures pour élucider l'impact cérébral / comportemental global ainsi que les mécanismes de la toxicité du manioc chez les adultes ayant une dépendance alimentaire au manioc cyanogène.

Mots-clés: Manioc; Troubles neurocognitifs; Cyanure; Fonctionnement quotidien; Hypertension

Reçu le 5 juillet 2022, Accepté le 5 novembre 2022

1 Department of Neurology, University of Kinshasa, Democratic Republic of Congo

2 Nutrition National Program & School of Public Health, Ministry of Health, Democratic Republic of Congo; †In memoriam

3 Departments of Psychiatry, Neurology, and Psychology and School of Nursing, University of Michigan, Ann Arbor MI, USA

4 Department of Psychiatry and Neurology & Ophthalmology, Michigan State University, East Lansing MI, USA

5 Department of Tropical Medicine, University of Kinshasa & Institute National de Recherche Biomédicale, Democratic Republic of Congo

6 Department of Neurology, Oregon Health & Science University, Portland OR, USA

Impoverished populations in rural Africa are at risk for the aforementioned neurological diseases because of their dietary dependency on food from drought-tolerant varieties of cassava a.k.a tapioca or yucca, which contain high concentrations of two cyanogenic glucosides notably linamarin and lotaustralin, in a ~97:3 concentration ratio (11-12). Overt diseases e.g. konzo or TAN surface when toxic thresholds are exceeded and subjects have limited detoxification capabilities because of low dietary protein intake (13). Cassava toxicity is commonly seen in times of famine often caused by flood, drought, pestilence, or war. In these times, poor populations are forced

common in Congolese adults relying on cyanogenic cassava as the main source of food. Our study findings warrant further studies to elucidate the overall lifespan brain/behavioral burden and mechanisms of cassava toxicity among adults with dietary dependency on cyanogenic cassava as the main source of food.

Keywords: Cassava; Cognitive; Impairments; Cyanide; Daily Functioning; Hypertension

Received: July 5th, 2022

Accepted: November 5th, 2022

<https://dx.doi.org/10.4314/aamed.v16i2.2>

Introduction

Chronic reliance on cyanogenic cassava in the settings of hunger, inadequate detoxification of its tubers, or yet to be fully documented individual susceptibility, can lead to neurological impairments and disability. These include acute encephalopathy, sub-acute spastic para paresis (a.k.a. konzo), tropical ataxic neuropathy (TAN), and/or, reportedly, a cerebellar-parkinsonism-dementia syndrome (CPDS) (1-3). Outbreaks of konzo have since occurred in several countries of sub-Saharan Africa including Angola, Cameroon, Central African Republic, Democratic Republic of the Congo (DRC), Tanzania, Uganda, Mozambique, and recently in Zambia (4-6). Konzo was named after a local designation in *kiyaka*, a local language spoken in DRC and means “tied legs” in reference to the scissoring spastic gait of affected subjects (6). Children and women of childbearing age are at higher risk for reasons that have yet to be known (8-10).

to adopt shortcuts in processing methods including a reduction in cassava processing time. Under these conditions, higher residual amounts of cyanogenic compounds are ingested and first converted into cyanide, a highly toxic compound. Naturally, defense mechanisms would then convert cyanide into the purportedly less toxic thiocyanate (SCN) which in turn is excreted in urine making it a suitable marker of food (cassava) cyanogenic exposure (12).

Neurological effects of food dependency on cassava represent a significant contribution to the global burden of disease. Unofficial estimates indicate that up to 100,000 cases of konzo may



have already occurred across the African continent (14). We also showed that cognition may be impaired in addition to the paralysis in children with konzo and those with no overt paralysis still perform worse at neuropsychological testing compared to children from non-konzo areas suggesting that pervasive cognition deficits have commonplace in konzo-affected areas (10, 15-16). Whether poor cognition may extend into adolescent and adult life impacting functioning and economic productivity still needs to be determined. In this study, we screened for dementia in the adult population of Kahemba, a konzo heavily affected district of the Democratic Republic of Congo (DRC), which relies on cassava farming for its subsistence (7, 9) and draw attention to a high prevalence of mild cognitive impairments (MCI) in this population.

Methods

Study Population. The present study was carried out in the district of Kahemba, southern Bandundu, nowadays the Kwango Province of the DRC. Residents of Kahemba heavily rely on cassava farming and possess very limited livestock for subsistence. During the last decade, Kahemba has faced a severe humanitarian crisis due to population displacement following the end of the Angola war. Because of food shortages, residents were forced to adopt shortcuts in cassava processing methods, a phenomenon that led to food cyanogenic exposure and poisonings. A sudden rise in the number of cases with konzo was since documented with a point-prevalence of the disease as high as 20% in certain villages (17).

Study Design. We carried out a cross-sectional study that consecutively enrolled household consenting members. Source population: these subjects are parents of children enrolled in previously studies that assessed cassava toxicity and neurodevelopmental outcomes in Kahemba (10). Inclusion criteria: only consented heads of households and couples who were permanent residents of Kahemba participated were included. The vast majority of the subjects (96 %) lived in Kahemba prior to the 2009 major outbreak of konzo (7). Exclusion criteria: subjects with a

history of neurological disability and/or mental illnesses were excluded from the study.

Data collection tools and parameters of interest

A questionnaire was used to collect data on sociodemographic characteristics, medical history, and current health status. Cognitive impairment, anxiety and/or depression, neurological evaluation, and cassava cyanogenic exposure, were assessed using tools that are outlined below. Cognition status was the main outcome variable, of which changes were assessed in relation to sociodemographic, clinical status, nutrition and cyanogenic exposure levels.

Cognitive Status, Functioning, and Depression Screening

The community screening interview for dementia (CSID) developed to screen for dementia across cultures and resource settings was used for study purposes (18). The CSID consists of a set of direct cognitive screening tasks for non-literate and literate populations, in addition to an assessment of the everyday functional status through an interview with a close informant. An overall cognitive score is calculated which ranges from 0 to 34 with lower scores indicating cognitive impairment. A cutoff point of CSID score < 25 was considered as the threshold for cognitive impairment in this study (19-20). The informant score ranges from 0 to 40 with scores > 16 indicating functional impairments in everyday life activities. Study participants were further examined by a certified Congolese neuropsychiatrist and classified as having either normal cognition, or mild cognitive impairment (MCI), or dementia based on the criteria listed in the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV). Signs indicative of depression and/or anxiety were assessed through a structured interview using the Goldberg Depressive Anxiety Scale (GDAS), an instrument previously used in the DRC (21); a GDAS score ≥ 22 was indicative of the depressed mood. The severity of the tendency to depression was graded as mild (scores 22 - 35), moderate (scores 36 - 53), or severe (scores ≥ 54).

Neurological Evaluation. A Congolese board-certified neuropsychiatrist performed



neurological examinations on all subjects to assess cranial nerves, motor and sensory functions as well as reflexes and sphincter control. The diagnosis of konzo (cassava-associated spastic paraparesis) was based on the following WHO criteria for the disease: (1) a visible symmetric spastic abnormality of gait while walking or running; (2) a history of onset of less than 1 week followed by a non-progressive course in a formerly healthy person; and (3) bilaterally exaggerated knee or ankle jerks without signs of disease of the spine (22). Anthropometric measurements (height, weight, mid-upper arm circumference) were also made at the time of the neurological evaluation to determine nutrition status as a function on body mass index (BMI).

Ascertainment of Cassava Cyanogenic Exposure
During the initial visit to Kahemba, samples of cassava flour from 18 consenting households were collected and found to have cyanide concentrations of 30 to 200 ppm, well-above the 10-ppm safe limit proposed by the WHO (23). We have previously reported higher levels of urinary SCN in children with konzo relative to their respective controls from our study area (15, 24). In the present study, urine samples from study participants were analyzed using the same previously used urinary SCN picrate measurement D1kit. In this analysis, a color chart is used with 10 shades of color from yellow to brown, corresponding to 0–100 mg SCN/l urine (ppm). The results in ppm are then multiplied by 17.2 to convert to $\mu\text{mol SCN} / \text{l}$ according the manufacturer protocol (25).

Ethics statement

Informed consent was obtained verbally by investigators fluent in Lingala and/or Kikongo, the local spoken languages. The research protocol was approved by the Institutional Review Board of the DRC Ministry of Health.

Statistical Analyses

Initial analyses consisted of Student t-tests, as well as Chi-square tests to compare key clinical and biological characteristics across study groups as deemed appropriate. Logistic regression was first used to assess relationships between

neurocognitive status (CSID normal score ≥ 25 vs. CSID poor score < 25) and age, gender, body mass index ($\leq 18.4 \text{ kg/m}^2$ for malnutrition), systolic and diastolic blood pressure, glycemia, smoking, alcohol consumption, depression, functioning, neurological and konzo status, and urinary concentrations in SCN as independent variables. Predictors significantly associated to the main outcome at the .10 significance threshold were then retained in models for multivariable analyses that assessed the aforementioned associations after adjusting for age, gender, nutrition status, exposure levels, konzo status and/or hypertension at the .05 significance level. All analyses were carried out using the STATA software (STATA, STATA inc. version 16.1).

Results

Sociodemographic Characteristics of Study Subjects


Of the 406 study participants (203 household couples; men/women ratio: 1/1), the majority (42%) were aged between 31 and 45 years regardless of gender and the 61 – 75 age group (12 subjects) was the least represented group making 3% of the study subjects. Most subjects (84.7 %, overall) had post-elementary education regardless of gender. The average (SD) duration of stay in Kahemba was 33.3 (13.9) years and most subjects (96.6 %) have been living in Kahemba prior to the major 2019 konzo outbreak.

Clinical and Biological Characteristics of Study Subjects

More than half of the subjects (54%, overall) suffered chronic undernutrition. Alcohol consumption was reported in 267 (65.8 %) subjects. Overall, fewer subjects (6.7 %) had hypertension, hyperglycemia (13.5 %), and smoking habits (5.9%). Subjects were heavily exposed to cassava cyanogens as indicated by concentrations of urinary SCN above $350 \mu\text{mol/l}$ in the majority (78%) of subjects regardless of gender. Of the total 406 study participants, 240 (59.1 %) subjects had at least one subject with konzo in their households while 38 (9.4 %) (26 men and 12 women) had konzo themselves.



CSID performance scores followed a normal distribution (**Figure 1**). Depressed mood was reported in 258 (63.5 %) of the study participants, almost exclusively in the group that performed well at the CSID screening. Of these, 135 were prone to mild depression, 97 to moderate, 26 to severe depression as per the Golberg scale.



One Hundred forty-nine (36.7 %) subjects poorly performed at the CSID screening. One hundred thirty-six (33.5 %) [69 women and 67 men, mean

(SD) age: 39 (14.4), range 18 – 73 years) and 13 (3.2 %) [7 women and 6 men, mean (SD) age: 32 (2.6) years, range 27 - 36) fulfilled the criteria for MCI and dementia, respectively. Key clinical and biological characteristics across study groups are summarized in **Figure 2** and **Table 1**.

Figure 1. Performance scores at the CSID screening followed a normal distribution regardless of gender.




Figure 2. Patterns of neurocognition status showed several domains affected at neurological screening.

Clinical and Biological Predictors of CSID status

Increases in urinary thiocyanate excretion (by 350 micromol/l and HT were associated with domain-specific neurocognitive impairments. By removing the HT (predictor of non-amnesic multidomain deficits) from the model, increase in urinary thiocyanate excretion remained a significant predictor of most deficits except for the non-amnesic unidomain impairment (Table 2).

Table 2. Clinical and Biological Predictors of Neurocognition Status (Univariable multimodal logistical regression of risk factors of neurocognitive impairment by type of disorder)

Predictors	MCI (n=136)				MNCD (n=13)
	Non amnesic		Amnésic		
	Single unidomain OR (95% CI)	Multidomain OR (95% CI)	Single unidomain OR (95% CI)	Multidomain OR (95% CI)	
Age/year	0.96 (0.89-.1.04)	0.9 (0.9 -1.03)	0.93 (0.77-1.12)	1.00 (0.97-1.04)	0.93 (0.82 -1.04)
Men	0.69 (0.15-3.15)	1.5 (0.6 -3.5)	1.37 (0.9-0.0)	0.79 (0.37-1.69)	0.46 (0.07-3.29)
Undernutrition (BMI ≤ 18.4 kg/m ²)	0.24 (0.04 -1.33)	1.9 (0.7 -4.6)	0.41 (0.02-7.79)	1.01 (0.47-2.20)	0.39 (0.06-2.62)
Thiocyanate /350 μmol/l	0.49 (0.2 -1.07)	2.3 (1.6 -3.2)	0.02 (0.00-0.96)	0.70 (0.5-0.97)	2.18 (1.07-4.45)
Konzo in households	2.04 (0.37 - 11.14)	0.72 (0.3-1.6)	18.56 (0.7- 437.0)	2.12 (0.93-4.87)	0.43 (0.06 - 2.99)
Hypertension	--	7.7 (1.5-39.0)	71.7 (0.1-4742.6)	4.43 (0.85-22.99)	--
By moving out HT in the model					
Age/year	0.96 (0.89-.1.04)	1.00 (0.9-1.04)	0.98 (0.87-1.12)	1.01 (0.98-1.04)	0.93 (0.82 -1.04)

Ann. Afr. Med., vol. 16, n° 2, Mars 2023

This is an open article distributed under the terms of the Creative Commons Attribution Non-Commercial License

(<http://creativecommons.org/licenses/bync/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited



Men	0.67 (0.15-3.06)	1.4 (0.6 -3.34)	-	0.81 (0.38-1.73)	0.47 (0.07-3.32)
Undernutrition (BMI $\leq 18.4 \text{ kg/m}^2$)	0.24 (0.04 -1.33)	1.79 (0.7 -4.3)	0.34 (0.02-5.13)	0.94 (0.44-2.02)	0.40 (0.06-2.65)
Thiocyanate /350 $\mu\text{mol/l}$	0.48 (0.22 -1.05)	2.3 (1.6 -3.21)	0.0 (0.0-0.9)	0.72 (0.52 -0.99)	2.17 (1.07-4.43)
Konzo in households	2.06 (0.38 -11.1)	0.71 (0.3-1.6)	20.4 (1 - 416.15)	1.97 (0.87-4.47)	0.41 (0.06 – 2.8)

Legend: OR = Odd Ratio at multinomial logistic regression; n = number, BMI = Body Mass Index; MCI = Mild Cognitive Impairment. MNCD = Major Neurocognitive Disorder; HT = Hypertension.

Impairments in daily living activities increased as performance at the CSID decreased (Spearman $r = -.2$, $p < 0.01$), figure 3.




Figure 3. Impairments in daily living activities increased as performance at the

CSID decreased both in women and in men, and overall.

Discussion

The main purpose of this study was to assess cognitive status and daily life functioning of adults relying on cyanogenic cassava as the main source of food. One hundred thirty-six (33.5 %) and 13 subjects (3.2 %) had MCI and dementia, respectively. Patterns of neurocognition functioning showed several domains (verbal fluency, attention, memory, and/or visuo-spatial processing) affected at neurological screening. The very high proportion of MCI (33.5 %), with overt impact on functioning, raises serious concerns on economic productivity in konzo-affected areas (26).

We have previously conducted electrophysiological and neurodevelopmental studies that suggested neurocognitive deficits in konzo (10, 27). Preschool-children, school age children, and adolescent have neurodevelopmental disabilities associated with cassava cyanogenic exposure (10, 15-16). This study provides the first systematic attempt to determine whether cognition may be affected in

adults relying on cyanogenic cassava as the main source of food. The prevalence of dementia (3.2 %) is similar to those from studies across rural sub-Saharan Africa (28) suggesting no or little impact of the cyanogenic diet on the occurrence of dementia. The association between domain-specific cognition deficits and levels of urinary SCN is a significant finding, as it appears to be consistent of findings from other studies conducted in sub-Saharan Africa suggesting possible neurotoxicity links (2, 10, 15-16, 29-30). However, a robust study design with markers of direct cyanide load (e.g., blood cyanide) and metabolism of sulfur amino acids would provide clear insights on whether excretion of urinary SCN is either a sign of marked toxicity or a sign of protective mechanisms against cyanide toxicity. The high proportion of depressed mood and/or the impact of hypertension on cognition as revealed in our study population, despite a small number of subjects with the condition, warrant further investigations. This present study remains, however, an important step in revealing the extent to which cognition and daily life



functioning may be affected among heavy cassava reliant populations; providing insights into the existence of preventable causes of cognition deficits among these populations.

Global Health Importance and Perspectives

Neurocognitive deficits in adults are common among Congolese adults who depend on cyanogenic cassava as their primary food source. Much of sub-Saharan Africa is becoming increasingly dependent on high-yield varieties of cassava that thrive in ecological degraded zones. Cassava is increasingly used as food-thickening ingredient. It is also exported to feed animals and produce snack for human consumption. Therefore, it is critical to continue brain/behavior research that will help us better understand its neurotoxic properties. Longitudinal studies, combined with studies of more sensitive biomarkers of neuropathogenesis, will help determine whether neurocognitive deficit is on a cassava-associated continuum of neurodegeneration and whether neurodevelopmental deficits in younger populations extend into adulthood.

Strengths and weaknesses of the study

This study is the first demonstration, in adults from an environment affected by konzo, of an obvious link between putative neurocognitive deficits and cassava cyanogenic exposure. Causal relationships may not be established due to lack of reliable markers of blood cyanide load and limits inherent to cross-sectional designs. Longitudinal studies will help determine whether MCI is on a continuum of a cassava-associated neurodegeneration and/or whether neurodevelopmental deficits in younger populations extend into adulthood.

Conflict of interest statement

Authors have no conflicts of interest to report.

Author contributions

Victor Hutu, MD, University of Kinshasa, participated in the field testing and drafting of the manuscript.

Daniel Okitundu, MD PhD, University of Kinshasa, participated in the field testing, analysis of data, and drafting of the manuscript.

Michael JBoivin, PhD MPH, Michigan State University, participated in the design of the study, analysis of the data, and drafting of manuscript.

Bruno Giordani, PhD, University of Michigan, participated in data analysis and drafting of manuscript.

Dieudonné Mumba MD PhD, Institut National de Recherches Biomédicales, participated in the study design, biochemical analyses, and drafting of manuscript.

Desire Tshala-Katumbay, MD PhD, Oregon Health and Sciences University, was the study principal investigator, participated in the design of study, oversight of neurological exam and data analysis, and drafting of the manuscript.



Acknowledgements

We thank the community of Kahemba in the Kwango (previously known as Bandundu) Province of DR Congo for their participation in the current research project.

Funding Statement

This study was funded by a grant from the National Institute of Environmental Health and Science: R01ES019841 (PI: Tshala-Katumbay). The study sponsor had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript. IRB approval for this study was obtained from the Ministry of Health from the Government of the Democratic Republic of the Congo.



Table 1. Clinical and biological characteristics of study by gender and cognition status

Legend: % = percentage; N/n = Number; SD= Standard Deviation. N = Number, SBP = Systolic blood Pressure. DBP=Diastolic blood Pressure.

Variables	All (N=406)	Men (N=203)	Women (N=203)	CSID ≤ 25 (Impaired N=149)	CSID > 25 (Normal N=257)	MCI (N=136)	Dementia (N=13)
Mean (SD) age	38.37 (11.12)	39.33 (10.96)	37.41 (11.21)	38.35 (9.2)	38.4 (13.9)	39 (14.4)	32.1 (2.6)
Men (n, %)	203 (50)	203 (50)	-	75(18,5)	128(31,5)	69(17)	6(1,5)
Women (n, %)	203 (50)	-	203(50)	74(18,2)	129(31,8)	67(16, 6)	7(1,7)
post-primary education level	344 (85)	180 (44.3)	164 (40.4)	126 (31.0)	218 (53.7)	115 (28.3)	11 (2.7)
Mean (SD)BMI (kg/m ²)	18.83 (3.1)	18.7 (2.9)	19 (3.3)	18.7 (3.0)	18.9 (3.0)	18.7 (3.0)	18.6 (2.0)
Undernutrition (BMI ≤ 18.4 kg/m ²)	220 (54, 2)	115(28,3)	105(25,9)	82(20,2)	138(34)	75(18,5)	7(1,7)
Mean (SD) SBP (mmHg)	118 (20.5)	117.8 (20.3)	118.25 (20.5)	115 (16.3)	119.8 (21.5)	114.6 (15.9)	118.9 (21.0)
Mean (SD) DBP (mmHg)	71.5 (13.1)	71.4 (13.2)	72 (13)	68.4 (11)	73.3 (13.8)	68 .1 (11)	71.5 (13.4)
Hypertension (n, %)	19(4, 7)	10 (2,5)	9(2, 2)	*14(3, 4)	*5(1,23)	*14(3, 4)	*0(0)
Mean (SD) Glycemia (mg %)	89.78 (35)	88.3 (33.2)	91.3 (36.8)	89.19 (34)	90.2 (34.6)	90.2 (37)	76 .7 (15)
Hyperglycemia (n, %)	55(13, 5)	26(6,4)	29(7, 1)	20(4, 9)	35(8, 6)	19 (4.7)	1(0,2)
Smoking (n, %)	24(5, 9)	13(3, 2)	11(2,7)	13(3,2)	11(2,7)	10 (0.5)*	3 (0.7)*
Alcohol consumption (n, %)	267(65,8)	135(33,3)	132(32,5)	98(24,1)	169(41,6)	89 (21.9)	9 (2.2)
Konzo subject (n, %)	38(9,4)	26(6,4)	12(2,9)	16(3,9)	22(5,4)	15 (3.7)	1(0,2)
Palmomental reflex (n, %)	349(86)	172 (42,4)	177(43,6)	*135(33,3)	*214(52,7)	124 (30.5)	11(2,7)
Snout reflex (n, %)	205 (50,5)	110 (27,1)	95(23,4)	77(19)	128(31,5)	69 (17)	8 (2)
Mean (SD) USCN (μmol/l) (N=242)	949.6 (518.3)	961.9 (534)	935.25 (501.5)	€1049(479.7)	€889.62 (532.8)	€1050.4 (479.1)	€1032(525.5)
USCN 100 – 350 (n, %)	54(22,3)	32(13,2)	22(9,1)	17(7)	37(15,3)	17(7)	0
USCN > 350 (n, %)	188(77,7)	98(40,5)	90(37,2)	71(17, 5)	117(48,3)	66(27.3)	5(2,1)
Mean (SD) CSID Score	25.8 (2.3)	25.8 (2.4)	25.8 (2.2)	23.2 (0.9)	27.4 (1.2)	23.2 (1.0)	23 (0.6)
Mean (SD) Golberg Score	29.6 (14.0)	29.37 (13.3)	29.8 (13.9)	15.6 (3.8)	37.6 (10.6)	16(3.4)	15(3.2)
Depressed Mood (n, %)	258(63,5)	129(31,8)	129(31,8)	1(0,2)	257(63,3)	1(0,2)	0(0)
Mean (SD) informant score	16.8 (8.7)	19.37 (8.8)	14.18 (7.8)	21.5 (8.4)	14.1 (7.7)	21.1 (8.6)	23(0.6)
MCI (n, %)	136(33.5)	69(17)	67(16, 6)	136(33.5)	-	136(33.5)	-
Dementia (n, %)	13(3,2)	6(1,5)	7(1,7)	13(3,2)	-	0	13(3,2)

USCN=Urinary concentration of Thiocyanate.MCI = Mild Cognitive Impairment. MNCD = Major Neurocognitive Disorder. * = significant difference in khi-square test. € = significant difference in Student's test-t. BMI=Body Mass Index, Goldberg Score ≥ 22 = Depressed Mood, CSID Score < 25 = Cognition Deficit.

Ann. Afr. Med., vol. 16, n° 2, Mars 2023

This is an open article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licences/bync/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited



References

1. Osuntokun BO. Cassava diet, chronic cyanide intoxication and neuropathy in the Nigerian Africans. *World Rev Nutr Diet.* 1981; **36**:141–173.
2. Olowule O, Onabolu A, Link H, Rosling H. Persistence of tropical ataxic neuropathy in a Nigerian community. *J Neurol Neurosurg Psychiatry.* 2000; **69** (1): 96-101.
3. Adamolekun B. Neurological disorders associated with cassava diet: a review of putative etiological mechanisms. *Metab Brain Dis.* 2011 Mar; **26** (1):79-85.
4. Mlingi N.L., Z. Bainbridge Z. Reduction of cyanogen levels during sun-drying of cassava in Tanzania. *Acta Hort.* 1994. **375**:233-239.
5. Ciglenecki I., Eyema R., Kabanda C., Taafo F., Mekaoui H., Urbaniak V.. Konzo outbreak among refugees from Central African Republic in Eastern region, Cameroon. *Food Chem Toxicol* 2011; **49** (3):579-582.
6. Siddiqi OK, Kapina M, Kumar R, Ngomah MA, Kabwe P, Mazaba ML, *et al.* Konzo outbreak in the Western Province of Zambia. *Neurology.*2020; **94** (14):e1495-e1501.
7. ACF International. Étude préliminaire du projet Amélioration de la diète et éradication de l'intoxication alimentaire appelée Konzo dans le Kwango. Disponible sur <http://rdchumanitaire.net/index.php/documentation>, Mai 2015. Accédé le 10 Octobre 2016.
8. Okitundu LD, Imbula EB, N'situ MA, Ndjukendi OA, Kamanga MT. Affects de la mère pendant la grossesse, relation mère-bébé, santé et développement du nourrisson à Kinshasa. *Pan African Medical Journal.* 2020;**36**: 203.
9. Banea JP, Bradbury JH, Mandombi C, Nahimana D, Denton IC, Kuwa N, *et al.* Control of konzo by detoxification of cassava flour in three villages in the Democratic Republic of Congo. *Food Chem. Toxicol.*2013; **60**:506-513.
10. Boivin MJ, Okitundu D, Makila-Mabe BG, Sombo MT, Mumba D, Tylleskar T, *et al.* Neuropsychological effects of konzo: a neuromotor disease associated with poorly processed cassava. *Pediatrics* 2013 ; **131** (4) : e1231-e1239.
11. Jorgensen K., Morant A V., Morant M., Jensen N.B., Olsen CE., Kannangara R., *et al.* Biosynthesis of the cyanogenic glucosides linamarin and lotaustralin in cassava: isolation, biochemical characterization, and expression pattern of CYP71E7, the oxime-metabolizing cytochrome P450 enzyme. *Plant physiology* 2011; **155** (1):282–292.
12. Zuk M, Pelc K, Szperlik J, Sawula A, Szopa J. Metabolism of the Cyanogenic Glucosides in Developing Flax: Metabolic Analysis, and Expression Pattern of Genes. *Metabolites.* 2020 Jul 14; **10** (7):288.
13. Cliff J., Muquingue H, Nhassico D, Nzwalo H. , Bradbury HJ.. Konzo and continuing cyanide intoxication from cassava in Mozambique. *Food and Chemical Toxicology.* 2011; **49** (3):631–635.
14. Diasolua ND. “Konzo, Cassava Toxicity and Associated Nutritional Factors. *CCDN NEWS* 2005. **6**:1–2.
15. Okitundu LD, Sombo MT, Bumoko MG, Banea MJ, Mumba ND, Boivin MJ, *et al.* Troubles socio-émotionnels de l'enfant en milieu Konzo, un syndrome paralytique de nature épidémique associé à une intoxication cyanhydrique d'origine alimentaire en Afrique sub-saharienne. *Pan African Medical Journal.* 2018; **31**:118.
16. Kashala-Abotnes E, Sombo MT, Okitundu LD, Kunyu M, Bumoko MG, Tylleskar T, *et al.* Dietary cyanogen exposure and early child neurodevelopment: An observational study from the Democratic Republic of Congo. *PloS one.* 2018;**13** (4):e0193261.



17. Banea JP, Bradbury JH., Nahimana D., Denton IC., Mashukano N., Kuwa, N. Survey of the konzo prevalence of village people and their nutrition in Kwilu District, Bandundu Province, DRC. *African Journal of Food Science* 2015; **9** (2): 43-50.
18. Hall KS, Gao S, Emsley CL, Ogunniyi AO, Morgan O, Hendrie HC. Community screening interview for dementia (CSID'); performance in five disparate study sites. *Int J Geriatr Psychiatry*. 2000. **15** (6): 521–531.
19. Fatai KS, Sunday AB, Musa AW, Bukar B, Bukbuk DN, Kida I. Cognitive function in HIV-seropositive Nigerians without AIDS, *Journal of the Neurological Sciences* 2008;**267**:142-146.
20. Arinzechi EO, Ogunrin OA, Nwosu CM, Nwani PO, Enwereji KO, Asomugha LA, *et al.* A community-based case-control study of prevalence and pattern of cognitive impairments in patients with epilepsy residing in South Eastern Nigeria. *J Neurosci Rural Pract* 2016; **7**:405-411.
21. Imbula EB, Okitundu LD, Mampunza MS. Prévalence et facteurs de risque de la dépression maternelle postnatale à Kinshasa. *Médecine et Santé Tropicales*. 2012; **22** (4):379-384.
22. OMS. Konzo, a distinct type of upper motoneuron disease. *Weekly Epidemiological records* 2015; **30** (71):225-232.
23. FAO/WHO Expert Committee on Food Additives (83rd, 2017: Geneva, Switzerland). (2017). Evaluation of certain contaminants in food: eighty-third report of the Joint FAO/WHO Expert Committee on Food Additives. WHO Technical Report Series; 1002 World Health Organization. Disponible sur <https://apps.who.int/iris/handle/10665/254893>
24. Banea JP., Bradbury JH., Nahimana D., Denton IC., Foster MP., Mekob N, *et al.* Health factors associated with persistent konzo in four villages in the Democratic Republic of Congo (DRC). *African Journal of Food Science*. 2016; **10** (1):1-6.
25. Bradbury MG, Egan SV, Bradbury HJ. Picrate paper kits for determination of total cyanogens in cassava roots and all forms of cyanogens in cassava products. *J Sci Food Agric*. 1999; **79**: 593-601.
26. Tshala-Katumbay D, Kazadi K, Tylleskar T, Eeg-Olofsson KE. Impairments, disabilities and handicap pattern in konzo – a non-progressive spastic para/tetraparesis. *Disability and Rehabilitation* 2001; **23** (16): 731-736.
27. Tshala Katumbay D, Lukusa VM, Eeg-Olofsson KE. EEG findings in Konzo: a spastic para/tetraparesis of acute onset. *Clin Electroencephalogr*. 2000 Oct; **31** (4):196-200.
28. Mbelesso, P., Tabo, A., Guerchet, M., Mouanga AM, Bandzouzi B, Houinato D *et al.* Épidémiologie des démences chez les personnes âgées dans le troisième arrondissement de la ville de Bangui (République Centrafricaine). *Bull. Soc. Pathol. Exot*. 2012; **105**:388–395.
29. Villanueva I, Alva-Sanchez C, Pacheco-Rosado J. The role of thyroid hormones as inductors of oxidative stress and neurodegeneration. *Oxid Med Cell Longev*. 2013; **2013**: 218145.
30. Makila-Mabe BG, Kikandau KJ, Sombo TM, Okitundu DL, Mwanza JC, Boivin MJ, *et al.* Serum 8, 12-Isoprostane F2-VI marker of oxidative damage and cognition deficits in children with konzo. *PlosOne* 2014; **9** (9):e107191.

Cite this paper as: Kabamba VH, Luwa DO, Banea JPM, Giordani B, Boivin MJ, Mumba-Ngoyi D, *et al.* Screening for neurocognitive deficits in adult populations reliant on toxic cassava as the main source of food. *Ann Afr Med* 2023; **16** (2): e5028-e5038. <https://dx.doi.org/10.4314/aamed.v16i2.2>