

Pure Endoscopic Excision of Parenchymal Brain Tumors: Feasibility, Risks, Advantages and Realities - A Beginners Perspective

Manish Budhiraja¹ Ashis Pathak¹ Harmandeep Brar¹ Rahat Brar²

¹ Department of Neurosurgery, Fortis Hospital, Mohali, India

² Department of Radiodiagnosis, Fortis Hospital, Mohali, India

Address for correspondence Ashis Pathak, MS, MCh., MNAMS, FICS.,
Department of Neurosurgery, Fortis Hospital, Mohali, India
(e-mail: ashispathak2006@rediffmail.com).

Arq Bras Neurocir 2020;39(3):201–206.

Abstract

Background Neuroendoscopy is gaining popularity and is reaching new realms. Young neurosurgeons are exploring the various possibilities associated with the use of neuroendoscopy. Neuroendoscopy in excision of parenchymal brain tumors is less explored, and young neurosurgeons should be aware of the realities. The present article is an approach to put forward the difficulties faced by a young neurosurgeon and the lessons learnt.

Objective To report the experience of surgical excision of parenchymal brain tumors, in selected cases, using pure endoscopic approach and to discuss its feasibility, technical benefits, risks and comparison with conventional microscopic excision.

Method Eight patients of variable age group with parenchymal brain tumors were operated upon by a single surgeon and followed up for a period varying from 6 months to 2 years. Data regarding operating time, illumination, clarity of the field, size of craniotomy, blood loss and course of recovery was evaluated. All of the tumors were resected using rigid high definition zero and 30° endoscope.

Results Out of eight cases, seven had lesions in the supratentorial and one in the infratentorial location. The age group ranged from 27 to 74 years old. Near to gross total resection was achieved in all except two cases. All of the patients recovered well without any significant morbidity or mortality. Hospital stay was reduced by 1 day on average.

Conclusion Excision of parenchymal brain tumors via pure endoscopic method is a safe and efficient procedure. Although there is an initial period of learning curve, it is not steep for those already practicing neuroendoscopy, but the approach has its advantages.

Keywords

- clarity
- illumination
- learning curve
- magnification
- neuroendoscopy

Introduction

Neuroendoscopic excision of intraventricular and sellar tumors is a well-established neurosurgical technique, but the use of this approach for parenchymal tumors has not gained popularity. Several surgeons have tried combined open and endoscopic approaches with success.^{1–4} However, pure endoscopic resection of parenchymal brain tumors is

not routinely in vogue, the possible reasons being unfamiliarity with the technique and apprehension of incomplete resection at the depth. Although the open microscopic excision procedure is perceived to be better due to the availability of high end microscopes, advanced magnetic resonance imaging (MRI) sequences and support of neuronavigation, the latest neuroendoscopes offer better clarity of vision.⁵ Contrarily, the endoscopic approach for parenchymal brain

received
November 30, 2019
accepted
December 23, 2019

DOI <https://doi.org/10.1055/s-0040-1702975>.
ISSN 0103-5355.

Copyright © 2020 by Thieme Revinter
Publicações Ltda, Rio de Janeiro, Brazil

License terms



tumor excision is a safer alternative as it allows easy entry to the tumor through the parafascicular corridor of white matter, offers better clarity of image because of the proximity of the endoscope to the brain and tumor surface.⁵ Also, the smaller size of craniotomy and the narrow corridor of entry is sufficient to allow bimanual manipulation of the tumor with clear visibility of the hidden corners. The technique therefore is minimally invasive and strictly adheres to proven microsurgical principles.^{6–8} Eight patients with parenchymal brain tumors were operated upon using the pure endoscopic method. Comparison on various parameters was done with the purely microscopic approach. The feasibility of this approach along with associated risk, benefits and misconceptions are discussed.

Summary

A total of 8 patients (► **Table 1**), with age ranging from 27 to 74 years old, with a follow-up period of 2 months to 2 years are presented. These include three females and five male patients, out of which seven had supratentorial and one had infratentorial location of the lesions. We were able to achieve near to gross total resection in all the lesions except in two, as one lesion had close proximity to the vessels and there was misleading frozen section report in the other. Three lesions were low grade, four were high grade and one was a hemangioblastoma. Only one patient had significant deficit in the postoperative period in the form of opposite side weakness with aphasia, which recovered with time. None of the remaining patients developed any remarkable postoperative deficits.

Operative Technique

All of the patients underwent contrast-enhanced MRI under neuronavigation protocol. Following standard neuroanaesthetic technique for induction, the heads of the patients were positioned and fixed with the help of a Sugita head frame (Head support of OT table company name, Mizuho, Made in Japan, Made in Japan). Using stealth navigation system, the

position of the tumor was confirmed and a limited craniotomy was planned centering on the main bulk of the lesion. Although the initial two cases had a slightly larger craniotomy as a precautionary learning curve measure, the subsequent exposures were reduced to almost half the size of standard exposures. Dural openings were similar to cruciate exposure. The site of cortisectomy was decided based on neuronavigation guidance depending upon the shortest route of entry and eloquence of structure(s) in the vicinity. Cortisectomies were tailored to the optimal space requirement for the introduction of the endoscope and maneuverability of the operating instruments. The initial debulking was performed using a zero degree scope mounted on an endoscope holder which allowed the freedom for bimanual excision. Subsequently, a 30° scope was introduced for visualization of the surrounding lesion in blind corners of the cavity using minimal retraction. There was no need for the use of other angled scopes. Tumor excision followed the same principles of cautery and suction (CUSA) evacuation using navigation guidance. In the initial 2 cases, the microscope was brought in at the end of surgery to confirm the definition of the tumor-brain interface, which reaffirmed good tumor clearance. In initial cortisectomies, the margins were not supported leading to subpial hemorrhages and cortical changes. Subsequent use of gloves strips circumvented the issue. Hemostasis was achieved in the usual manner followed by complete dural closure in every case.

Results

A total of eight patients of intraparenchymal lesions were operated in the present series, out of which seven were supratentorial and one was in infratentorial location. The age group ranged between 27 and 74 years old. Near to gross total resection was achievable in all of the cases, except in two (► **Fig. 1, 2 and 3**). One patient with a dominant frontotemporal lobe tumor developed hemiparesis and motor aphasia in the postoperative period, which recovered gradually. The remaining patients had no neurological deficit (► **Table 2**). Considering

Table 1 Details of the patients including types of tumors and postoperative complications

Case No	Age/Sex	Location	Resection	Histopathology	Complication
1.	27 yrs/F	Left temporal (► Fig. 1)	Near total	Diffuse infiltrating astrocytoma Grade II	Nil
2.	34 yrs/F	Left posterior frontal (► Fig. 2)	Gross total	Oligodendroglioma Grade II	Right hemiparesis with aphasia
3.	57 yrs/M	Left cerebellar	Gross total	Hemangioblastoma Grade I	Nil
4.	74 yrs/M	Right parieto occipital (► Fig. 3)	Gross total	Glioblastoma multiforme Grade IV	Nil
5.	36 yrs/F	Left perisylvian	Gross total	Astrocytoma grade III	Nil
6.	64 yrs/M	Right parieto occipital	Subtotal	Glioblastoma Multiforme Grade IV	Nil
7.	33 yrs/M	Right Frontal	Near total	Oligodendroglioma grade II	Nil
8.	59 yrs/M	Left Fronto-Parietal	Gross total	Glioblastoma Multiforme Grade IV	Nil

Abbreviations: F, female; M, male.




Fig. 1 (A) Axial T2 weighted magnetic resonance imaging showing the heterogeneous hyperintense lesion in the anterior temporal lobe showing ill-defined margins. (B) Axial computed tomography image showing postoperative changes in the left temporal lobe and foci of air in the operative bed. (C) Intraoperative endoscopic view during tumor excision showing partly fibrous tumor.




Fig. 2 (A) T2 weighted sagittal image of the brain showing lesion in the left frontal lobe. (B) Sagittal computed tomography image showing postoperative changes in the left frontal lobe with small foci of air at the postoperative site and extra axial nondependent air in the frontal region. (C) Intraoperative endoscopic image showing tumor tissue. (D) Endoscopic view after tumor excision showing the surgical lining at the margins.

the evaluation of the endoscopic technique with microscopic procedure, the parameters compared were: operating time, illumination, clarity, magnification, blood loss, size of craniotomy, postoperative imaging. The surgical time was slightly




Fig. 3 (A and B) Preoperative axial and sagittal T1 contrast magnetic resonance imaging showing large cystic tumor with enhancing nodule in the left cerebellum. (C and D) Postoperative axial and sagittal plain computed tomography images showing complete excision with no tumor bed hematoma.

longer than the microsurgical technique, most likely due to our learning curve, but blood loss was comparable to that of the microscopic technique. The illumination, clarity and magnification were at par with the microscope, but the blind corners were much better visualized with good clarity (►Table 2; ►Fig. 1, 2). We saw a significant reduction in craniotomy size (►Fig. 3). The extent of excision was >95% in all cases except one; although the number of patients was not adequate to reach any conclusion, adequate visualization of blind corners definitely helped in excising the tumors with a good limit of confidence and safety with minimal damage to the normal tissue due to small cortisectomies. The patients recovered well and were discharged home on average 1 day earlier than those who were submitted to the routine microscopic approach with large craniotomy (►Table 2 and 3). A concise comparison between the standard microscopic approach and the pure endoscopic approach, with emphasis on selected important points, is shown in ►Table 4.

Discussion

Pure endoscopic resection of intraparenchymal brain tumors is a minimally invasive approach that is not routinely practiced by neurosurgeons. To the best of our knowledge, until now, there are two major series with 21 and 48 cases, respectively, by Kasam et al¹ and Plaha et al² with few other sporadic reports (►Table 5). The reason for the lower popularity of this technique is due to the unfamiliarity with the procedure, the long learning curve and apprehension about inadequate exposure and inadequate visibility through the endoscope.

Table 2 Evaluation of Parameters in pure endoscopic method

S No	Parameters	Case I (► Fig. 1)	Case II (► Fig. 2)	Case III	Case IV (► Fig. 3)	Case V	Case VI	Case VII	Case VIII
1	Surgical Time	2 hrs 15 mins	3 hrs 25 mins	2 hr 45 mins	2 hr 50 mins	2 hrs 30 mins	2 hrs 10 mins	1 hr 50 mins	2 hrs 10 mins
2	Illumination & Clarity	Very good specially at corners	Very good specially at corners	Very good specially at corners	Very good specially at corners	Very good specially at corners	Very good specially at corners	Very good specially at corners	Very good specially at corners
3	Magnification	Tumor brain interface well differentiated	Tumor brain interface well differentiated	Tumor brain interface well differentiated	At par with microscope with little difficulty in hemostasis at depth	At par with microscope with little difficulty in hemostasis at depth	Tumor brain interface well differentiated	At par with microscope with little difficulty in hemostasis at depth	Tumor brain interface well differentiated
4	Blood Loss	150–200 ml	200 ml	200 ml	250–300 ml	400 ml	150–200 ml	100–150 ml	150–180 ml
5	Craniotomy Size	70% of microscopic	Standard	50% of microscopic	40% of microscopic	60–70% of microscopic	Standard (extensive edema)	60% of microscopic	60–70% of microscopic
6	Tumor bed hematoma	No	Small	No	Significant but without mass effect	Significant but no mass effect	Small	Nil	Small

Table 3 Evaluation of parameters in randomly selected cases of pure microscopic excision of parenchymal tumors

S No	Parameters	Case I	Case II	Case III	Case IV	Case V	Case VI	Case VII	Case VIII
1	Surgical Time	1 hrs 50 mins	1 hrs 45 mins	2 hr 30 mins	2 hr 15 mins	2 hrs	2 hrs 10 mins	1 hr 50 mins	2 hrs 25 mins
2	Illumination & Clarity	Poor at corners no visualization of undermining edges	Poor at corners no visualization of undermining edges	Poor at corners no visualization of undermining edges	Poor at corners no visualization of undermining edges	Poor at corners no visualization of undermining edges	Poor at corners no visualization of undermining edges	Poor at corners no visualization of undermining edges	Poor at corners no visualization of undermining edges
3	Magnification	Good	Good	Good	Good	Good	Good	Good	Good
4	Blood Loss	100–120 ml	100–120 ml	100–150 ml	300 ml	100 ml	150–200 ml	100–120 ml	150 ml
5	Craniotomy Size	Large	Decompressive	Standard according to tumor size	Standard according to tumor size	Standard according to tumor size	Standard according to tumor size	Standard according to tumor size	Standard according to tumor size
6	Tumor bed hematoma	No	No	Small	No	Small	No	No	No

Table 4 Comparison of pure endoscopic with pure microscopic excision in parenchymal brain tumors

S No	Parameters	Pure Microscopic	Pure Endoscopic
1.	Surgical Time	Less time consuming	Comparatively more due to learning curve initially
2.	Illumination and clarity	Poor at Blind Corners	Neatly visualized and especially undermined edges
3.	Magnification	Good with 3D vision	At par with microscope but with comparatively difficult depth perception
4.	Blood Loss	Better controlled	Sometimes difficult to control due to difficult depth perception Microscope can aid at the end of surgery in case of difficulty
5.	Craniotomy Size	Large craniotomy	Mini craniotomy size reduction by minimum of 20–30%
6.	Tumor bed hematoma	Lower incidence	Incidence may be higher due to learning curve initially.

Table 5 Summary of major studies of pure endoscopic approach for excision of parenchymal brain tumors

Author	Method	Type of endoscope	Extent of resection	Limitations
Kassam et al, 2009 ¹	Neuroendoport Conduit 11.5 mm	0-degree endoscope	Total resection 38%, near total 28.6%, sub-total 33.3% multiple	Manipulations of conduit required to achieve maximal resection. Conduit cannot be used for tumors reaching pial surface
Jo et al, 2011 ⁴	Transparent tubular conduit 11 mm	0-degree endoscope	Gross total resection in all cases	Small lesions (3 cm) limited
Otsuki et al., 1990 ³	Tubular conduit on stereotactic frame	0-degree endoscope	Total resection in 8 lesions; biopsy or aspiration in 7	Small lesions limited by size of conduit
Plaha et al, 2014 ²	Nontubular access corridor 10 mm	30-degree endoscope	Total resection 48%, > 95% resection 70%	Needs further development of microsurgical instruments and access corridor
Present study	Nontubular access corridor 10 - 15 mm	0 and 30-degree endoscope	Near total to gross total resection	Limited number of patients & variability in types of lesions

Although it is perceived that endoscopic brain tumor resection requires a long learning curve, neurosurgeons familiar with endoscopic transphenoidal procedures would already have a hang of working in a narrow, rigid corridor within the bounds of limited space. Selected entry through cortisectomy using a safe surgical corridor under neuro-navigation guidance gives an easy access to the lesions. However, the maneuvering of the endoscope and the instruments need to be more gentle because of the risk of retraction injury to the normal parenchyma around the port of entry. Although the use of rigid conduits has been claimed to be less harmful as compared with normal brain retractors,⁹ we agree with the proposed argument that a constant pressure on the brain would add to ischemic insults apart from compression effect on the surrounding normal parenchyma.^{2,10} According to our experience, the use of a pulsatile retractor combined with latex glove lining provides the advantage of minimizing traction effect and allowing the desired exposure needed for resection. As a beginner, it is suggested to perform the initial cases with slightly larger exposure so as to have a leeway to revert to microscopic excision if the surgeon is inconvenienced at any stage.

Comparing the illumination and visibility between microscopic versus endoscopic excision, undoubtedly the microscope gives better resolution. However, it is also true that for deeper entry corridors there is considerable loss of light leading to poor visibility and resolution in microscopic techniques in deeper areas. This problem is compounded by manipulation of instruments at the depth. Contrarily, the endoscope has the advantage of better optical resolution as the light source is nearer to the target. This enhances the visibility and adds advantage to better differentiation of the brain-tumor interface without compromising the magnification of the image.

Resection of a lesion without fresh neurological deficit remains a major challenge in all intraparenchymal lesions. The microscopic vision is limited by straight projection of light rays at the depth of the surgical field, which is different from endoscopic visualization as it provides a wide-angled panoramic view, giving better clarity at the depth. Moreover, the use of an angled endoscope can help in viewing the blind corners without much manipulation or retraction.

Our Learning Experience

Usually a single surgeon will suffice in the microscopic approach, but in the endoscopic approach there may be the need of an assistant to hold the scope and navigate. In the present series, the need for two persons was obviated by mounting the endoscope on a holder which gave an unhindered opportunity for bimanual excision by a single surgeon. However, it may take a while to switch over and get oriented from the resolution of a microscope to endoscope. As a beginner, it is always safe to begin with a bigger exposure and have a fall back option to bring in the microscope, if the situation demands. As it happened in one of our cases, it was decided to use the microscope for a brief period since there was difficulty in manipulating between the vessels of the sylvian fissure in a peri-sylvian lesion (case no 5, ► **Table 1**).

Hence, there should not be any dogma to use technology interchangeably in situations on demand.

Cortisectomy length of 1.5 to 2 cm, which was initially supported by cotton patties, led to widening of the cortisectomy margins and damage to the edges. Subsequently, it was planned to insulate the edges with surgical superimposed with glove patties with overlying cotton patties to hold them in place. Using this technique, the damage to the surrounding brain and to the extension of the cortisectomy margins was reduced to minimum, thus achieving near total resection in most of the cases avoiding unwanted cortical injury.

The resection of a lesion is facilitated by the initial use of a zero degree scope. Subsequent to reasonable debulking/excision of the tumor, the use of a 30° scope with minimal sector wise retraction of the cortisectomy margins helps in achieving a total excision of the lesion, even at blind corners, through good resolution and visibility. The use of more obtuse angled scopes, in our experience, is not necessary.

Outcome and Complications

The level of resection achieved in previous studies varies from 29% to > 95%.¹⁻⁴ We achieved near to gross total resection in all cases and the illumination, magnification and clarity was excellent. However, this is a small series to comment upon the exact extent of the excision of the lesions in a wide variety of cases. The blood loss in all cases was comparable to that of microscopic excision, with no postoperative tumor bed hematoma in any case. One patient with oligodendroglioma (case no 2, ► **Table 1**) situated in a strategic location in the dominant hemisphere, developed aphasia and right side weakness which improved gradually over a period of time. Although there was the issue of depth perception initially, that was overcome with subsequent experience. It is most important that the surgeon conducts a critical evaluation of clinical and radiological findings before embarking upon endoscopic excision, and also weighs the plausibility of conversion to microscopic excision to avoid any adventurous complications. The issue of falling brain and managing brisk bleeding in a vascular tumor bed may at times blind the vision through a scope. Hence a steady and slow resection of vascular lesions in a controlled way is likely to achieve the goal of satisfactory resection. As mentioned earlier, there should not be any hesitation to resort to the help of a microscope in situations of brisk bleeding. The resection of surface tumors, in our experience, turned out to be an easier option with hardly any need for retraction. This technique was demonstrated to be accurate and safe, and possibly will be expanded to remove other intraparenchymal lesions in the future.¹¹

Conclusion

In spite of its innovative and beneficial aspects, neuroendoscopy, like any other diagnostic and treatment technique, has some risks. The most significant is perhaps the risk of local injury to the surrounding structures and normal brain. Other risks of neuroendoscopy include hemorrhage (with an associated difficulty in hemostasis) leading to raised intracranial

pressure. However, this procedure has high future potentials to establish itself as a minimally invasive technique, although it still remains in its nascent phase. With ongoing development of endoscopic instruments and advanced surgical techniques including multiport approaches, endoscopic surgery will be expanded beyond intraventricular and skull base lesions to intraparenchymal brain lesions. These advances will be important for the future of endoscope-assisted microsurgery. In the future, neuroendoscopy is expected to become routine in modern neurosurgical practice. Institutions should develop training programs for young neurosurgeons.¹²

Conflict of Interests

The authors have no conflict of interests to declare.

References

- 1 Kassam AB, Engh JA, Mintz AH, Prevedello DM. Completely endoscopic resection of intraparenchymal brain tumors. *J Neurosurg* 2009;110(01):116–123
- 2 Plaha P, Livermore LJ, Voets N, Pereira E, Cudlip S. Minimally invasive endoscopic resection of intraparenchymal brain tumors. *World Neurosurg* 2014;82(06):1198–1208
- 3 Otsuki T, Jokura H, Yoshimoto T. Stereotactic guiding tube for open-system endoscopy: a new approach for the stereotactic endoscopic resection of intra-axial brain tumors. *Neurosurgery* 1990;27(02):326–330
- 4 Jo KW, Shin HJ, Nam DH, et al. Efficacy of endoport-guided endoscopic resection for deep-seated brain lesions. *Neurosurg Rev* 2011;34(04):457–463
- 5 Teo C, Mobbs R. Neuroendoscopy. In: Setti S, Rengachary, Richard G. Ellenbogen, eds. *Principals of Neurosurgery*. 2nd ed. London, UK: Elsevier Lifesciences; 2004: chapter 10
- 6 Kelly PJ, Goerss SJ, Kall BA. The stereotaxic retractor in computer-assisted stereotaxic microsurgery. Technical note. *J Neurosurg* 1988;69(02):301–306
- 7 Kelly PJ, Kall BA, Goerss S, Earnest F IV. Computer-assisted stereotaxic laser resection of intra-axial brain neoplasms. *J Neurosurg* 1986;64(03):427–439
- 8 Russell SM, Kelly PJ. Volumetric stereotaxy and the supratentorial occipitotemporal approach in the resection of posterior hippocampus and parahippocampal gyrus lesions. *Neurosurgery* 2002;50(05):978–988
- 9 Zhong J, Dujovny M, Perlin AR, Perez-Arjona E, Park HK, Diaz FG. Brain retraction injury. *Neurol Res* 2003;25(08):831–838
- 10 Yokoh A, Sugita K, Kobayashi S. Intermittent versus continuous brain retraction. An experimental study. *J Neurosurg* 1983;58(06):918–923
- 11 Sgouros S. *Neuroendoscopy: current status and future trends*. ed 1. Heidelberg: Springer Science & Business Media; 2013
- 12 Shim KW, Park EK, Kim D-S, Choi J-U. Neuroendoscopy: Current and Future Perspectives. *J Korean Neurosurg Soc* 2017;60(03):322–326