

ACR Appropriateness Criteria[®]

Asymptomatic Patient at Risk for Coronary Artery Disease: 2021 Update

Expert Panel on Cardiac Imaging: *Brian B. Ghoshhajra, MD, MBA^a, Sandeep S. Hedgire, MD^b, Lynne M. Hurwitz Kowek, MD^c, Garth M. Beache, MD^d, Richard K. J. Brown, MD^e, Andrew M. Davis, MD, MPH^f, Joe Y. Hsu, MD^g, Thomas V. Johnson, MD^h, Gregory A. Kicska, MD, PhDⁱ, Seth J. Kligerman, MD^j, Diana Litmanovich, MD^k, Christopher D. Maroules, MD^l, Nandini Meyersohn, MD^m, Mark G. Rabbat, MDⁿ, Todd C. Villines, MD^o, Samuel Wann, MD^p, Suhny Abbata, MD^q*

Abstract

Coronary atherosclerotic disease is a leading cause of mortality and morbidity due to major cardiovascular events in the United States and abroad. Risk stratification and early preventive measures can reduce major cardiovascular events given the long latent asymptomatic period. Imaging tests can detect subclinical coronary atherosclerosis and aid initiation of targeted preventative efforts based on patient risk. A summary of available imaging tests for low-, intermediate-, and high-risk asymptomatic patients is outlined in this document.

The American College of Radiology Appropriateness Criteria are evidence-based guidelines for specific clinical conditions that are reviewed annually by a multidisciplinary expert panel. The guideline development and revision include an extensive analysis of current medical literature from peer reviewed journals and the application of well-established methodologies (RAND/UCLA Appropriateness Method and Grading of Recommendations Assessment, Development, and Evaluation or GRADE) to rate the appropriateness of imaging and treatment procedures for specific clinical scenarios. In those instances where evidence is lacking or equivocal, expert opinion may supplement the available evidence to recommend imaging or treatment.

^aPanel Vice Chair, Massachusetts General Hospital, Boston, Massachusetts.

^bResearch Author, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, Assistant Clinical Director CV Division, Vascular Imaging.

^cPanel Chair, Duke University Medical Center, Durham, North Carolina, Director, Cardiovascular Imaging, Medical Director of CT; Duke University Medical Center.

^dUniversity of Louisville School of Medicine, Louisville, Kentucky.

^eUniversity of Utah, Department of Radiology and Imaging Sciences, Salt Lake City, Utah.

^fUniversity of Chicago Medical Center, Chicago, Illinois, American College of Physicians, Associate Vice-Chair for Quality, Department of Medicine, University of Chicago.

^gKaiser Permanente, Los Angeles, California.

^hSanger Heart and Vascular Institute, Charlotte, North Carolina, Cardiology Expert.

ⁱUniversity of Washington, Seattle, Washington.

^jUniversity of California San Diego, San Diego, California.

^kHarvard Medical School, Boston, Massachusetts.

^lNaval Medical Center Portsmouth, Portsmouth, Virginia. Section Chief, Cardiovascular Imaging, Naval Medical Center Portsmouth, Board of Directors, Society of Cardiovascular Computed Tomography.

^mMassachusetts General Hospital, Boston, Massachusetts.

ⁿLoyola University Medical Center, Maywood, Illinois, Society for Cardiovascular Magnetic Resonance.

^oUniversity of Virginia Health Center, Charlottesville, Virginia, Society of Cardiovascular Computed Tomography.

^pAscension Healthcare Wisconsin, Milwaukee, Wisconsin, Nuclear Cardiology Expert.

^qSpecialty Chair, UT Southwestern Medical Center, Dallas, Texas.

Corresponding author: Brian Ghoshhajra, MD, MBA, Cardiovascular Imaging, Department of Radiology, Massachusetts General Hospital, 55 Fruit Street, Boston, MA 02114; e-mail: bghoshhajra@mgh.harvard.edu.

The American College of Radiology seeks and encourages collaboration with other organizations on the development of the ACR Appropriateness Criteria through society representation on expert panels. Participation by representatives from collaborating societies on the expert panel does not necessarily imply individual or society endorsement of the final document. Reprint requests to: publications@acr.org.

The authors state that they have no conflict of interest related to the material discussed in this article. Dr Wann is self-employed. All other authors are non-partner/non-partnership track employees.

The ACR Appropriateness Criteria documents are updated regularly. Please go to the ACR website at www.acr.org/ac to confirm that you are accessing the most current content.

Disclaimer: The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Key Words: Appropriateness Criteria, Appropriate Use Criteria, Asymptomatic, AUC, Coronary atherosclerotic disease, CT coronary calcium score, CTA coronary arteries

J Am Coll Radiol 2021;18:S2-S12. Copyright © 2021 Published by Elsevier Inc. on behalf of American College of Radiology

ACR Appropriateness Criteria® Asymptomatic Patient at Risk for Coronary Artery Disease: 2021 Update. [Variants 1 to 3](#) and [Tables 1](#) and [2](#).

Variant 1. Asymptomatic patient. Low risk for coronary artery disease. Initial imaging.

Procedure	Appropriateness Category	Relative Radiation Level
CT coronary calcium	Usually Not Appropriate	☢☢☢
US echocardiography transthoracic resting	Usually Not Appropriate	○
CTA coronary arteries with IV contrast	Usually Not Appropriate	☢☢☢
MRA coronary arteries without and with IV contrast	Usually Not Appropriate	○
MRA coronary arteries without IV contrast	Usually Not Appropriate	○
MRI heart function and morphology without and with IV contrast	Usually Not Appropriate	○
MRI heart function and morphology without IV contrast	Usually Not Appropriate	○
MRI heart function with stress without and with IV contrast	Usually Not Appropriate	○
MRI heart function with stress without IV contrast	Usually Not Appropriate	○
Radiography chest	Usually Not Appropriate	☢
SPECT or SPECT/CT MPI rest and stress	Usually Not Appropriate	☢☢☢☢
US echocardiography transthoracic stress	Usually Not Appropriate	○

Variant 2. Asymptomatic patient. Intermediate risk for coronary artery disease. Initial imaging.

Procedure	Appropriateness Category	Relative Radiation Level
CT coronary calcium	Usually Appropriate	☢☢☢
CTA coronary arteries with IV contrast	May Be Appropriate	☢☢☢
MRA coronary arteries without and with IV contrast	Usually Not Appropriate	○
MRI heart function and morphology without and with IV contrast	Usually Not Appropriate	○
SPECT or SPECT/CT MPI rest and stress	Usually Not Appropriate	☢☢☢☢
US echocardiography transthoracic resting	Usually Not Appropriate	○
US echocardiography transthoracic stress	Usually Not Appropriate	○
MRA coronary arteries without IV contrast	Usually Not Appropriate	○
MRI heart function and morphology without IV contrast	Usually Not Appropriate	○
MRI heart function with stress without and with IV contrast	Usually Not Appropriate	○
MRI heart function with stress without IV contrast	Usually Not Appropriate	○
Radiography chest	Usually Not Appropriate	☢

Variant 3. Asymptomatic patient. High risk for coronary artery disease. Initial imaging.

Procedure	Appropriateness Category	Relative Radiation Level
CT coronary calcium	May Be Appropriate	☼☼☼
CTA coronary arteries with IV contrast	May Be Appropriate	☼☼☼
US echocardiography transthoracic resting	Usually Not Appropriate	○
MRA coronary arteries without and with IV contrast	Usually Not Appropriate	○
MRI heart function and morphology without and with IV contrast	Usually Not Appropriate	○
MRI heart function and morphology without IV contrast	Usually Not Appropriate	○
MRI heart function with stress without and with IV contrast	Usually Not Appropriate	○
MRI heart function with stress without IV contrast	Usually Not Appropriate	○
SPECT or SPECT/CT MPI rest and stress	Usually Not Appropriate	☼☼☼☼
US echocardiography transthoracic stress	Usually Not Appropriate	○
MRA coronary arteries without IV contrast	Usually Not Appropriate	○
Radiography chest	Usually Not Appropriate	☼

SUMMARY OF LITERATURE REVIEW

Introduction/Background

In the United States, atherosclerotic cardiovascular disease (ASCVD) remains the leading cause of death for both men and women [1]. Although improvements in awareness, knowledge, and medications have led to a decrease in death rates, the burden of disease remains very high [1,2]. Identification of patients who may benefit from early intervention prior to development of symptoms has been shown to reduce mortality and morbidity [3].

Cardiovascular disease prevention has traditionally been based on the assessment of a patient's conventional risk factor profile, a combined evaluation based on genetic, social, physiological, and environmental factors [4]. Risk assessment for ASCVD is intended to aid in determining the appropriate lifestyle changes and pharmacological interventions to reduce a patient's risk of adverse cardiovascular outcomes (eg, myocardial infarction, stroke, cardiovascular death). A global risk score, such as the Framingham Risk Score (FRS), Reynolds risk score, or Systematic Coronary Risk Evaluation, is used to categorize a patient's risk as low, intermediate, or high risk. These risk factors are strong population-based markers but poor individual discriminators of coronary atherosclerotic disease (CAD), and many individuals with one or more risk factors do not experience a cardiac event [4]. Risk calculators are also bound by the underlying prior data that informs them, and as population health

and lifestyle changes, risk calculators become less accurate [5]. More recently, the 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol, suggested the use of a pooled cohort risk calculator in patients to categorize their 10-year risk of ASCVD as low risk (<5%), borderline risk (5% to <7.5%), intermediate risk (7.5% to <20%), or high risk (>20%) [3].

There is a well-established discordance between the prognostic accuracy of current risk estimation scores versus imaging when directly measuring the burden of atherosclerosis for the assessment of individual ASCVD risk as a guide to optimally manage preventive therapies [6]. Imaging allows for the detection of subclinical coronary atherosclerosis. Patients with familial hyperlipidemia in particular have a high prevalence of subclinical coronary atherosclerosis that is independently associated with cardiovascular risk [7]. The coronary artery calcium (CAC) score is a validated measure of overall coronary atherosclerotic burden, the strongest known imaging measure of risk in asymptomatic individuals. Individual data derived from this and other imaging tests provide useful prognostic information for patient management and can complement current risk prediction models [8].

The purpose of this document is to discuss the use of diagnostic imaging tests in asymptomatic patients who are at elevated risk of future cardiovascular events related to

Table 1. Appropriateness category names and definitions

Appropriateness Category Name	Appropriateness Rating	Appropriateness Category Definition
Usually Appropriate	7, 8, or 9	The imaging procedure or treatment is indicated in the specified clinical scenarios at a favorable risk-benefit ratio for patients.
May Be Appropriate	4, 5, or 6	The imaging procedure or treatment may be indicated in the specified clinical scenarios as an alternative to imaging procedures or treatments with a more favorable risk-benefit ratio, or the risk-benefit ratio for patients is equivocal.
May Be Appropriate (Disagreement)	5	The individual ratings are too dispersed from the panel median. The different label provides transparency regarding the panel's recommendation. "May be appropriate" is the rating category and a rating of 5 is assigned.
Usually Not Appropriate	1, 2, or 3	The imaging procedure or treatment is unlikely to be indicated in the specified clinical scenarios, or the risk-benefit ratio for patients is likely to be unfavorable.

atherosclerosis. The tests are to improve targeted preventive efforts based on patient risk and are aimed at identification of CAD.

Appropriate imaging tests in patients who have a known diagnosis of CAD, cardiac symptoms, history of a coronary event, or prior intervention can be found in other ACR Appropriateness Criteria. See the ACR Appropriateness Criteria® topics on "Acute Nonspecific Chest Pain—Low Probability of Coronary Artery Disease" [9], "Chronic Chest Pain-Noncardiac Etiology Unlikely; Low to Intermediate Probability of Coronary Artery Disease" [10], and "Chronic Chest Pain—High Probability of Coronary Artery Disease" [11] for further information and to guide imaging. Please note that the topic of preoperative cardiac evaluation is a distinct and evolving topic, and there is not a current ACR Appropriateness Criteria concerning preoperative evaluations (regarding cardiac or noncardiac surgery); this topic is not considered to be within the scope of this document.

Special Imaging Considerations

Imaging findings in CAD range from direct visualization of plaque and perfusion deficits to ventricular dilatation and infarct [12]. Radiographs depict static findings of late CAD, such as severe coronary artery calcification and heart size, whereas multidetector CT (MDCT) can directly visualize plaque in the coronary arteries. MDCT of the heart is performed with electrocardiogram (ECG) synchronization and can be performed without or with intravenous (IV) contrast—the former defined as CT CAC score—and can assess overall calcific burden of the coronary arteries only; the latter is CT angiography (CTA) coronaries with IV contrast that allows assessment of both noncalcified and calcified plaque and any resultant visualized nonobstructive or obstructive coronary stenosis.

Resting image modalities (ultrasound [US], scintigraphy, CT, and MRI) depict late findings of CAD, such as ventricular dilatation and wall-motion abnormalities, and can

Table 2. Relative radiation level designations

RRL	Adult Effective Dose Estimate Range (mSv)	Pediatric Effective Dose Estimate Range (mSv)
0	0	0
☼	<0.1	<0.03
☼☼	0.1-1	0.03-0.3
☼☼☼	1-10	0.3-3
☼☼☼☼	10-30	3-10
☼☼☼☼☼	30-100	10-30

Note: Relative radiation level (RRL) assignments for some of the examinations cannot be made, because the actual patient doses in these procedures vary as a function of a number of factors (eg, region of the body exposed to ionizing radiation, the imaging guidance that is used). The RRLs for these examinations are designated as "varies."

directly visualize the morphology of infarcted myocardial segments. Intravascular US, an invasive technique, can detect both calcified and noncalcified plaque. Myocardial perfusion can be assessed by stress, rest, and/or delayed imaging, which can be accomplished by US (via assessment of wall-motion changes at stress versus rest, or with contrast echocardiography); cardiac perfusion scintigraphy (via comparison of first-pass radiotracer perfusion to the ventricle at stress versus rest, or measurement of coronary blood flow); and MRI (via comparison of wall-motion or first-pass gadolinium enhancement of the ventricle during stress versus rest). Cardiac MRI is performed with ECG synchronization and may be performed without and with IV contrast and before and after vasodilators or inotropes if stress myocardial perfusion assessment is desired. MR angiography (MRA) of the coronary arteries is possible without or with IV contrast, but it only depicts luminal blood and cannot depict calcified plaque. Myocardial scarring, infarction, and viability can be assessed by cardiac MRI or cardiac PET.

For the purposes of distinguishing between CT and CTA, ACR Appropriateness Criteria topics use the definition in the ACR–NASCI–SIR–SPR Practice Parameter for the Performance and Interpretation of Body Computed Tomography Angiography (CTA) [13]:

CTA uses a thin-section CT acquisition that is timed to coincide with peak arterial or venous enhancement. The resultant volumetric dataset is interpreted using primary transverse reconstructions as well as multiplanar reformations and 3-D renderings.

All elements are essential: 1) timing, 2) reconstructions/reformats, and 3) 3-D renderings. Standard CTs with contrast also include timing issues and reconstructions/reformats. Only in CTA, however, is 3-D rendering a **required** element. This corresponds to the definitions that the CMS has applied to the Current Procedural Terminology codes.

Initial Imaging Definition

Initial imaging is defined as imaging at the beginning of the care episode for the medical condition defined by the variant. More than one procedure can be considered usually appropriate in the initial imaging evaluation when:

- There are procedures that are equivalent alternatives (ie, only one procedure will be ordered to provide the clinical information to effectively manage the patient's care)

OR

- There are complementary procedures (ie, more than one procedure is ordered as a set or simultaneously where each procedure provides unique clinical information to effectively manage the patient's care).

DISCUSSION OF PROCEDURES BY VARIANT

Variant 1: Asymptomatic patient. Low risk for coronary artery disease. Initial imaging

CT Coronary Calcium. The CAC score first became available and validated with electron beam CT, and in the modern era, MDCT is used to acquire this data. Several multicenter trials have assessed the use of CAC in asymptomatic patients at intermediate risk for CAD. In the Multi-Ethnic Study of Atherosclerosis (MESA), Joshi et al [14] followed 6,814 participants including 13% with a CAC score >100 and 34% with a nonzero CAC score with low risk over a period of 10.4 years and found an event rate of 2.4% (33 events). CAC was present in 76% of participants with an event. In the multivariate analysis, only CAC >100 was predictive of coronary heart disease [14]. Recently, in a large cohort of 14,169 low-risk patients with a family history of CAD, Dudum et al showed that a calcium score of >100 had a 2.2 times higher risk for all-cause mortality, 4.3 times higher cardiovascular specific mortality, and 10.4 times higher risk of coronary heart disease, than patients with a zero calcium score [15,16]. Mitchell et al [17] evaluated 23,637 subjects with a mean age of 50.0 ± 8.5 years and low burden of traditional risk factors, they noted relative adjusted subhazard ratio for CAC 1 to 100, 101 to 400, and >400 was 2.2, 3.8, and 5.9 for myocardial infarction; 1.2, 1.4, and 1.9 for stroke; 1.4, 2.0, and 2.8 for major adverse cardiovascular events (MACE); and 1.2, 1.5 and 2.1 for death ($P < .0001$) over a median follow-up period of 11.4 years. Among subjects without traditional risk factors ($n = 6,208$; mean age 43.8 ± 4.4 years), the presence of any CAC (>0 ; $n = 848$) was associated with an increased risk of MACE (adjusted subhazard ratio: 1.67; 95% confidence interval [CI], 1.16-2.39) noting the fact that young individuals without traditional risk factors are typically not offered preventive therapy with statins. Carr et al [18] prospectively enrolled a much younger cohort of 3,043 patients with mean [SD] age 40.3 [3.6] years in the Coronary Artery Risk Development in Young Adults (CARDIA) study and noted that 10.2% of patients had CAC, with a geometric mean Agatston score of 21.6 (interquartile range, 17.3-26.8). During 12.5 years of follow-up, 57 coronary heart disease events and 108 cardiovascular events were observed. After adjusting for demographics, risk factors, and treatments, patients with any CAC experienced a 5-fold increase in coronary heart disease events (hazard ratio [HR], 5.0; 95% CI, 2.8-8.7) and 3-fold increase in cardiovascular events (HR, 3.0; 95% CI, 1.9-4.7), and those with a CAC score of ≥ 100 had an incidence of 22.4 deaths per 100 participants (HR, 3.7; 95% CI, 1.5-10.0).

CTA Coronary Arteries. There is no relevant literature supporting the use of coronary CTA (CCTA) in asymptomatic patients at low risk of CAD. Choi et al [19] identified atherosclerotic plaques in 215 of 1,000 middle-aged asymptomatic patients with 2% prevalence of plaque in the low-risk group.

US Echocardiography Transthoracic Resting. There is no relevant literature supporting the use of transthoracic echocardiography (TTE) resting in asymptomatic patients at low risk of CAD.

US Echocardiography Transthoracic Stress. There is no relevant literature supporting the use of TTE stress in asymptomatic patients at low risk of CAD. The sensitivity and specificity of the test is 72% to 83% and 84% to 95%, respectively, for identification of ischemic myocardium and has been validated only in elevated risk populations [20].

MRA Coronary Arteries. There are very limited data on the utility of MRA in asymptomatic patients, which demonstrated a low yield in a small cohort of non-randomized asymptomatic self-referred patients. In this cohort of 341 patients, 3.8% were found to have significant CAD $\geq 50\%$ stenosis in a protocol that also included MRA, MRI perfusion, and delayed-enhancement imaging; 0.9% of the cohort underwent percutaneous coronary intervention after CAD was detected by cardiac MRI and MRA and was found to have good correlation with stress perfusion MRI in the 13 positive patients, 3 of which were confirmed with invasive angiography [21].

MRI Heart Function With Stress. There is no relevant literature supporting the use of MRI heart function with stress in asymptomatic patients at low risk of CAD.

MRI Heart Function and Morphology. Weir-McCall et al [22] demonstrated an overall low utility of resting MRI via abnormal late gadolinium enhancement in asymptomatic low-risk volunteers (0.67% of whom were found to have abnormalities, including myocardial infarction), with only 0.2% of volunteers having a previously unrecognized myocardial infarction.

Radiography Chest. There is no relevant literature to support the use of chest radiographs to evaluate asymptomatic patients at low risk of CAD.

SPECT/CT MPI Rest and Stress. There is no relevant literature supporting the use of single-photon emission computed tomography (SPECT)/CT myocardial perfusion imaging (MPI) rest and stress in asymptomatic patients at low risk of CAD. Stress SPECT sensitivity and specificity for detection of obstructive CAD ($\geq 50\%$ diameter stenosis) are 74% and 79%, respectively, in symptomatic patients [23].

Variant 2: Asymptomatic patient. Intermediate risk for coronary artery disease. Initial imaging

CT Coronary Calcium. Kondos et al [24] found that any measurable coronary calcium was independently related to hard (death and myocardial infarction) and soft (revascularization procedure) events in men and women at low to intermediate pretest risk; this finding provided incremental prognostic information over conventional risk factors.

Many trials have found evidence of the prognostic use of a CAC score. Shaw et al [25] followed 10,377 asymptomatic patients for 5 ± 3.5 years and found a CAC score to be an independent predictor of death that increased proportionally relative to baseline, with an adjusted relative risk of 1.6, 1.7, 2.5, and 4 for CAC scores of 11 to 100, 101 to 400, 401 to 1,000, and $>1,000$, respectively. Budoff et al [26] also demonstrated incremental risk beyond age, gender, ethnicity, and cardiac risk factors, in evaluating data from 25,253 asymptomatic patients who had a 10-year adjusted survival rate of 99.4% for a CAC score of 0, and 87.8% for a score $>1,000$. In the CARDIA study, of the 2,831 participants, the CAC score and prevalence increased, and an increase in FRS with a score >0 was observed in 7.3%, 20.2%, 19.1%, and 44.8% of individuals with FRSs of 0% to 2.5%, 2.6% to 5%, 5.1% to 10%, and $>10\%$, respectively. A CAC score of ≥ 100 was observed in 1.3%, 2.4%, and 3.5% of those with FRSs of 0% to 2.5%, 2.6% to 5%, and 5.1% to 10%, respectively, without significant change in stratification according to sex and race. The yield of CAC was deemed considerable in the intermediate-risk group [27]. Additionally, the MESA of 6,779 initially asymptomatic individuals also showed CAC as an independent predictor of cerebrovascular events when CAC analysis was stratified by sex or race or ethnicity and improved discrimination for a cerebrovascular event when added to the full model (C-statistic: 0.744 versus 0.755) [28]. Although the authors of the study do not specify the pretest risk of the overall patient population, the presence of hypertension and treated hypertension as well as the degree of coronary calcification was higher in the group containing individuals who had cerebrovascular events.

In the MESA, Polonsky et al [29] used the CAC score, in conjunction with their conventional FRS, to evaluate 5,878 asymptomatic men and women. In that study, the net reclassification index was 25%, an additional 23% of subjects with events were reclassified to the high-risk category, and 13% of subjects without events were reclassified to the low-risk category [29]. The Heinz Nixdorf Recall study, a large population-based study with nearly 5,000 participants and a 5-year follow-up, demonstrated a net

reclassification index of 24% and 19% as high- and low-risk groups, respectively [30]. The recent AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APHA/ASPC/NLA/PCNA guideline on the management of blood cholesterol also suggests patients with a 10-year risk of 5% to 7.5% (borderline risk) in the presence of risk-enhancing factors, such as family history, of premature ASCVD; persistently elevated LDL-C levels ≥ 160 mg/dL; metabolic syndrome; chronic kidney disease; history of preeclampsia; or premature menopause (age <40 years); chronic inflammatory disorders (eg, rheumatoid arthritis, psoriasis, or chronic HIV); high-risk ethnic groups (eg, South Asian); and persistent elevations of triglycerides ≥ 175 mg/dL may benefit from a CAC score prior to initiating a statin therapy [3]. These studies have shown that CAC score predicts future mortality and major cardiac events and aids in improved risk stratification beyond the conventional risk factor–based scores alone. A positive calcium score can reclassify asymptomatic patients with a family history of premature CAD to a high-risk category and can even reclassify those individuals without risk factors as higher risk than those with multiple risk factors but no coronary artery calcification [31,32].

CTA Coronary Arteries. In a recent study, Di Cesare et al [33] showed utility of CCTA in asymptomatic patients at intermediate risk of stenosis detection in 112 out of 185 (60.5%) patients: 56 patients (30.2%) had mild stenosis, 49 patients (26.5%) had moderate stenosis, only 3 patients (1.6%) had severe stenosis, and in 4 patients (2.2%) evaluation could not be determined.

US Echocardiography Transthoracic Resting. There is no relevant literature supporting the use of TTE resting in asymptomatic patients at intermediate risk of CAD.

US Echocardiography Transthoracic Stress. The sensitivity and specificity of stress TTE is 85% and 89%, respectively, has been validated only in populations at elevated risk, and is best utilized to search for obstructive major epicardial coronary stenosis [34]. There is no relevant literature to support the use of stress TTE in asymptomatic patients at intermediate risk of CAD.

MRA Coronary Arteries. MRA of the coronary arteries can assess for arterial patency and pathologic wall thickening but not calcific burden, and it cannot reliably assess small distal vessels [35,36]. There is no relevant literature to support the use of MRA of the coronary arteries in asymptomatic patients at intermediate risk of CAD.

MRI Heart Function and Morphology. There is no relevant literature to support the use of MRI heart function and morphology to evaluate asymptomatic patients at intermediate risk of CAD.

MRI Heart Function With Stress. The IMPACT II study demonstrated a sensitivity of 67% and specificity of 61% for detection of ischemic heart disease, including 533 patients in the intermediate-risk population [37]. The vast majority of the patients in this study were symptomatic, with angina pectoris. The sensitivity and specificity of the test in the asymptomatic intermediate-risk population has not been validated.

Radiography Chest. There is no relevant literature to support the use of chest radiographs alone as a test to evaluate asymptomatic patients at intermediate risk of CAD.

SPECT/CT MPI Rest and Stress. Stress SPECT MPI is the most commonly used stress-imaging technique for patients with suspected or known CAD but has been primarily validated in symptomatic patients, and no studies have examined SPECT MPI alone in asymptomatic intermediate-risk patients. However, in combination with a calcium score of ≥ 400 , expert consensus has speculated that the test may be used for intermediate-risk patients [38]. However, there is no relevant literature to support the use of SPECT MPI in asymptomatic patients at intermediate risk of CAD, except in diabetic patients or those about to undertake a vigorous exercise program [39].

Variant 3: Asymptomatic patient. High risk for coronary artery disease. Initial imaging

CT Coronary Calcium. Studies to date indicate the high prevalence of calcific plaque burden in a high-risk patient, and further testing may be warranted to exclude epicardial stenosis. Data from a large study with 29,312 high-risk patients support the rationale of the “power of zero”; in absence of coronary calcification, only 0.56% participants developed a cardiovascular event during a mean follow-up period of 51 months [40]. A similar observation was also noted in the JUPITER population study in the MESA cohort in which event rates were 0.8 per 1,000 person years with zero calcium score versus 20.2 per 1,000 person years and a calcium score >100 . The estimated number needed to treat for 5 years was at 549 in the zero calcium score group versus 42 in the nonzero calcium score group [41].

CTA Coronary Arteries. There is an added value of coronary atherosclerotic plaque burden as a prognostic benefit in addition to the assessment for epicardial stenosis, such that in high-risk asymptomatic patients, CCTA examinations can be useful. Cho et al [42], in their study of 3,217 asymptomatic patients from the CONFIRM registry stratified according to the magnitude of their CAC score, found that CCTA did provide incremental value in

patients with a CAC score >100. The incremental value of CCTA over FRS was demonstrated in individuals with CAC scores >100 (likelihood ratio χ^2 , 25.34; increment in C-statistic, 0.24; net reclassification index, 0.62; all $P < .001$) but not with CAC scores ≤ 100 (all $P > .05$). For subgroups with CAC score >100, the utility of CCTA for predicting the study end point was evident among individuals whose CAC score ranged from 101 to 400; the observed predictive benefit attenuated with increasing CAC score. In a high-risk population of 665 asymptomatic patients, the multivariate analysis, adjusted for age, gender, and CAC score, obstructive CAD on CCTA ($\geq 50\%$ luminal stenosis) was a significant predictor of adverse events (HR, 5.9; CI, 1.3-26.1). Dedic et al [43] also showed that the addition of CCTA to age and gender, plus a CAC score, increased the C-statistic from 0.81 to 0.84 and resulted in a total net reclassification index of 0.19 ($P < .01$). Incremental value of the CCTA was also demonstrated by Plank et al [44] in a series of 711 patients, where prevalence of a zero calcium score was 306 (43%); out of those, 100 (32.7%) had noncalcified plaque only seen on CTA. With a mean follow-up period of 2.65 years, MACE rate was 0% in CAD negative and higher (1.2%) in CAD positive by CTA.

In a large multicenter registry of 27,125 patients (which included both symptomatic and asymptomatic patients), Min et al [45] found that CCTA improved discrimination by maximal stenosis, number of obstructive vessels, and segment stenosis score (C-index 0.77, 0.77, and 0.78, respectively) beyond age, gender, and CAC score (C-index 0.64) in a small subset of 400 asymptomatic patients. Similarly, CCTA findings improved risk reclassification by per patient maximal stenosis (integrated discrimination improvement [IDI] index, 0.03; $P = .03$) and number of obstructive vessels (IDI index, 0.06; $P = .002$), and by trend for segment stenosis score (IDI, 0.03; $P = .06$).

In the FACTOR-64 study of 900 asymptomatic patients with both type 1 and 2 diabetes, with a mean follow-up time of 4.0 years, the primary outcome event rates were not significantly different between the CCTA and the control groups (6.2% [28 events] versus 7.6% [34 events]; HR, 0.80 [95% CI, 0.49-1.32]; $P = .38$) [46]. A study of 517 asymptomatic subjects also showed that CAD- and plaque-positive remodeling increased MACE prediction compared with a model based on 10-year FRS, carotid disease, and CAC score estimation. In the diabetes subgroup, the percentage of segments with remodeled plaque was the only predictor of MACE [47].

In the asymptomatic diabetic population in patients between the ages of 55 and 74, Halon et al [48] demonstrated 2,242 plaques in 499 subjects with 24 patients with acute coronary syndromes during median

follow-up of 9.2 years. Additional imaging parameters, like plaque volume (upper versus lower quartile HR, 6.9; 95% CI, 1.6-30.8; $P = .011$), percentage of low-density plaque content <50 Hounsfield units (HR, 14.2; 95% CI, 1.9-108; $P = .010$), and mild plaque calcification (HR versus all other plaques, 3.3; 95% CI, 1.5-7.3; $P = .004$), predicted plaque events univariately after adjustment by clinical risk score. In this series, a culprit plaque event occurred in 13 of 376 (3.5%) high-risk plaques (plaques with ≥ 2 risk predictors) versus 11 of 1,866 (0.6%) in non-high-risk plaques ($P < .0001$), at 12 of 343 (3.5%) stenotic sites ($\geq 50\%$) versus 12 of 1,899 (0.6%) nonstenotic sites ($P < .0001$), and in 7 of 131 (5.3%) high-risk plaques with stenosis ($P < .0001$ versus all others) [48]. A systemic review and meta-analysis composed of 10 studies, including 5,012 asymptomatic participants with diabetes who underwent CCTA, found that presence of obstructive CAD on CCTA (versus nonobstructive or no CAD) was associated with a significantly elevated risk of adverse events (summary HR, 4.07; 95% CI, 2.30-7.21). Beller et al [49] observed estimated HR for nonobstructive plaque (versus no CAD) was 2.17 (95% CI, 1.11-4.25). The pooled HRs per unit for segment stenosis score and segment involvement score were 1.44 (95% CI, 0.98-2.12) and 1.73 (95% CI, 1.07-2.80) respectively. The authors concluded that the presence and extent of CAD on CCTA were strong, independent predictors of cardiovascular events in asymptomatic individuals with diabetes.

US Echocardiography Transthoracic Resting. There is no relevant literature supporting the use of resting TTE in asymptomatic patients at high risk of CAD.

US Echocardiography Transthoracic Stress. TTE performed at rest and stress can assess for inducible wall-motion abnormalities, thus revealing ischemic heart disease. The sensitivity and specificity of the test is 72% to 83% and 84% to 95%, respectively, and has been validated only in symptomatic elevated risk populations and is best utilized to identify obstructive major epicardial coronary stenosis [20]. There is no relevant literature to support the use of stress TTE in asymptomatic patients at high risk of CAD.

MRA Coronary Arteries. MRA of the coronary arteries can assess for arterial patency and pathologic wall thickening but not calcific burden, and it cannot reliably assess small, distal vessels [35,36]. There is no relevant literature to support the use of MRA of the coronary arteries in asymptomatic patients at high risk of CAD.

MRI Heart Function With Stress. There is no relevant literature supporting the use of MRI heart function with stress in asymptomatic patients at high risk of CAD.

MRI Heart Function and Morphology. Cardiac MRI can assess resting left ventricular function and potential ischemic scar burden. There is no relevant literature supporting the use of cardiac MRI as a screening test in asymptomatic patients at high risk of CAD.

Radiography Chest. There is no relevant literature supporting the use of chest radiographs over other modalities as a screening test in asymptomatic patients at high risk of CAD.

SPECT/CT MPI Rest and Stress. Stress SPECT MPI is the most commonly used stress-imaging technique for patients with suspected or known CAD but has been primarily validated in symptomatic patients. Stress SPECT pooled sensitivity and specificity for detection of obstructive CAD ($\geq 50\%$ diameter stenosis) was 74% and 79%, respectively, as validated chiefly in symptomatic patients [50]. Young et al [51], in their study Detection of Ischemia in Asymptomatic Diabetics (DIAD), a randomized controlled trial, enrolled and randomized 1,123 asymptomatic participants with type 2 diabetes to either an adenosine-stress radionuclide MPI or no screening imaging, showed the positive predictive value of having moderate or large MPI defects was only 12% with 7 nonfatal myocardial ischemia's and 8 cardiac deaths (2.7%) in the screened MPI group and 10 nonfatal myocardial ischemia's and 7 cardiac deaths (3.0%) among the not-screened group (HR, 0.88; 95% CI, 0.44-1.88; $P = .73$). Overall, cardiac event rates were not significantly reduced by MPI screening for myocardial ischemia over 4.8 years. In a prospective multicenter BARDOT trial, 22% of asymptomatic high-risk patients with diabetes had abnormal SPECT, but those with abnormal SPECT randomized to medical versus invasive-medical strategies had similar event rates ($P = .215$) [52]. For a selected subgroup of asymptomatic patients with diabetes, data suggest routine use of SPECT as a screening test is likely to have a lower yield as well as limited effect on clinical outcomes [53]. However, the most recent ACC/AHA/ASNC Appropriate Use Criteria for SPECT MPI states SPECT would be useful when the calcium score is >400 or 100 to 400 if the patient is at high risk of CAD [38].

SUMMARY OF RECOMMENDATIONS

- **Variant 1:** Imaging is usually not appropriate for the initial evaluation of an asymptomatic patient with low risk of coronary artery disease.
- **Variant 2:** CT coronary calcium is usually appropriate for the initial imaging of an asymptomatic patient with intermediate risk of coronary artery disease.

- **Variant 3:** CT coronary calcium or CTA coronary arteries with IV contrast may be appropriate for the initial imaging of an asymptomatic patient with high risk of coronary artery disease.

SUPPORTING DOCUMENTS

The evidence table, literature search, and appendix for this topic are available at <https://acsearch.acr.org/list>. The appendix includes the strength of evidence assessment and the final rating round tabulations for each recommendation.

For additional information on the Appropriateness Criteria methodology and other supporting documents go to www.acr.org/ac.

RELATIVE RADIATION LEVEL INFORMATION

Potential adverse health effects associated with radiation exposure are an important factor to consider when selecting the appropriate imaging procedure. Because there is a wide range of radiation exposures associated with different diagnostic procedures, a relative radiation level (RRL) indication has been included for each imaging examination. The RRLs are based on effective dose, which is a radiation dose quantity that is used to estimate population total radiation risk associated with an imaging procedure. Patients in the pediatric age group are at inherently higher risk from exposure, because of both organ sensitivity and longer life expectancy (relevant to the long latency that appears to accompany radiation exposure). For these reasons, the RRL dose estimate ranges for pediatric examinations are lower as compared with those specified for adults (see Table 2). Additional information regarding radiation dose assessment for imaging examinations can be found in the ACR Appropriateness Criteria® Radiation Dose Assessment Introduction document [54].

REFERENCES

1. Cooper R, Cutler J, Desvigne-Nickens P, et al. Trends and disparities in coronary heart disease, stroke, and other cardiovascular diseases in the United States: findings of the national conference on cardiovascular disease prevention. *Circulation* 2000;102:3137-47.
2. Greenland P, Alpert JS, Beller GA, et al. 2010 ACCF/AHA guideline for assessment of cardiovascular risk in asymptomatic adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2010;56:e50-103.
3. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2019;73:e285-350.
4. Greenland P, Knoll MD, Stamler J, et al. Major risk factors as antecedents of fatal and nonfatal coronary heart disease events. *JAMA* 2003;290:891-7.

5. Goff DC Jr, Lloyd-Jones DM, Bennett G, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014;63:2935-59.
6. Wilson SR, Lin FY, Min JK. Role of coronary artery calcium score and coronary CT angiography in the diagnosis and risk stratification of individuals with suspected coronary artery disease. *Curr Cardiol Rep* 2011;13:271-9.
7. Perez de Isla L, Alonso R, Muniz-Grijalvo O, et al. Coronary computed tomographic angiography findings and their therapeutic implications in asymptomatic patients with familial hypercholesterolemia. Lessons from the SAFEHEART study. *J Clin Lipidol* 2018;12:948-57.
8. Detrano R, Guerci AD, Carr JJ, et al. Coronary calcium as a predictor of coronary events in four racial or ethnic groups. *N Engl J Med* 2008;358:1336-45.
9. American College of Radiology. ACR Appropriateness Criteria®: Acute Nonspecific Chest Pain—Low Probability of Coronary Artery Disease. Available at: <https://acsearch.acr.org/docs/69401/Narrative/>. Accessed September 30, 2020.
10. Shah AB, Kirsch J, Bolen MA, et al. ACR Appropriateness Criteria® Chronic Chest Pain-Noncardiac Etiology Unlikely-Low to Intermediate Probability of Coronary Artery Disease. *J Am Coll Radiol* 2018;15:S283-90.
11. Akers SR, Panchal V, Ho VB, et al. ACR Appropriateness Criteria® Chronic Chest Pain-High Probability of Coronary Artery Disease. *J Am Coll Radiol* 2017;14:S71-80.
12. Nesto RW, Kowalchuk GJ. The ischemic cascade: temporal sequence of hemodynamic, electrocardiographic and symptomatic expressions of ischemia. *Am J Cardiol* 1987;59:23C-30C.
13. American College of Radiology. ACR–NASCI–SIR–SPR Practice Parameter for the Performance and Interpretation of Body Computed Tomography Angiography (CTA). Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/body-cta.pdf>. Accessed September 30, 2020.
14. Joshi PH, Patel B, Blaha MJ, et al. Coronary artery calcium predicts cardiovascular events in participants with a low lifetime risk of cardiovascular disease: The Multi-Ethnic Study of Atherosclerosis (MESA). *Atherosclerosis* 2016;246:367-73.
15. Hecht H, Blaha MJ, Berman DS, et al. Clinical indications for coronary artery calcium scoring in asymptomatic patients: expert consensus statement from the Society of Cardiovascular Computed Tomography. *J Cardiovasc Comput Tomogr* 2017;11:157-68.
16. Dudum R, Dzaye O, Mirbolouk M, et al. Coronary artery calcium scoring in low risk patients with family history of coronary heart disease: validation of the SCCT guideline approach in the coronary artery calcium consortium. *J Cardiovasc Comput Tomogr* 2019;13:21-5.
17. Mitchell JD, Paisley R, Moon P, Novak E, Villines TC. Coronary artery calcium and long-term risk of death, myocardial infarction, and stroke: the Walter Reed Cohort Study. *JACC Cardiovasc Imaging* 2018;11:1799-806.
18. Carr JJ, Jacobs DR Jr, Terry JG, et al. Association of coronary artery calcium in adults aged 32 to 46 years with incident coronary heart disease and death. *JAMA Cardiol* 2017;2:391-9.
19. Choi EK, Choi SI, Rivera JJ, et al. Coronary computed tomography angiography as a screening tool for the detection of occult coronary artery disease in asymptomatic individuals. *J Am Coll Cardiol* 2008;52:357-65.
20. Heijnenbroek-Kal MH, Fleischmann KE, Hunink MG. Stress echocardiography, stress single-photon-emission computed tomography and electron beam computed tomography for the assessment of coronary artery disease: a meta-analysis of diagnostic performance. *Am Heart J* 2007;154:415-23.
21. Song KD, Kim SM, Choe YH, et al. Integrated cardiac magnetic resonance imaging with coronary magnetic resonance angiography, stress-perfusion, and delayed-enhancement imaging for the detection of occult coronary artery disease in asymptomatic individuals. *Int J Cardiovasc Imaging* 2015;31(Suppl 1):77-89.
22. Weir-McCall JR, Fitzgerald K, Papagiorgopulo CJ, et al. Prevalence of unrecognized myocardial infarction in a low-intermediate risk asymptomatic cohort and its relation to systemic atherosclerosis. *Eur Heart J Cardiovasc Imaging* 2017;18:657-62.
23. Mordi IR, Badar AA, Irving RJ, Weir-McCall JR, Houston JG, Lang CC. Efficacy of noninvasive cardiac imaging tests in diagnosis and management of stable coronary artery disease. *Vasc Health Risk Manag* 2017;13:427-37.
24. Kondos GT, Hoff JA, Sevrakov A, et al. Electron-beam tomography coronary artery calcium and cardiac events: a 37-month follow-up of 5635 initially asymptomatic low- to intermediate-risk adults. *Circulation* 2003;107:2571-6.
25. Shaw LJ, Raggi P, Schisterman E, Berman DS, Callister TQ. Prognostic value of cardiac risk factors and coronary artery calcium screening for all-cause mortality. *Radiology* 2003;228:826-33.
26. Budoff MJ, Shaw LJ, Liu ST, et al. Long-term prognosis associated with coronary calcification: observations from a registry of 25,253 patients. *J Am Coll Cardiol* 2007;49:1860-70.
27. Okwuosa TM, Greenland P, Ning H, Liu K, Lloyd-Jones DM. Yield of screening for coronary artery calcium in early middle-age adults based on the 10-year Framingham Risk Score: the CARDIA study. *JACC Cardiovasc Imaging* 2012;5:923-30.
28. Gibson AO, Blaha MJ, Arnan MK, et al. Coronary artery calcium and incident cerebrovascular events in an asymptomatic cohort. The MESA Study. *JACC Cardiovasc Imaging* 2014;7:1108-15.
29. Polonsky TS, McClelland RL, Jorgensen NW, et al. Coronary artery calcium score and risk classification for coronary heart disease prediction. *JAMA* 2010;303:1610-6.
30. Erbel R, Mohlenkamp S, Moebus S, et al. Coronary risk stratification, discrimination, and reclassification improvement based on quantification of subclinical coronary atherosclerosis: the Heinz Nixdorf Recall study. *J Am Coll Cardiol* 2010;56:1397-406.
31. Mulders TA, Taraboanta C, Franken LC, et al. Coronary artery calcification score as tool for risk assessment among families with premature coronary artery disease. *Atherosclerosis* 2016;245:155-60.
32. Nasir K, Rubin J, Blaha MJ, et al. Interplay of coronary artery calcification and traditional risk factors for the prediction of all-cause mortality in asymptomatic individuals. *Circ Cardiovasc Imaging* 2012;5:467-73.
33. Di Cesare E, Patriarca L, Panebianco L, et al. Coronary computed tomography angiography in the evaluation of intermediate risk asymptomatic individuals. *Radiol Med* 2018;123:686-94.
34. Picano E, Molinaro S, Pasanisi E. The diagnostic accuracy of pharmacological stress echocardiography for the assessment of coronary artery disease: a meta-analysis. *Cardiovasc Ultrasound* 2008;6:30.
35. Cruz G, Atkinson D, Henningsson M, Botnar RM, Prieto C. Highly efficient nonrigid motion-corrected 3D whole-heart coronary vessel wall imaging. *Magn Reson Med* 2017;77:1894-908.
36. Moghari MH, Roujol S, Henningsson M, et al. Three-dimensional heart locator for whole-heart coronary magnetic resonance angiography. *Magn Reson Med* 2014;71:2118-26.
37. Schwitzer J, Wacker CM, Wilke N, et al. MR-IMPACT II: Magnetic Resonance Imaging for Myocardial Perfusion Assessment in Coronary Artery Disease trial: perfusion-cardiac magnetic resonance vs. single-photon emission computed tomography for the detection of coronary artery disease: a comparative multicentre, multivendor trial. *Eur Heart J* 2013;34:775-81.
38. Hendel RC, Abbott BG, Bateman TM, et al. The role of radionuclide myocardial perfusion imaging for asymptomatic individuals. *J Nucl Cardiol* 2011;18:3-15.
39. Multimodality Writing Group for Stable Ischemic Heart Disease Wolk MJ, Bailey SR, et al. ACCF/AHA/ASE/ASNC/HFSA/HRS/SCAI/SCCT/SCMR/STS 2013 multimodality appropriate use criteria for the detection and risk assessment of stable ischemic heart disease: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Heart Association, American Society of Echocardiography, American Society of Nuclear

Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *J Card Fail* 2014;20:65-90.

40. Sarwar A, Shaw LJ, Shapiro MD, et al. Diagnostic and prognostic value of absence of coronary artery calcification. *JACC Cardiovasc Imaging* 2009;2:675-88.
41. Blaha MJ, Budoff MJ, DeFilippis AP, et al. Associations between C-reactive protein, coronary artery calcium, and cardiovascular events: implications for the JUPITER population from MESA, a population-based cohort study. *Lancet* 2011;378:684-92.
42. Cho I, Chang HJ, B OH, et al. Incremental prognostic utility of coronary CT angiography for asymptomatic patients based upon extent and severity of coronary artery calcium: results from the COronary CT Angiography EvaluationN For Clinical Outcomes InterNational Multicenter (CONFIRM) study. *Eur Heart J* 2015;36:501-8.
43. Dedic A, Ten Kate GJ, Roos CJ, et al. Prognostic value of coronary computed tomography imaging in patients at high risk without symptoms of coronary artery disease. *Am J Cardiol* 2016;117:768-74.
44. Plank F, Friedrich G, Dichtl W, et al. The diagnostic and prognostic value of coronary CT angiography in asymptomatic high-risk patients: a cohort study. *Open Heart* 2014;1:e000096.
45. Min JK, Labounty TM, Gomez MJ, et al. Incremental prognostic value of coronary computed tomographic angiography over coronary artery calcium score for risk prediction of major adverse cardiac events in asymptomatic diabetic individuals. *Atherosclerosis* 2014;232:298-304.
46. Muhlestein JB, Lappe DL, Lima JA, et al. Effect of screening for coronary artery disease using CT angiography on mortality and cardiac events in high-risk patients with diabetes: the FACTOR-64 randomized clinical trial. *JAMA* 2014;312:2234-43.
47. Guaricci AI, Lorenzoni V, Guglielmo M, et al. Prognostic relevance of subclinical coronary and carotid atherosclerosis in a diabetic and nondiabetic asymptomatic population. *Clin Cardiol* 2018;41:769-77.
48. Halon DA, Lavi I, Barnett-Griness O, et al. Plaque morphology as predictor of late plaque events in patients with asymptomatic type 2 diabetes: a long-term observational study. *JACC Cardiovasc Imaging* 2019;12:1353-63.
49. Beller E, Meinel FG, Schoeppe F, et al. Predictive value of coronary computed tomography angiography in asymptomatic individuals with diabetes mellitus: systematic review and meta-analysis. *J Cardiovasc Comput Tomogr* 2018;12:320-8.
50. Takx RA, Blomberg BA, El Aidi H, et al. Diagnostic accuracy of stress myocardial perfusion imaging compared to invasive coronary angiography with fractional flow reserve meta-analysis. *Circ Cardiovasc Imaging* 2015;8:1-7.
51. Young LH, Wackers FJ, Chyun DA, et al. Cardiac outcomes after screening for asymptomatic coronary artery disease in patients with type 2 diabetes: the DIAD study: a randomized controlled trial. *JAMA* 2009;301:1547-55.
52. Zellweger MJ, Maraun M, Osterhues HH, et al. Progression to overt or silent CAD in asymptomatic patients with diabetes mellitus at high coronary risk: main findings of the prospective multicenter BARDOT trial with a pilot randomized treatment substudy. *JACC Cardiovasc Imaging* 2014;7:1001-10.
53. Budoff MJ, Raggi P, Beller GA, et al. Noninvasive cardiovascular risk assessment of the asymptomatic diabetic patient: the Imaging Council of the American College of Cardiology. *JACC Cardiovasc Imaging* 2016;9:176-92.
54. American College of Radiology. ACR Appropriateness Criteria® radiation dose assessment introduction. Available at: <https://www.acr.org/-/media/ACR/Files/Appropriateness-Criteria/RadiationDoseAssessmentIntro.pdf>. Accessed September 30, 2020.