

Masseteric-Facial Anastomosis: A Report of Three Cases

Anastomose massetérico-facial: Um relato de três casos

Breno Alexander Bispo¹ Paulo Eduardo Albuquerque Zito Raffa²
 Pedro Henrique Simm Pires de Aguiar³ André Alexandre Bocchi¹
 Maria Eugênia Martins Publio Correa⁴ Victor de Oliveira Correa⁴
 Carlos Alberto de Almeida Castro Junior⁵ Paulo Henrique Pires de Aguiar^{6,7,8}
 Roberto Sergio Martins⁹

¹ Faculdade de Medicina São Leopoldo Mandic, Campinas, São Paulo, SP, Brazil

² Faculdade de Medicina de Catanduva, Catanduva, São Paulo, SP, Brazil

³ Pontifícia Universidade Católica de São Paulo, Brazil

⁴ Faculdade de Medicina, Universidade Santo Amaro, São Paulo, SP, Brazil

⁵ Fellowship in Neurosurgery, Hospital Santa Paula, São Paulo, SP, Brazil

⁶ Department of Research and Innovation, Laboratory of Cellular and Molecular Biology, Faculdade de Medicina do ABC, Santo André, São Paulo, SP, Brazil

⁷ Pain Center and Department of Neurosurgery, Hospital Santa Paula, São Paulo, São Paulo, SP, Brazil

⁸ Department of Neurology, Faculdade de Medicina de Sorocaba, Pontifícia Universidade Católica de São Paulo, Sorocaba, São Paulo, SP, Brazil

⁹ Peripheral Nerve Surgery Unit, Division of Functional Neurosurgery, Instituto de Psiquiatria, Faculdade de Medicina da Universidade de São Paulo, São Paulo, São Paulo, SP, Brazil

Address for correspondence Breno Alexander Bispo, Bachelor of Medicine, Rua Santo Antônio 431, apto. 93, Cambuí, Campinas, São Paulo, SP, Brazil (e-mail: brenobispo26@gmail.com).

Arq Bras Neurocir 2021;40(4):e380–e386.

Abstract

Keywords

- facial paralysis
- facial anastomosis
- masseteric nerve
- nerve graft

Intoduction The pathways of the facial nerve are variable, and knowledge of that is essential. The worst impact caused by facial paralysis is related to quality of life, especially regarding the self-esteem and social acceptance on the part of the patients, leading to social isolation and disruption on their mental health.

Case Report A 33-year-old female patient, with a stage-T3 acoustic neurinoma, presented with a moderate dysfunction (grades II to III) according to the House-Brackmann (HB) Facial Nerve Grading System. A 43-year-old male patient, with a stage-T4B trigeminal schwannoma, underwent a resective surgery and presented grade-VI dysfunction according to the HB scale. And a female patient with a stage-T4A acoustic neurinoma presented grade-IV dysfunction according to the HB scale.

Discussion We performed a literature review of papers related to surgeries for masseteric-facial nerve anastomosis and compiled the results in table; then, we compared these data with those obtained from our cases.

received
August 9, 2020
accepted
February 22, 2021
published online
August 31, 2021

DOI <https://doi.org/10.1055/s-0041-1730338>.
ISSN 0103-5355.

© 2021. Sociedade Brasileira de Neurocirurgia. All rights reserved. This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
 Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

Conclusion The masseteric nerve is the one that shows the best prognosis among all the cranial nerves that could be used, but it is also necessary to perform well the surgical technique to access the facial branch and consequently achieve a better masseteric-facial nerve anastomosis.

Resumo

Introdução As vias do nervo facial são variáveis, e o conhecimento delas é essencial. O pior impacto causado pela paralisia facial está relacionado à qualidade de vida, principalmente no que se refere à autoestima e aceitação social dos pacientes, e leva ao isolamento social e a transtornos na saúde mental.

Relato de caso Paciente do sexo feminino, de 33 anos, com neurinoma acústico de estágio T3, apresentou disfunção moderada (graus II a III) na escala de nervo facial de House-Brackmann (HB). Paciente do sexo masculino, de 43 anos, portador de schwannoma trigeminal de estágio T4B, foi submetido a cirurgia de ressecção do tumor e apresentou, no pós-operatório, disfunção de grau VI na escala de HB. E paciente do sexo feminino com neurinoma acústico de estágio T4A, apresentava disfunção de grau IV na escala de HB.

Discussão Realizamos uma revisão da literatura sobre os principais trabalhos relacionados às cirurgias de anastomose massetérico-facial, compilamos os resultados em uma tabela, e, em seguida, comparamos os dados obtidos com nossos casos.

Conclusão O nervo massetérico é o que apresenta melhor prognóstico entre todas os nervos cranianos que poderiam ser utilizados; entretanto, para isso é necessário realizar uma boa técnica cirúrgica para garantir um adequado acesso ao ramo facial e, consequentemente realizar uma melhor anastomose massetérico-facial

Palavras-chave

- ▶ paralisia facial
- ▶ anastomose facial
- ▶ nervo massetérico
- ▶ enxerto de nervo

Introduction

The facial nerve (CN VII), is a mixed cranial nerve with motor, parasympathetic, and sensory branches;¹ it is made up of four nuclei that serve different functions, such as: movement of muscles that produce facial expression; parasympathetic innervation to the lacrimal, submaxillary, and submandibular glands; the sensation of the external ear; and the sensation of taste.² The pathways of the facial nerve are variable, and that knowledge is essential to perform an accurate physical diagnosis and safe and effective surgical intervention in the head and neck.

This nerve is vulnerable to several injuries which can result in facial paralysis. Facial paralysis is one of the most common cranial neuropathies, and its main clinical repercussion in patients is a dysfunction in facial expression³. The facial paralysis can range from the slightest asymmetry to complete paralysis on one or both sides of the face, and can be classified through the House-Brackmann (HB) Facial Nerve Grading System, which was developed as an indicator of severity; however, it is also used as a progress record for facial paralysis treatments. The grades of involvement on this scale range from I to VI (► **Table 1**); grades I and II have good outcomes, grades III and IV represent moderate dysfunction, and grades V and VI have bad outcomes.⁴

The worst impact caused by facial paralysis is related to quality of life, especially regarding the self-esteem and social acceptance on the part of the patients, leading to social

isolation and disruption in their mental health. Moreover, facial paralysis increases the risk of corneal damage and infection.⁵ Therefore, several treatment options have emerged, including reanimation surgery via anastomosis. The surgical reanimation of facial movements and, above all, of the facial expression are a challenge for all microsurgeons, and knowledge of the surgical techniques becomes fundamental to provide the best treatment. In patients with facial paralysis, the local cranial nerves represent a good source of motor axons that can be redirected for facial reanimation.⁶ Besides that, masseteric-facial nerve anastomosis has been shown to be an excellent surgical treatment for the facial paralysis in the recent academic literature.

Masseteric-facial nerve anastomosis can be performed through a surgery to redirect the motor axons of the trigeminal cranial nerve (masseteric motor branch), which is extremely abundant in motor axons, to reverse and resuscitate the facial muscles, recovering facial expression. In the present paper, we report three cases of masseteric-facial anastomosis, and compare the data obtained from them with the data from a literature review on facial-anastomosis surgeries (► **Table 2**).

Methods

Surgical Technique

Under general anesthesia, a modified rhytidectomy incision is marked (as it provides good exposure and has the advantage of resulting in less complications and better cosmetic

Table 1 Categorization of facial lesion through the House-Brackmann Facial Nerve Grading System

Grade	Description	Characteristics	
I	Normal	Normal facial function in all areas	
II	Mild dysfunction	<i>Gross:</i>	Slight weakness noticeable on close inspection; may have very slight synkinesis
		<i>At rest:</i>	Normal symmetry and tone
		<i>Motion:</i>	Forehead: moderate to good function Eye: complete closure with minimal effort Mouth: slight asymmetry
III	Moderate dysfunction	<i>Gross:</i>	Obvious but no disfiguring difference between two sides; noticeable but not severe synkinesis, contracture, and/or hemifacial spasm
		<i>At rest:</i>	Normal symmetry and tonus
		<i>Motion:</i>	Forehead: slight to moderate movement Eye: complete closure with effort Mouth: slightly weak with maximum effort
IV	Moderately severe dysfunction	<i>Gross:</i>	Obvious weakness and/or disfiguring asymmetry
		<i>At rest:</i>	Normal symmetry and tone
		<i>Motion:</i>	Forehead: none Eye: incomplete closure Mouth: asymmetry with maximum effort
V	Severe dysfunction	<i>Gross:</i>	Only barely perceptible motion
		<i>At rest:</i>	Asymmetry
		<i>Motion:</i>	Forehead: none Eye: incomplete closure Mouth: slight movement
VI	Total paralysis		No movement

outcomes⁷), and epinephrine (dilution: 1:200,000) is infiltrated subcutaneously along the incision line. A point 3 cm anterior to the tragus and 1 cm inferior to the zygomatic arch (surface landmark for the masseteric nerve) is also marked. Skin flaps are elevated anteriorly to this point, and the greater auricular nerve is delineated in its entire course until its entry into the parotid gland (► **Figure 1**). If the need arises, this can be used as an interposition nerve graft. The parotid gland is cut in half, and the entire facial nerve trunk, from its exit from the stylomastoid foramen and the first 2 cm of the main branches after bifurcation to the buccinator and obicularis oculi muscles, is exposed and divided close to the exit from the stylomastoid foramen (to gain additional nerve length for the anastomosis). The masseteric nerve is then identified with the zygomatic arch, the posterior border of the masseter muscle, and the lower part of the temporal nerve as standard landmarks. It is advisable to enter the masseter ~ 1 cm anterior to the posterior border. As the nerve lies deep at the level of the lower border of the zygomatic arch, detachment of the masseter from the zygomatic arch is not necessary. Care must be taken to preserve the branches of the facial nerve over the masseter. The masseteric nerve lies 1.5 cm to 2.0 cm deep from the muscle surface within its belly closer to its internal surface than its external surface. Gentle dissection of the muscle fibers usually reveals the nerve, which courses from its origin deep into the zygomatic arch pointing toward the oral commissure. For the anastomosis, the proximal branches

can be spared, preventing total denervation of the masseter muscle.⁸ The masseteric nerve is sectioned, and the anastomosis is performed directly to the most affected portion of the facial nerve (the mouth, the eye or even the main trunk). (► **Figure 2**)

Case Report

First Case

A 33-year-old female patient, with a stage-T3 acoustic neuroma according to the Samii classification.⁹ The patient had a grade-IV to V facial paralysis on the right side, according to the HB scale (► **Table 1**). She was submitted to an elective surgery for tumor removal and radiotherapy (radiosurgery) in the meatal lesion, which decreased the grade of the paralysis to II to III according to the HB scale. After 9 months, she underwent another elective surgery for masseteric-facial nerve anastomosis, which resulted in full recovery of function four months postoperatively (► **Figure 3**).

Second Case

A 43-year-old male patient, with a stage-T4B trigeminal schwannoma. The patient underwent a resective surgery and presented grade-VI paralysis according to the HB scale. After six months of the resective surgery, the patient underwent a hypoglossal-facial anastomosis surgery to improve facial function, which was not possible due to the technical conditions.

Table 2 Summary of the literature review

Author (year)	Preoperative HB grade (n)	Results	Conclusion
Bianchi et al. ¹¹ (2018)	HB VI (6)	4 patients were evaluated as grade III in the scale and 2 patients were evaluated as grade II.	The masseteric/cross facial nerve grafting might be used in patients with palsies for 20 to 24 months and achieve great results.
Altamami et al. ¹² (2019)	HB VI (7)	The masseteric anastomosis of the facial nerve did not show improvement on forehead movements; however, there was some improvement in the movement of the mouth and smile.	The masseteric-facial nerve anastomosis is a viable option for patients with facial nerve paralysis.
Biglioli et al. ¹⁵ (2018)	HB VI (24)	11 patients were HB grade II postoperatively; 11 patients were HB grade III postoperatively; and 2 patients were HB grade IV postoperatively.	This surgical technique showed to be safe and effective, improving the degree of paralysis.
Lee et al. ¹⁰ (2020)	Not available	3 patients had an excellent resting symmetry postoperatively, and all 4 had good eye closure. 3 had good oral excursion, and 1 had excellent. 3 patients were evaluated with good smiles. 2 patients were evaluated with excellent, and 1, with good oral continence.	The masseteric anastomosis shows good results when it comes down to dynamic facial motion.
Vincent et al. ¹³ (2019)	HB III or IV (7)	HB scores ranged from II to III	After masseteric facial nerve transfer, the patients evolved with significant improvement in smile symmetry and lower facial synkinesis as measured with the eFace tool.
Sakthivel et al. ⁸ (2020)	HB VI (6)	1 HB I, 2 patients HB II and 3 patients HB V	Compared to other techniques masseteric facial nerve anastomosis is a versatile powerful early facial dynamic reanimation tool with almost negligible morbidity for patients with complete facial nerve paralysis.
Gray et al. ¹⁷ (2020)	HB V or VI (8)	Preoperative smile deviation in commissure excursion of 8.8 mm, and a postoperative deviation of 3.8 mm. The deviation angle of the smile improved from 10.3 degrees to 5.2 degrees.	Single-stage masseteric-zygomatic nerve transfer is a useful procedure for patients with synkinesis caused by facial nerve paralysis.
Hontanilla et al. ¹⁶ (2018)	Not available	Spontaneous recovery of the smile during more than 50% of the patient's daily life was of 80% in the group submitted to cross-facial nerve grafting, and of 55.5% in group submitted to masseteric-facial nerve anastomosis, which, surprisingly presented a higher degree of satisfaction.	With the masseteric nerve, better symmetry, a higher degree of recovery, and an increased level of satisfaction are achieved in a single-stage operation. Furthermore, both nerve sources are able to restore spontaneity in more than 50% of the patient's daily life, with no significant differences between them.
Mohanty et al. ¹⁸ (2018)	HB VI (3) IV (1)	Motion was observed 8 days after external neurolysis of the neurovascular bundle in patient 1, and 4 months in the second free functional muscle transfer group (patients 2, 3, and 4). The average excursion and philtral correction were of 11.2 mm and 2.39 mm respectively.	Reusing the masseteric nerve to innervate another free functional muscle transfer is a feasible and reliable option if there are no obvious contraindications to another free functional muscle transfer and masseter muscle contraction is clearly demonstrable preoperatively.

Abbreviation: HB, House-Brackmann Facial Nerve Grading System.

Subsequently, a masseteric-facial anastomosis surgery was attempted, which, unfortunately, even after 2 years, did not achieve good results, and the patient remained with altered facial function, with grade-VI paralysis on the HB scale.

Third Case

A female patient with a stage-T4A acoustic neuroma; after the resective surgery, she presented grade-IV paralysis on the HB scale, that is, a significant impairment in facial function. She was

submitted to a masseteric-facial nerve anastomosis surgery, and, two months postoperatively, presented grade-III paralysis on the HB scale, being able to close the eyes more and smile somewhat more widely, expressing excellent surgical results.

Discussion

Incomplete facial paralysis still remains a surgical challenge. Techniques such as masseteric-facial anastomosis are




Fig. 1 Representation of the surgical incision and the division of the branches of the facial nerve (CN VII) in the parotid gland.¹⁹




Fig. 2 (A) Surgical Approach using the modified rhytidectomy incision; (B) opening of the masseteric fascia; (C) inspection of the masseteric nerve; (D) anastomosis of the masseteric nerve; (E) inspection of the preservation of the other nerves; (F) suture procedures.

gaining popularity and showing good results. We performed a literature review on the main papers related to masseteric-facial nerve anastomosis surgeries and compiled the results obtained in ► **Table 2**.

We compared the data obtained from the literature with the data from our cases. In our first case, the patient had a moderate dysfunction (grades II to III on the HB scale), and, just four months after the masseteric-facial anastomosis, she

evolved to complete recovery of function. Analyzing the third case, one can see that an excellent evolution has also occurred, for the patient, who had a moderately severe dysfunction (grade IV on the HB scale), evolved to grade III in two months, and was able to close the eyes more and smile somewhat more widely. Unfortunately, this did not happen with our second case, who, even after two years of the masseteric-facial anastomosis, did not show a significant




Fig. 3 (A) Bilateral and harmonious movement of the facial muscles four months after the surgery; (B) normal movement of the face muscles on the right side four months after the surgery; (C) normal movement of the muscles of the left side (unaffected side); (D) bilateral and harmonious movement of the facial muscles four months after the surgery.

evolution. Considering the three cases herein reported, the third had a good outcome, the first had an excellent outcome and fast evolution, and the second presented bad outcome; therefore two out of three patients presented good evolution.

Our results corroborate with those of Lee et al.¹⁰ (2020), who reported that 75% of the patients who underwent masseteric innervation obtained excellent results regarding the symmetry of the face at rest. They also included in their results an improvement in the ability to close the eyes, in the movement of the mouth, and in the ability to smile. We can also mention the studies by Bianchi et al.¹¹ (2018) and Altamami et al.¹² (2019), in which recovery was successful, with the patients evolving from grade VI preoperatively to grades lower than IV on the HB scale after masseteric-facial nerve anastomosis. However, Vincent et al.¹³ (2019) found no statistically significant difference in their study, but suggest that masseteric-to-facial nerve transfer with selective neurectomy provides a significant improvement in the ability to smile, with a long-term decrease in synkinesis among patients unresponsive to botulinum toxin therapy or who, for other reasons, do not wish to continue with regular injections.

In addition, Biglioli et al.¹⁴ (2018) presented a new surgical technique for the treatment of patients with segmental midface paresis (with or without synkinesis). This method comprises two surgeries with a gap of one year between them, which consist of a masseteric neuroorrhaphy combined with cross-face nerve grafting. Of the 20 patients in the study, 8 presented muscular contracture at rest in the middle third of the face preoperatively, and, postoperatively, partial-to-complete relaxation of the middle face was observed in all of these cases. In the preoperative period, synkinesis was present in 11 of the 20 patients, and, after the surgery, it was completely resolved in 7 patients; in 3 patients, synkinesis was reduced, although it was still present in response to more intense stimuli than in the preoperative period; and, in 1 out of 11 patients, synkinesis

remained present and showed no improvement. Moreover, in another study, Biglioli et al.¹⁵ (2018), analyzed 24 patients with dense unilateral facial palsy (HB grade VI) who underwent surgery between January 2013 and February 2016. The surgery employed was a new technique with triple neural inputs: the use of the masseteric nerve and 30% of the hypoglossus nerve fibers as quantitative sources was associated with the contralateral facial nerve (incorporated via two cross-face nerve grafts) as a qualitative source. In all patients, the facial paralysis improved, as did their grades on the HB scale. Recovery of the facial paralysis showed its first signs two to eight months after surgery, and continued through the entire postoperative period. During an 18-month follow-up period, the median grade of those 24 patients was III on the modified HB scale: 11 patients (45.83%) were grade II; 11 patients (45.83%), grade III; and 2 (8.33%), grade IV. Both studies showed that other surgical techniques can be used for the treatment of facial paralysis; however, masseteric-facial anastomosis is still being performed as an important step in the recovery of these patients. To compare two different surgical techniques, the cross-facial nerve grafting (CFNG) and the masseteric-facial nerve anastomosis, mainly used in the aforementioned papers, Hontanilla et al.¹⁶ (2018) divided the 28 patients in their sample into 2 groups: 10 in group 1 (patients submitted to CFNG) and 18 in group 2 (patients submitted to masseteric-facial nerve anastomosis). In total, 80% of the CFNG group had spontaneous recovery of their smiles during more than 50% of their daily lives. In contrast, in group 2, the recovery was of just 55.5%. Nevertheless, satisfaction with the treatment was greater among group 2. This paper also showed that, with the masseteric-facial nerve anastomosis, better symmetry can be achieved, with a greater recovery and a higher degree of satisfaction on the part of the patients.

Sakthivel et al.⁸ (2020) described the surgical technique and the effectiveness of the masseteric nerve in early facial reanimation. All patients in their sample underwent end-to-

end anastomosis, except for one, in whom a greater auricular interposition graft was used. In all cases, the facial muscles showed signs of recovery in 2 to 5 months, when movement of the cheek musculature was noticed when the patients activated their masseter muscle. Eye movement started appearing after 6 to 9 months (in 3 cases), and forehead movements, after 18 months (in 1 case). According to the modified HB scale, one patient had grade-I function, two patients had grade-II function, and three had grade-V function. There was no morbidity, except for one patient who underwent interposition graft and had numbness in the ear lobule. None of the patients could feel the loss of function of the masseteric nerve. Additionally, in the study by Gray et al.¹⁷ (2020), 8 patients with eye closure or smile excursion synkinesis were submitted to the surgery, with results that made the authors realize that the single-stage masseteric-zygomatic nerve transfer is a useful procedure for patients with synkinesis caused by facial nerve paralysis.

Conclusion

Due to the obstruction of the innervation of the facial muscles by tumor growth, masseteric-facial nerve anastomosis, after the removal of the tumor, had to be performed for the total reinnervation of such muscles and the skin in the cases herein reported. The masseteric nerve is the one that shows the best prognosis among all the cranial nerve that could be used, due to its efficient neuroplasticity, and the rapid and effective motor rehabilitation of the facial muscles. However, this fact alone does not lead to a better prognosis when compared with the other cases in the literature, it is also necessary to perform the surgical technique well to access the facial branch, and consequently perform a better masseteric-facial anastomosis.

Conflict of Interest

The authors have no conflict of interests to declare.

References

- 1 Toulgoat F, Sarrazin JL, Benoudiba F, et al. Facial nerve: from anatomy to pathology. *Diagn Interv Imaging* 2013;94(10):1033–1042. Doi: 10.1016/j.diii.2013.06.016
- 2 Sanders RD, Gillig PM. The trigeminal (V) and facial (VII) cranial nerves: Head and face sensation and movement. *Psychiatry (Edmont Pa)* 2010;7(01):13–16
- 3 Batista KT, Cauhi AF. Surgical rehabilitation for facial paralysis. *Rev Bras Cir Plást* 2007;22(04):253–260
- 4 House JW, Brackmann DE. Facial nerve grading system. *Otolaryngol Head Neck Surg* 1985;93(02):146–147. Doi: 10.1177/019459988509300202
- 5 Davies J, Al-Hassani F, Kannan R. Facial nerve disorder: a review of the literature. *Int J Surg Oncol (NY)* 2018; 3(07):e65. Doi: 10.1097/IJ9.0000000000000065
- 6 Jeziorowski A, Tirapelle R, Saciloto A, et al. Masseteric-facial anastomosis on treatment of facial palsy. *ACM Arq Catarin Med* 2007;36(supl. 1):165–168
- 7 Xu ZJ, Chen LS, Zhan JD, et al. [Modified rhytidectomy incision and modified Blair incision contrast research in superficial parotid gland tumor resection]. *Lin Chung Er Bi Yan Hou Tou Jing Wai Ke Za Zhi* 2017; 31(21):1684–1687. Doi: 10.13201/j.issn.1001-1781.2017.21.015
- 8 Sakthivel P, Singh CA, Thakur A, Thirumeni G, Raveendran S, Sharma SC. Masseteric-Facial Nerve Anastomosis: Surgical Techniques and Outcomes-A Pilot Indian study. *Indian J Otolaryngol Head Neck Surg* 2020;72(01):92–97. Doi: 10.1007/s12070-019-01758-z
- 9 Wu H, Zhang L, Han D, et al. Summary and consensus in 7th International Conference on acoustic neuroma: An update for the management of sporadic acoustic neuromas. *World J Otorhinolaryngol Head Neck Surg* 2016;2(04):234–239. Doi: 10.1016/j.wjorl.2016.10.002
- 10 Lee YS, Ahn JH, Park HJ, et al. Dual Coaptation of Facial Nerve Using Masseteric Branch of Trigeminal Nerve for Iatrogenic Facial Palsy: Preliminary Reports. *Ann Otol Rhinol Laryngol* 2020;129(05): 505–511. Doi: 10.1177/0003489419893722
- 11 Bianchi B, Varazzani A, Pedrazzi G, et al. Masseteric coaptation and crossfacial nerve grafting: Is it still applicable 22 months after the onset of facial palsy? *Microsurgery* 2018;38(08):860–866. Doi: 10.1002/micr.30296
- 12 Altamami NM, Zaouche S, Vertu-Ciolino D. A comparative retrospective study: hypoglossofacial versus masseterofacial nerve anastomosis using Sunnybrook facial grading system. *Eur Arch Otorhinolaryngol* 2019;276(01):209–216. Doi: 10.1007/s00405-018-5186-y
- 13 Vincent AG, Bevans SE, Robitschek JM, Wind GG, Hohman MH. Masseteric-to-Facial Nerve Transfer and Selective Neurectomy for Rehabilitation of the Synkinetic Smile. *JAMA Facial Plast Surg* 2019;21(06):504–510. Doi: 10.1001/jamafacial.2019.0689
- 14 Biglioli F, Soliman M, El-Shazly M, et al. Use of the masseteric nerve to treat segmental midface paresis. *Br J Oral Maxillofac Surg* 2018;56(08):719–726. Doi: 10.1016/j.bjoms.2018.07.023
- 15 Biglioli F, Allevi F, Rabbiosi D, et al. Triple innervation for re-animation of recent facial paralysis. *J Craniomaxillofac Surg* 2018; 46(05):851–857. Doi: 10.1016/j.jcms.2018.02.014
- 16 Hontanilla B, Olivás J, Cabello Á, Marré D. Cross-Face Nerve Grafting versus Masseteric-to-Facial Nerve Transposition for Re-animation of Incomplete Facial Paralysis: A Comparative Study Using the FACIAL CLIMA Evaluating System. *Plast Reconstr Surg* 2018;142(02):179e–191e. Doi: 10.1097/PRS.0000000000004612
- 17 Gray ML, Hu S, Gorbea E, Mashkevich G. Masseteric-zygomatic nerve transfer for the management of eye closure-smile excursion synkinesis. [published online ahead of print, 2020 Apr 4] *Am J Otolaryngol* 2020;41(04):102479. Doi: 10.1016/j.amjoto.2020.102479